

Rules for integrands of the form $(a + b \sin[c + d x])^n$

1. $\int (b \sin[c + d x])^n dx$

1. $\int (b \sin[c + d x])^n dx$ when $2 n \in \mathbb{Z}$

1. $\int (b \sin[c + d x])^n dx$ when $n > 1$

1: $\int \sin[c + d x]^n dx$ when $\frac{n-1}{2} \in \mathbb{Z}^+$

Derivation: Integration by substitution

Basis: If $\frac{n-1}{2} \in \mathbb{Z}$, then $\sin[c + d x]^n = -\frac{1}{d} \text{Subst}\left[\left(1 - x^2\right)^{\frac{n-1}{2}}, x, \cos[c + d x]\right] \partial_x \cos[c + d x]$

Rule: If $\frac{n-1}{2} \in \mathbb{Z}^+$, then

$$\int \sin[c + d x]^n dx \rightarrow -\frac{1}{d} \text{Subst}\left[\int (1 - x^2)^{\frac{n-1}{2}} dx, x, \cos[c + d x]\right]$$

Program code:

```
Int[sin[c_.+d_.*x_]^n_,x_Symbol]:= 
-1/d*Subst[Int[Expand[(1-x^2)^((n-1)/2),x],x,x,Cos[c+d*x]] /; 
FreeQ[{c,d},x] && IGtQ[(n-1)/2,0]
```

2. $\int (b \sin[c + d x])^n dx$ when $n > 1$

1: $\int \sin[c + d x]^2 dx$

- Derivation: Algebraic expansion

- Basis: $\sin[z]^2 = \frac{1}{2} - \frac{\cos[2z]}{2}$

- Rule:

$$\int \sin[c + d x]^2 dx \rightarrow \frac{x}{2} - \frac{\sin[2c + 2d x]}{4d}$$

Program code:

```
Int[sin[c_.+d_.*x_/2]^2,x_Symbol] :=  
  x/2 - Sin[2*c+d*x]/(2*d) /;  
FreeQ[{c,d},x]
```

2: $\int (b \sin[c + d x])^n dx$ when $n > 1$

Reference: G&R 2.510.2 with $q \rightarrow 0$, CRC 299

Reference: G&R 2.510.5 with $p \rightarrow 0$, CRC 305

- Derivation: Sine recurrence 3a with $A \rightarrow 0$, $B \rightarrow a$, $C \rightarrow b$, $m \rightarrow m - 1$, $n \rightarrow -1$

- Derivation: Sine recurrence 1b with $A \rightarrow 0$, $B \rightarrow 0$, $C \rightarrow b$, $a \rightarrow 0$, $m \rightarrow -1$, $n \rightarrow n - 1$

- Rule: If $n > 1$, then

$$\int (b \sin[c+dx])^n dx \rightarrow -\frac{b \cos[c+dx] (b \sin[c+dx])^{n-1}}{d n} + \frac{b^2 (n-1)}{n} \int (b \sin[c+dx])^{n-2} dx$$

Program code:

```
Int[(b.*sin[c.+d.*x_])^n,x_Symbol] :=
(* -Cot[c+d*x]*(c*Sin[c+d*x])^n/(d*n) + b^2*(n-1)/n*Int[(b*Sin[c+d*x])^(n-2),x] *)
-b*Cos[c+d*x]*(b*Sin[c+d*x])^(n-1)/(d*n) + b^2*(n-1)/n*Int[(b*Sin[c+d*x])^(n-2),x] /;
FreeQ[{b,c,d},x] && GtQ[n,1] && IntegerQ[2*n]
```

2: $\int (b \sin[c+dx])^n dx$ when $n < -1$

Reference: G&R 2.510.3 with $q \rightarrow 0$, CRC 309

Reference: G&R 2.510.6 with $p \rightarrow 0$, CRC 313

Reference: G&R 2.552.3

Derivation: Sine recurrence 3a with $A \rightarrow 0$, $B \rightarrow a$, $C \rightarrow b$, $m \rightarrow m - 1$, $n \rightarrow -1$ inverted

Derivation: Sine recurrence 2a with $A \rightarrow 1$, $B \rightarrow 0$, $C \rightarrow 0$, $a \rightarrow 0$, $m \rightarrow 0$

Rule: If $n < -1$, then

$$\int (b \sin[c+dx])^n dx \rightarrow \frac{\cos[c+dx] (b \sin[c+dx])^{n+1}}{b d (n+1)} + \frac{n+2}{b^2 (n+1)} \int (b \sin[c+dx])^{n+2} dx$$

Program code:

```
Int[(b.*sin[c.+d.*x_])^n,x_Symbol] :=
Cos[c+d*x]*(b*Sin[c+d*x])^(n+1)/(b*d*(n+1)) +
(n+2)/(b^2*(n+1))*Int[(b*Sin[c+d*x])^(n+2),x] /;
FreeQ[{b,c,d},x] && LtQ[n,-1] && IntegerQ[2*n]
```

3. $\int (b \sin[c + d x])^n dx$ when $-1 \leq n \leq -1$

1. $\int \sin[c + d x]^n dx$ when $-1 \leq n \leq -1$

1. $\int \sin[c + d x]^n dx$ when $n^2 = 1$

1: $\int \sin[c + d x] dx$

Reference: G&R 2.01.5, CRC 290, A&S 4.3.113

Reference: G&R 2.01.6, CRC 291, A&S 4.3.114

Derivation: Primitive rule

Basis: $\partial_x \cos[c + d x] = -d \sin[c + d x]$

Rule:

$$\int \sin[c + d x] dx \rightarrow -\frac{\cos[c + d x]}{d}$$

Program code:

```
Int[sin[c_.+Pi/2+d_.*x_],x_Symbol]:=  
  Sin[c+d*x]/d /;  
FreeQ[{c,d},x]
```

```
Int[sin[c_.+d_.*x_],x_Symbol]:=  
  -Cos[c+d*x]/d /;  
FreeQ[{c,d},x]
```

$$\text{x: } \int \frac{1}{\sin[c + d x]} dx$$

— Note: This rule not necessary since *Mathematica* automatically simplifies $\frac{1}{\sin[z]}$ to $\csc[z]$.

— Rule:

$$\int \frac{1}{\sin[c + d x]} dx \rightarrow \int \csc[c + d x] dx$$

— Program code:

```
(* Int[1/sin[c_.+d_.*x_],x_Symbol] :=
  Int[Csc[c+d*x],x] /;
  FreeQ[{c,d},x] *)
```

2. $\int \sin[c + d x]^n dx$ when $n^2 = \frac{1}{4}$

1: $\int \sqrt{\sin[c + d x]} dx$

Derivation: Primitive rule

- Basis: $\partial_x \text{EllipticE}\left[\frac{1}{2} \left(x - \frac{\pi}{2}\right), 2\right] = \frac{\sqrt{\sin[x]}}{2}$

- Rule:

$$\int \sqrt{\sin[c + d x]} dx \rightarrow \frac{2}{d} \text{EllipticE}\left[\frac{1}{2} \left(c - \frac{\pi}{2} + d x\right), 2\right]$$

- Program code:

```
Int[Sqrt[sin[c_.+d_.*x_]],x_Symbol]:=  
 2/d*EllipticE[1/2*(c-Pi/2+d*x),2];;  
FreeQ[{c,d},x]
```

$$\text{2: } \int \frac{1}{\sqrt{\sin[c + d x]}} dx$$

Derivation: Primitive rule

Basis: $\partial_x \text{EllipticF}\left[\frac{1}{2} \left(x - \frac{\pi}{2}\right), 2\right] = \frac{1}{2 \sqrt{\sin[x]}}$

Rule:

$$\int \frac{1}{\sqrt{\sin[c + d x]}} dx \rightarrow \frac{2}{d} \text{EllipticF}\left[\frac{1}{2} \left(c - \frac{\pi}{2} + d x\right), 2\right]$$

Program code:

```
Int[1/Sqrt[sin[c_.+d_.*x_]],x_Symbol]:=  
 2/d*EllipticF[1/2*(c-Pi/2+d*x),2]/;  
FreeQ[{c,d},x]
```

2: $\int (b \sin[c + d x])^n dx$ when $-1 < n < -1$

Derivation: Piecewise constant extraction

Basis: $\partial_x \frac{(b \sin[c + d x])^n}{\sin[c + d x]^n} = 0$

Rule: If $-1 < n < -1$, then

$$\int (b \sin[c + d x])^n dx \rightarrow \frac{(b \sin[c + d x])^n}{\sin[c + d x]^n} \int \sin[c + d x]^n dx$$

Program code:

```
Int[(b_*sin[c_+d_*x_])^n_,x_Symbol] :=
  (b*Sin[c+d*x])^n/Sin[c+d*x]^n*Int[Sin[c+d*x]^n,x] /;
FreeQ[{b,c,d},x] && LtQ[-1,n,1] && IntegerQ[2*n]
```

2: $\int (b \sin[c + d x])^n dx$ when $2n \notin \mathbb{Z}$

Derivation: Piecewise constant extraction and integration by substitution

Basis: $\partial_x \frac{\cos[c + d x]}{\sqrt{\cos[c + d x]^2}} = 0$

Basis: $\frac{\cos[c + d x]}{\sqrt{\cos[c + d x]^2}} \frac{\cos[c + d x]}{\sqrt{1 - \sin[c + d x]^2}} = 1$

Basis: $\cos[c + d x] F[b \sin[c + d x]] = \frac{1}{b d} \text{Subst}[F[x], x, b \sin[c + d x]] \partial_x (b \sin[c + d x])$

Rule: If $2n \notin \mathbb{Z}$, then

$$\begin{aligned}
\int (b \sin[c + d x])^n dx &\rightarrow \frac{\cos[c + d x]}{\sqrt{\cos[c + d x]^2}} \int \frac{\cos[c + d x] (b \sin[c + d x])^n}{\sqrt{1 - \sin[c + d x]^2}} dx \\
&\rightarrow \frac{\cos[c + d x]}{b d \sqrt{\cos[c + d x]^2}} \text{Subst} \left[\int \frac{x^n}{\sqrt{1 - \frac{x^2}{b^2}}} dx, x, b \sin[c + d x] \right] \\
&\rightarrow \frac{\cos[c + d x] (b \sin[c + d x])^{n+1}}{b d (n+1) \sqrt{\cos[c + d x]^2}} \text{Hypergeometric2F1} \left[\frac{1}{2}, \frac{n+1}{2}, \frac{n+3}{2}, \sin[c + d x]^2 \right]
\end{aligned}$$

– Alternate rule: If $2 n \notin \mathbb{Z}$, then

$$\int (b \sin[c + d x])^n dx \rightarrow -\frac{\cos[c + d x] (b \sin[c + d x])^{n+1}}{b d (\sin[c + d x]^2)^{\frac{n+1}{2}}} \text{Hypergeometric2F1} \left[\frac{1}{2}, \frac{1-n}{2}, \frac{3}{2}, \cos[c + d x]^2 \right]$$

Program code:

```

(* Int[(b.*sin[c.+d.*x_])^n_,x_Symbol] :=
  Cos[c+d*x]/(b*d*Sqrt[Cos[c+d*x]^2])*Subst[Int[x^n/Sqrt[1-x^2/b^2],x],x,b*Sin[c+d*x]] /;
  FreeQ[{b,c,d,n},x] && Not[IntegerQ[2*n] || IntegerQ[3*n]] *)

Int[(b.*sin[c._+d._*x_])^n_,x_Symbol] :=
  Cos[c+d*x]*(b*Sin[c+d*x])^(n+1)/(b*d*(n+1)*Sqrt[Cos[c+d*x]^2])*Hypergeometric2F1[1/2,(n+1)/2,(n+3)/2,Sin[c+d*x]^2] /;
  FreeQ[{b,c,d,n},x] && Not[IntegerQ[2*n]]

```

2: $\int (a + b \sin[c + d x])^2 dx$

– Derivation: Algebraic expansion

– Basis: $(a + b z)^2 = \frac{1}{2} (2 a^2 + b^2) + 2 a b z - \frac{1}{2} b^2 (1 - 2 z^2)$

– Rule:

$$\int (a + b \sin[c + d x])^2 dx \rightarrow \frac{(2 a^2 + b^2) x}{2} - \frac{2 a b \cos[c + d x]}{d} - \frac{b^2 \cos[c + d x] \sin[c + d x]}{2 d}$$

Program code:

```
Int[(a_+b_.*sin[c_.+d_.*x_])^2,x_Symbol]:=  
  (2*a^2+b^2)*x/2 - 2*a*b*Cos[c+d*x]/d - b^2*Cos[c+d*x]*Sin[c+d*x]/(2*d) /;  
FreeQ[{a,b,c,d},x]
```

3. $\int (a + b \sin[c + d x])^n dx$ when $a^2 - b^2 = 0$

1. $\int (a + b \sin[c + d x])^n dx$ when $a^2 - b^2 = 0 \wedge 2 n \in \mathbb{Z}$

1. $\int (a + b \sin[c + d x])^n dx$ when $a^2 - b^2 = 0 \wedge 2 n \in \mathbb{Z}^+$

1: $\int (a + b \sin[c + d x])^n dx$ when $a^2 - b^2 = 0 \wedge n \in \mathbb{Z}^+$

Derivation: Algebraic expansion

Rule: If $a^2 - b^2 = 0 \wedge n \in \mathbb{Z}^+$, then

$$\int (a + b \sin[c + d x])^n dx \rightarrow \int \text{ExpandTrig}[(a + b \sin[c + d x])^n, x] dx$$

Program code:

```
Int[(a_+b_.*sin[c_.+d_.*x_])^n_,x_Symbol]:=  
  Int[ExpandTrig[(a+b*sin[c+d*x])^n,x],x] /;  
FreeQ[{a,b,c,d,n},x] && EqQ[a^2-b^2,0] && IGtQ[n,0]
```

2. $\int (a + b \sin[c + dx])^n dx$ when $a^2 - b^2 = 0 \wedge n + \frac{1}{2} \in \mathbb{Z}^+$

1: $\int \sqrt{a + b \sin[c + dx]} dx$ when $a^2 - b^2 = 0$

Derivation: Singly degenerate sine recurrence 1b with $A \rightarrow c$, $B \rightarrow d$, $m \rightarrow \frac{1}{2}$, $n \rightarrow -1$, $p \rightarrow 0$

Rule: If $a^2 - b^2 = 0$, then

$$\int \sqrt{a + b \sin[c + dx]} dx \rightarrow -\frac{2b \cos[c + dx]}{d \sqrt{a + b \sin[c + dx]}}$$

Program code:

```
Int[Sqrt[a+b.*sin[c.+d.*x_]],x_Symbol]:=  
-2*b*Cos[c+d*x]/(d*Sqrt[a+b*Sin[c+d*x]]) /;  
FreeQ[{a,b,c,d},x] && EqQ[a^2-b^2,0]
```

2: $\int (a + b \sin(c + dx))^n dx$ when $a^2 - b^2 = 0 \wedge n - \frac{1}{2} \in \mathbb{Z}^+$

Reference: G&R 2.555.? inverted

Derivation: Singly degenerate sine recurrence 1b with $A \rightarrow c$, $B \rightarrow d$, $n \rightarrow -1$, $p \rightarrow 0$

Rule: If $a^2 - b^2 = 0 \wedge n - \frac{1}{2} \in \mathbb{Z}^+$, then

$$\int (a + b \sin(c + dx))^n dx \rightarrow -\frac{b \cos(c + dx) (a + b \sin(c + dx))^{n-1}}{d n} + \frac{a (2n - 1)}{n} \int (a + b \sin(c + dx))^{n-1} dx$$

Program code:

```
Int[(a_+b_.*sin[c_.+d_.*x_])^n_,x_Symbol]:=  
-b*Cos[c+d*x]*(a+b*Sin[c+d*x])^(n-1)/(d*n)+  
a*(2*n-1)/n*Int[(a+b*Sin[c+d*x])^(n-1),x]/;  
FreeQ[{a,b,c,d},x] && EqQ[a^2-b^2,0] && IGtQ[n-1/2,0]
```

2. $\int (a + b \sin[c + dx])^n dx$ when $a^2 - b^2 = 0 \wedge 2n \in \mathbb{Z}^+$

1: $\int \frac{1}{a + b \sin[c + dx]} dx$ when $a^2 - b^2 = 0$

Reference: G&R 2.555.3', CRC 337', A&S 4.3.134'/5'

- Derivation: Singly degenerate sine recurrence 2a with $A \rightarrow 1$, $B \rightarrow 0$, $m \rightarrow -1$, $n \rightarrow 0$, $p \rightarrow 0$

- Rule: If $a^2 - b^2 = 0$, then

$$\int \frac{1}{a + b \sin[c + dx]} dx \rightarrow -\frac{\cos[c + dx]}{d(b + a \sin[c + dx])}$$

- Program code:

```
Int[1/(a+b.*sin[c.+d.*x_]),x_Symbol] :=
-Cos[c+d*x]/(d*(b+a*Sin[c+d*x])) /;
FreeQ[{a,b,c,d},x] && EqQ[a^2-b^2,0]
```

2: $\int \frac{1}{\sqrt{a+b \sin[c+dx]}} dx$ when $a^2 - b^2 = 0$

Derivation: Integration by substitution

Basis: If $a^2 - b^2 = 0$, then $\frac{1}{\sqrt{a+b \sin[c+dx]}} = -\frac{2}{d} \text{Subst}\left[\frac{1}{2a-x^2}, x, \frac{b \cos[c+dx]}{\sqrt{a+b \sin[c+dx]}}\right] \partial_x \frac{b \cos[c+dx]}{\sqrt{a+b \sin[c+dx]}}$

Rule: If $a^2 - b^2 = 0$, then

$$\int \frac{1}{\sqrt{a+b \sin[c+dx]}} dx \rightarrow -\frac{2}{d} \text{Subst}\left[\int \frac{1}{2a-x^2} dx, x, \frac{b \cos[c+dx]}{\sqrt{a+b \sin[c+dx]}}\right]$$

Program code:

```
Int[1/Sqrt[a_+b_.*sin[c_._+d_._*x_]],x_Symbol]:= 
-2/d*Subst[Int[1/(2*a-x^2),x],x,b*Cos[c+d*x]/Sqrt[a+b*Sin[c+d*x]]]/;
FreeQ[{a,b,c,d},x] && EqQ[a^2-b^2,0]
```

3: $\int (a + b \sin[c + d x])^n dx$ when $a^2 - b^2 = 0 \wedge n < -1 \wedge 2n \in \mathbb{Z}$

Reference: G&R 2.555.?

Derivation: Singly degenerate sine recurrence 2a with $A \rightarrow 1$, $B \rightarrow 0$, $n \rightarrow 0$, $p \rightarrow 0$

Rule: If $a^2 - b^2 = 0 \wedge n < -1 \wedge 2n \in \mathbb{Z}$, then

$$\int (a + b \sin[c + d x])^n dx \rightarrow \frac{b \cos[c + d x] (a + b \sin[c + d x])^n}{a d (2n+1)} + \frac{n+1}{a (2n+1)} \int (a + b \sin[c + d x])^{n+1} dx$$

Program code:

```
Int[ (a+b.*sin[c.+d.*x_])^n_,x_Symbol] :=
  b*Cos[c+d*x]* (a+b*Sin[c+d*x])^n/(a*d*(2*n+1)) +
  (n+1)/(a*(2*n+1))*Int[ (a+b*Sin[c+d*x])^(n+1),x] /;
FreeQ[{a,b,c,d},x] && EqQ[a^2-b^2,0] && LtQ[n,-1] && IntegerQ[2*n]
```

2. $\int (a + b \sin[c + d x])^n dx$ when $a^2 - b^2 = 0 \wedge 2n \notin \mathbb{Z}$

x: $\int (a + b \sin[c + d x])^n dx$ when $a^2 - b^2 = 0 \wedge 2n \notin \mathbb{Z}$

Derivation: Piecewise constant extraction and integration by substitution

Basis: If $a^2 - b^2 = 0$, then $\partial_x \frac{\cos[c+d x]}{\sqrt{a-b \sin[c+d x]}} = 0$

Basis: If $a^2 - b^2 = 0$, then $\frac{a^2 \cos[c+d x]}{\sqrt{a+b \sin[c+d x]}} \frac{\cos[c+d x]}{\sqrt{a-b \sin[c+d x]}} = 1$

Basis: $\cos[c + d x] F[\sin[c + d x]] = \frac{1}{d} \text{Subst}[F[x], x, \sin[c + d x]] \partial_x \sin[c + d x]$

Note: If $3n \in \mathbb{Z}$, this results in a complicated expression involving elliptic integrals instead of a single hypergeometric

function.

- Rule: If $a^2 - b^2 = 0 \wedge 2n \notin \mathbb{Z}$, then

$$\int (a + b \sin[c + d x])^n dx \rightarrow$$

$$\frac{a^2 \cos[c + d x]}{\sqrt{a + b \sin[c + d x]} \sqrt{a - b \sin[c + d x]}} \int \frac{\cos[c + d x] (a + b \sin[c + d x])^{n-\frac{1}{2}}}{\sqrt{a - b \sin[c + d x]}} dx \rightarrow$$

$$\frac{a^2 \cos[c + d x]}{d \sqrt{a + b \sin[c + d x]} \sqrt{a - b \sin[c + d x]}} \text{Subst} \left[\int \frac{(a + b x)^{\frac{n-1}{2}}}{\sqrt{a - b x}} dx, x, \sin[c + d x] \right]$$

- Program code:

```
(* Int[(a+b.*sin[c.+d.*x_])^n_,x_Symbol]:=  
a^2*Cos[c+d*x]/(d*.Sqrt[a+b*Sin[c+d*x]]*Sqrt[a-b*Sin[c+d*x]])*Subst[Int[(a+b*x)^(n-1/2)/Sqrt[a-b*x],x],x,Sin[c+d*x]] /;  
FreeQ[{a,b,c,d,n},x] && EqQ[a^2-b^2,0] && Not[IntegerQ[2*n]] *)
```

1: $\int (a + b \sin[c + dx])^n dx$ when $a^2 - b^2 = 0 \wedge 2n \notin \mathbb{Z} \wedge a > 0$

Derivation: Piecewise constant extraction and integration by substitution

Rule: If $a^2 - b^2 = 0 \wedge 2n \notin \mathbb{Z} \wedge a > 0$, then

$$\int (a + b \sin[c + dx])^n dx \rightarrow$$

$$-\frac{2^{n+\frac{1}{2}} a^{n-\frac{1}{2}} b \cos[c + dx]}{d \sqrt{a + b \sin[c + dx]}} \text{Hypergeometric2F1}\left[\frac{1}{2}, \frac{1}{2} - n, \frac{3}{2}, \frac{1}{2} \left(1 - \frac{b \sin[c + dx]}{a}\right)\right]$$

Program code:

```
Int[(a_+b_.*sin[c_.+d_.*x_])^n_,x_Symbol]:=  
-2^(n+1/2)*a^(n-1/2)*b*Cos[c+d*x]/(d*Sqrt[a+b*Sin[c+d*x]])*Hypergeometric2F1[1/2,1/2-n,3/2,1/2*(1-b*Sin[c+d*x]/a)] /;  
FreeQ[{a,b,c,d,n},x] && EqQ[a^2-b^2,0] && Not[IntegerQ[2*n]] && GtQ[a,0]
```

2: $\int (a + b \sin(c + d x))^n dx$ when $a^2 - b^2 = 0 \wedge 2n \notin \mathbb{Z} \wedge a > 0$

Derivation: Piecewise constant extraction

Basis: $\partial_x \frac{(a+b \sin(c+d x))^n}{\left(1+\frac{b}{a} \sin(c+d x)\right)^n} = 0$

Rule: If $a^2 - b^2 = 0 \wedge 2n \notin \mathbb{Z} \wedge a > 0$, then

$$\int (a + b \sin(c + d x))^n dx \rightarrow \frac{a^{\text{IntPart}[n]} (a + b \sin(c + d x))^{\text{FracPart}[n]}}{\left(1 + \frac{b}{a} \sin(c + d x)\right)^{\text{FracPart}[n]}} \int \left(1 + \frac{b}{a} \sin(c + d x)\right)^n dx$$

Program code:

```
Int[(a+b.*sin[c.+d.*x_])^n_,x_Symbol] :=
  a^IntPart[n]* (a+b*Sin[c+d*x])^FracPart[n]/(1+b/a*Sin[c+d*x])^FracPart[n]*Int[(1+b/a*Sin[c+d*x])^n,x] /;
FreeQ[{a,b,c,d,n},x] && EqQ[a^2-b^2,0] && Not[IntegerQ[2*n]] && Not[GtQ[a,0]]
```

4. $\int (a + b \sin[c + dx])^n dx$ when $a^2 - b^2 \neq 0$

1. $\int (a + b \sin[c + dx])^n dx$ when $a^2 - b^2 \neq 0 \wedge 2n \in \mathbb{Z}$

1. $\int (a + b \sin[c + dx])^n dx$ when $a^2 - b^2 \neq 0 \wedge 2n \in \mathbb{Z}^+$

1. $\int \sqrt{a + b \sin[c + dx]} dx$ when $a^2 - b^2 \neq 0$

1: $\int \sqrt{a + b \sin[c + dx]} dx$ when $a^2 - b^2 \neq 0 \wedge a + b > 0$

Derivation: Primitive rule

Basis: If $a + b > 0$, then $\partial_x \text{EllipticE}\left[\frac{1}{2} \left(x - \frac{\pi}{2}\right), \frac{2b}{a+b}\right] = \frac{1}{2\sqrt{a+b}} \sqrt{a + b \sin[x]}$

Rule: If $a^2 - b^2 \neq 0 \wedge a + b > 0$, then

$$\int \sqrt{a + b \sin[c + dx]} dx \rightarrow \frac{2\sqrt{a+b}}{d} \text{EllipticE}\left[\frac{1}{2} \left(c - \frac{\pi}{2} + dx\right), \frac{2b}{a+b}\right]$$

— Program code:

```
Int[Sqrt[a+b.*sin[c.+d.*x.]],x_Symbol] :=
  2*Sqrt[a+b]/d*EllipticE[1/2*(c-Pi/2+d*x),2*b/(a+b)] /;
FreeQ[{a,b,c,d},x] && NeQ[a^2-b^2,0] && GtQ[a+b,0]
```

2: $\int \sqrt{a + b \sin[c + dx]} \, dx$ when $a^2 - b^2 \neq 0 \wedge a - b > 0$

Derivation: Primitive rule

Basis: If $a - b > 0$, then $\partial_x \text{EllipticE}\left[\frac{1}{2} \left(x + \frac{\pi}{2}\right), -\frac{2b}{a-b}\right] = \frac{1}{2\sqrt{a-b}} \sqrt{a + b \sin[x]}$

Rule: If $a^2 - b^2 \neq 0 \wedge a - b > 0$, then

$$\int \sqrt{a + b \sin[c + dx]} \, dx \rightarrow \frac{2\sqrt{a-b}}{d} \text{EllipticE}\left[\frac{1}{2} \left(c + \frac{\pi}{2} + dx\right), -\frac{2b}{a-b}\right]$$

Program code:

```
Int[Sqrt[a+b.*sin[c.+d.*x_]],x_Symbol]:=  
 2*Sqrt[a-b]/d*EllipticE[1/2*(c+Pi/2+d*x),-2*b/(a-b)] /;  
 FreeQ[{a,b,c,d},x] && NeQ[a^2-b^2,0] && GtQ[a-b,0]
```

3: $\int \sqrt{a + b \sin[c + dx]} \, dx$ when $a^2 - b^2 \neq 0 \wedge a + b \neq 0$

Derivation: Piecewise constant extraction

Basis: $\partial_x \frac{\sqrt{a+b f[x]}}{\sqrt{\frac{a+b f[x]}{a+b}}} = 0$

Note: Since $\frac{a}{a+b} + \frac{b}{a+b} = 1 > 0$, the above rule applies to the resulting integrand.

Rule: If $a^2 - b^2 \neq 0 \wedge a + b \neq 0$, then

$$\int \sqrt{a + b \sin[c + dx]} \, dx \rightarrow \frac{\sqrt{a + b \sin[c + dx]}}{\sqrt{\frac{a+b \sin[c+dx]}{a+b}}} \int \sqrt{\frac{a}{a+b} + \frac{b}{a+b} \sin[c + dx]} \, dx$$

Program code:

```
Int[Sqrt[a_+b_.*sin[c_.+d_.*x_]],x_Symbol] :=
  Sqrt[a+b*Sin[c+d*x]]/Sqrt[(a+b*Sin[c+d*x])/(a+b)]*Int[Sqrt[a/(a+b)+b/(a+b)*Sin[c+d*x]],x] /;
FreeQ[{a,b,c,d},x] && NeQ[a^2-b^2,0] && Not[GtQ[a+b,0]]
```

2: $\int (a + b \sin[c + dx])^n \, dx$ when $a^2 - b^2 \neq 0 \wedge n > 1 \wedge 2n \in \mathbb{Z}$

Derivation: Nondegenerate sine recurrence 1b with $A \rightarrow a c$, $B \rightarrow b c + a d$, $C \rightarrow b d$, $m \rightarrow -1 + m$, $n \rightarrow -1$, $p \rightarrow 0$

Rule: If $a^2 - b^2 \neq 0 \wedge n > 1 \wedge 2n \in \mathbb{Z}$, then

$$\int (a + b \sin[c + dx])^n \, dx \rightarrow -\frac{b \cos[c + dx] (a + b \sin[c + dx])^{n-1}}{d n} + \frac{1}{n} \int (a + b \sin[c + dx])^{n-2} (a^2 n + b^2 (n - 1) + a b (2 n - 1) \sin[c + dx]) \, dx$$

Program code:

```
Int[(a_+b_.*sin[c_.+d_.*x_])^n_,x_Symbol] :=
  -b*Cos[c+d*x]*(a+b*Sin[c+d*x])^(n-1)/(d*n) +
  1/n*Int[(a+b*Sin[c+d*x])^(n-2)*Simp[a^2*n+b^2*(n-1)+a*b*(2*n-1)*Sin[c+d*x],x],x] /;
FreeQ[{a,b,c,d},x] && NeQ[a^2-b^2,0] && GtQ[n,1] && IntegerQ[2*n]
```

2. $\int (a + b \sin[c + dx])^n dx$ when $a^2 - b^2 \neq 0 \wedge 2n \in \mathbb{Z}^-$

1. $\int \frac{1}{a + b \sin[c + dx]} dx$ when $a^2 - b^2 \neq 0$

1. $\int \frac{1}{a + b \sin[c + dx]} dx$ when $a^2 - b^2 > 0$

1: $\int \frac{1}{a + b \sin[c + dx]} dx$ when $a^2 - b^2 > 0 \wedge a > 0$

Note: Resulting antiderivative is continuous on the real line.

Rule: If $a^2 - b^2 > 0 \wedge a > 0$, let $q = \sqrt{a^2 - b^2}$, then

$$\int \frac{1}{a + b \sin[c + dx]} dx \rightarrow \frac{x}{q} + \frac{2}{dq} \operatorname{ArcTan}\left[\frac{b \cos[c + dx]}{a + q + b \sin[c + dx]}\right]$$

Program code:

```
Int[1/(a_+b_.*sin[c_.+d_.*x_]),x_Symbol]:=  
With[{q=Rt[a^2-b^2,2]},  
x/q+2/(d*q)*ArcTan[b*Cos[c+d*x]/(a+q+b*Sin[c+d*x])]];  
FreeQ[{a,b,c,d},x] && GtQ[a^2-b^2,0] && PosQ[a]
```

2: $\int \frac{1}{a + b \sin[c + dx]} dx$ when $a^2 - b^2 > 0 \wedge a \neq 0$

Note: Resulting antiderivative is continuous on the real line.

Rule: If $a^2 - b^2 > 0 \wedge a \neq 0$, let $q = \sqrt{a^2 - b^2}$, then

$$\int \frac{1}{a + b \sin[c + dx]} dx \rightarrow -\frac{x}{q} - \frac{2}{dq} \operatorname{ArcTan}\left[\frac{b \cos[c + dx]}{a - q + b \sin[c + dx]}\right]$$

Program code:

```
Int[1/(a+b.*sin[c.+d.*x_]),x_Symbol] :=
With[{q=Rt[a^2-b^2,2]},
-x/q - 2/(d*q)*ArcTan[b*Cos[c+d*x]/(a-q+b*Sin[c+d*x])] ];
FreeQ[{a,b,c,d},x] && GtQ[a^2-b^2,0] && NegQ[a]
```

2: $\int \frac{1}{a + b \sin[c + dx]} dx$ when $a^2 - b^2 \neq 0$

Reference: G&R 2.551.3, CRC 340, A&S 4.3.131

Reference: G&R 2.553.3, CRC 341, A&S 4.3.133

Derivation: Integration by substitution

Basis:

$$F[\sin[c+dx], \cos[c+dx]] = \frac{2}{d} \text{Subst}\left[\frac{1}{1+x^2} F\left[\frac{2x}{1+x^2}, \frac{1-x^2}{1+x^2}\right], x, \tan\left[\frac{1}{2}(c+dx)\right]\right] \partial_x \tan\left[\frac{1}{2}(c+dx)\right]$$

- Basis: $\frac{1}{a+b \sin[c+dx]} = \frac{2}{d} \text{Subst}\left[\frac{1}{a+2bx+ax^2}, x, \tan\left[\frac{1}{2}(c+dx)\right]\right] \partial_x \tan\left[\frac{1}{2}(c+dx)\right]$

- Basis: $\frac{1}{a+b \cos[c+dx]} = \frac{2}{d} \text{Subst}\left[\frac{1}{a+b+(a-b)x^2}, x, \tan\left[\frac{1}{2}(c+dx)\right]\right] \partial_x \tan\left[\frac{1}{2}(c+dx)\right]$

Note: $\tan\left[\frac{z}{2}\right] = \frac{\sin[z]}{1+\cos[z]}$

Rule: If $a^2 - b^2 \neq 0$, then

$$\begin{aligned} \int \frac{1}{a + b \sin[c + dx]} dx &\rightarrow \frac{2}{d} \text{Subst}\left[\int \frac{1}{a + 2bx + ax^2} dx, x, \tan\left[\frac{1}{2}(c+dx)\right]\right] \\ \int \frac{1}{a + b \cos[c + dx]} dx &\rightarrow \frac{2}{d} \text{Subst}\left[\int \frac{1}{a + b + (a-b)x^2} dx, x, \tan\left[\frac{1}{2}(c+dx)\right]\right] \end{aligned}$$

Program code:

```
Int[1/(a+b.*sin[c.+Pi/2+d.*x_]),x_Symbol]:=  
With[{e=FreeFactors[Tan[(c+d*x)/2],x]},  
2*e/d*Subst[Int[1/(a+b+(a-b)*e^2*x^2),x],x,Tan[(c+d*x)/2]/e]] /;  
FreeQ[{a,b,c,d},x] && NeQ[a^2-b^2,0]
```

```

Int[1/(a_+b_.*sin[c_.+d_.*x_]),x_Symbol] :=
With[{e=FreeFactors[Tan[(c+d*x)/2],x]}, 
2*e/d*Subst[Int[1/(a+2*b*e*x+a*e^2*x^2),x],x,Tan[(c+d*x)/2]/e]] /;
FreeQ[{a,b,c,d},x] && NeQ[a^2-b^2,0]

```

2. $\int \frac{1}{\sqrt{a + b \sin[c + d x]}} dx$ when $a^2 - b^2 \neq 0$

1: $\int \frac{1}{\sqrt{a + b \sin[c + d x]}} dx$ when $a^2 - b^2 \neq 0 \wedge a + b > 0$

Derivation: Primitive rule

Basis: If $a + b > 0$, then $\partial_x \text{EllipticF}\left[\frac{1}{2} \left(x - \frac{\pi}{2}\right), \frac{2b}{a+b}\right] = \frac{\sqrt{a+b}}{2 \sqrt{a+b \sin[x]}}$

Rule: If $a^2 - b^2 \neq 0 \wedge a + b > 0$, then

$$\int \frac{1}{\sqrt{a + b \sin[c + d x]}} dx \rightarrow \frac{2}{d \sqrt{a+b}} \text{EllipticF}\left[\frac{1}{2} \left(c - \frac{\pi}{2} + d x\right), \frac{2b}{a+b}\right]$$

Program code:

```

Int[1/Sqrt[a_+b_.*sin[c_.+d_.*x_]],x_Symbol] :=
2/(d*Sqrt[a+b])*EllipticF[1/2*(c-Pi/2+d*x),2*b/(a+b)] /;
FreeQ[{a,b,c,d},x] && NeQ[a^2-b^2,0] && GtQ[a+b,0]

```

2: $\int \frac{1}{\sqrt{a + b \sin[c + d x]}} dx$ when $a^2 - b^2 \neq 0 \wedge a - b > 0$

Derivation: Primitive rule

Basis: If $a - b > 0$, then $\partial_x \text{EllipticF}\left[\frac{1}{2} \left(x + \frac{\pi}{2}\right), -\frac{2b}{a-b}\right] = \frac{\sqrt{a-b}}{2 \sqrt{a+b \sin[x]}}$

Rule: If $a^2 - b^2 \neq 0 \wedge a - b > 0$, then

$$\int \frac{1}{\sqrt{a + b \sin[c + d x]}} dx \rightarrow \frac{2}{d \sqrt{a-b}} \text{EllipticF}\left[\frac{1}{2} \left(c + \frac{\pi}{2} + d x\right), -\frac{2b}{a-b}\right]$$

Program code:

```
Int[1/Sqrt[a+b.*sin[c.+d.*x_]],x_Symbol]:=  
 2/(d*Sqrt[a-b])*EllipticF[1/2*(c+Pi/2+d*x),-2*b/(a-b)] /;  
 FreeQ[{a,b,c,d},x] && NeQ[a^2-b^2,0] && GtQ[a-b,0]
```

3: $\int \frac{1}{\sqrt{a + b \sin[c + d x]}} dx$ when $a^2 - b^2 \neq 0 \wedge a + b \neq 0$

Derivation: Piecewise constant extraction

Basis: $\partial_x \frac{\sqrt{\frac{a+b f[x]}{a+b}}}{\sqrt{a+b f[x]}} = 0$

Note: Since $\frac{a}{a+b} + \frac{b}{a+b} = 1 > 0$, rule f1 applies to the resulting integrand.

Rule: If $a^2 - b^2 \neq 0 \wedge a + b \neq 0$, then

$$\int \frac{1}{\sqrt{a + b \sin[c + dx]}} dx \rightarrow \frac{\sqrt{\frac{a+b \sin[c+dx]}{a+b}}}{\sqrt{a + b \sin[c + dx]}} \int \frac{1}{\sqrt{\frac{a}{a+b} + \frac{b}{a+b} \sin[c + dx]}} dx$$

Program code:

```
Int[1/Sqrt[a+b.*sin[c.+d.*x.]],x_Symbol] :=
  Sqrt[(a+b*Sin[c+d*x.])/(a+b)]/Sqrt[a+b*Sin[c+d*x.]]*Int[1/Sqrt[a/(a+b)+b/(a+b)*Sin[c+d*x.]],x] /;
FreeQ[{a,b,c,d},x] && NeQ[a^2-b^2,0] && Not[GtQ[a+b,0]]
```

3: $\int (a + b \sin[c + dx])^n dx$ when $a^2 - b^2 \neq 0 \wedge n < -1 \wedge 2n \in \mathbb{Z}$

Reference: G&R 2.552.3

Derivation: Nondegenerate sine recurrence 1a with $A \rightarrow 1$, $B \rightarrow 0$, $C \rightarrow 0$, $m \rightarrow 0$, $p \rightarrow 0$

Rule: If $a^2 - b^2 \neq 0 \wedge n < -1 \wedge 2n \in \mathbb{Z}$, then

$$\int (a + b \sin[c + dx])^n dx \rightarrow -\frac{b \cos[c + dx] (a + b \sin[c + dx])^{n+1}}{d (n + 1) (a^2 - b^2)} + \frac{1}{(n + 1) (a^2 - b^2)} \int (a + b \sin[c + dx])^{n+1} (a (n + 1) - b (n + 2) \sin[c + dx]) dx$$

Program code:

```
Int[(a+b.*sin[c.+d.*x.])^n_,x_Symbol] :=
  -b*Cos[c+d*x.]* (a+b*Sin[c+d*x.])^(n+1)/(d*(n+1)*(a^2-b^2)) +
  1/((n+1)*(a^2-b^2))*Int[(a+b*Sin[c+d*x.])^(n+1)*Simp[a*(n+1)-b*(n+2)*Sin[c+d*x.],x],x] /;
FreeQ[{a,b,c,d},x] && NeQ[a^2-b^2,0] && LtQ[n,-1] && IntegerQ[2*n]
```

2: $\int (a + b \sin[c + d x])^n dx$ when $a^2 - b^2 \neq 0 \wedge 2 n \notin \mathbb{Z}$

Derivation: Piecewise constant extraction and integration by substitution

Basis: $\partial_x \frac{\cos[c+d x]}{\sqrt{1+\sin[c+d x]}} \sqrt{1-\sin[c+d x]} = 0$

Basis: $\cos[c + d x] = \frac{1}{d} \partial_x \sin[c + d x]$

Rule: If $a^2 - b^2 \neq 0 \wedge 2 n \notin \mathbb{Z}$, then

$$\begin{aligned} & \int (a + b \sin[c + d x])^n dx \rightarrow \\ & \frac{\cos[c + d x]}{\sqrt{1 + \sin[c + d x]}} \sqrt{1 - \sin[c + d x]} \int \frac{\cos[c + d x] (a + b \sin[c + d x])^n}{\sqrt{1 + \sin[c + d x]} \sqrt{1 - \sin[c + d x]}} dx \rightarrow \\ & \frac{\cos[c + d x]}{d \sqrt{1 + \sin[c + d x]} \sqrt{1 - \sin[c + d x]}} \text{Subst} \left[\int \frac{(a + b x)^n}{\sqrt{1+x} \sqrt{1-x}} dx, x, \sin[c + d x] \right] \end{aligned}$$

Program code:

```
Int[(a+b.*sin[c.+d.*x.])^n_,x_Symbol] :=
  Cos[c+d*x]/(d*.Sqrt[1+Sin[c+d*x]]*Sqrt[1-Sin[c+d*x]])*Subst[Int[(a+b*x)^n/(Sqrt[1+x]*Sqrt[1-x]),x],x,Sin[c+d*x]] /;
FreeQ[{a,b,c,d,n},x] && NeQ[a^2-b^2,0] && Not[IntegerQ[2*n]]
```

Rules for integrands of the form $(a + b \sin[c + d x] \cos[c + d x])^n$

1: $\int (a + b \sin[c + d x] \cos[c + d x])^n dx$

Derivation: Algebraic simplification

Basis: $\sin[z] \cos[z] = \frac{1}{2} \sin[2z]$

Rule:

$$\int (a + b \sin[c + d x] \cos[c + d x])^n dx \rightarrow \int \left(a + \frac{1}{2} b \sin[2c + 2dx] \right)^n dx$$

Program code:

```
Int[(a_+b_.*sin[c_._+d_._*x_]*cos[c_._+d_._*x_])^n_,x_Symbol]:=  
  Int[(a+b*Sin[2*c+2*d*x]/2)^n,x] /;  
  FreeQ[{a,b,c,d,n},x]
```