

Rules for integrands of the form $(a + b \tan[c + d x^n])^p$

1: $\int (a + b \tan[c + d x^n])^p dx$ when $\frac{1}{n} \in \mathbb{Z}^+ \wedge p \in \mathbb{Z}$

Derivation: Integration by substitution

Basis: If $-1 \leq n \leq 1 \wedge n \neq 0$, then $F[x^n] = \frac{1}{n} \text{Subst}\left[x^{\frac{1}{n}-1} F[x], x, x^n\right] \partial_x x^n$

Note: If $\frac{1}{n} \in \mathbb{Z}^-$, resulting integrand is not integrable.

Rule: If $\frac{1}{n} \in \mathbb{Z}^+ \wedge p \in \mathbb{Z}$, then

$$\int (a + b \tan[c + d x^n])^p dx \rightarrow \frac{1}{n} \text{Subst}\left[\int x^{\frac{1}{n}-1} (a + b \tan[c + d x])^p dx, x, x^n\right]$$

Program code:

```
Int[(a_.+b_.*Tan[c_._+d_._*x_._^n_])^p_.,x_Symbol]:=  
 1/n*Subst[Int[x^(1/n-1)*(a+b*Tan[c+d*x])^p,x],x,x^n] /;  
 FreeQ[{a,b,c,d,p},x] && IGtQ[1/n,0] && IntegerQ[p]
```

```
Int[(a_.+b_.*Cot[c_._+d_._*x_._^n_])^p_.,x_Symbol]:=  
 1/n*Subst[Int[x^(1/n-1)*(a+b*Cot[c+d*x])^p,x],x,x^n] /;  
 FreeQ[{a,b,c,d,p},x] && IGtQ[1/n,0] && IntegerQ[p]
```

x: $\int (a + b \tan[c + d x^n])^p dx$

— Rule:

$$\int (a + b \tan[c + d x^n])^p dx \rightarrow \int (a + b \tan[c + d x^n])^p dx$$

— Program code:

```
Int[(a_..+b_..*Tan[c_..+d_..*x_..^n_..])^p_.,x_Symbol] :=
  Unintegrable[(a+b*Tan[c+d*x^n])^p,x] /;
  FreeQ[{a,b,c,d,n,p},x]

Int[(a_..+b_..*Cot[c_..+d_..*x_..^n_..])^p_.,x_Symbol] :=
  Unintegrable[(a+b*Cot[c+d*x^n])^p,x] /;
  FreeQ[{a,b,c,d,n,p},x]
```

s: $\int (a + b \tan[c + d u^n])^p dx$ when $u = e + f x$

Derivation: Integration by substitution

Rule: If $u = e + f x$, then

$$\int (a + b \tan[c + d u^n])^p dx \rightarrow \frac{1}{f} \text{Subst}\left[\int (a + b \tan[c + d x^n])^p dx, x, u\right]$$

Program code:

```
Int[(a_..+b_..*Tan[c_..+d_..*u_..^n_..])^p_.,x_Symbol] :=
  1/Coefficient[u,x,1]*Subst[Int[(a+b*Tan[c+d*x^n])^p,x],x,u] /;
  FreeQ[{a,b,c,d,n,p},x] && LinearQ[u,x] && NeQ[u,x]
```

```
Int[(a_.+b_.*Cot[c_.+d_.*u_`^n_`])^p_,x_Symbol]:=  
 1/Coefficient[u,x,1]*Subst[Int[(a+b*Cot[c+d*x^n])^p,x],x,u] /;  
FreeQ[{a,b,c,d,n,p},x] && LinearQ[u,x] && NeQ[u,x]
```

N: $\int (a + b \tan u)^p dx$ when $u = c + d x^n$

Derivation: Algebraic normalization

– Rule: If $u = c + d x^n$, then

$$\int (a + b \tan u)^p dx \rightarrow \int (a + b \tan [c + d x^n])^p dx$$

Program code:

```
Int[(a_.+b_.*Tan[u_])^p_,x_Symbol]:=  
  Int[(a+b*Tan[ExpandToSum[u,x]])^p,x] /;  
FreeQ[{a,b,p},x] && BinomialQ[u,x] && Not[BinomialMatchQ[u,x]]
```

```
Int[(a_.+b_.*Cot[u_])^p_,x_Symbol]:=  
  Int[(a+b*Cot[ExpandToSum[u,x]])^p,x] /;  
FreeQ[{a,b,p},x] && BinomialQ[u,x] && Not[BinomialMatchQ[u,x]]
```

Rules for integrands of the form $(e x)^m (a + b \tan[c + d x^n])^p$

1. $\int x^m (a + b \tan[c + d x^n])^p dx$

1: $\int x^m (a + b \tan[c + d x^n])^p dx$ when $\frac{m+1}{n} \in \mathbb{Z}^+ \wedge p \in \mathbb{Z}$

Derivation: Integration by substitution

Basis: If $\frac{m+1}{n} \in \mathbb{Z}$, then $x^m F[x^n] = \frac{1}{n} \text{Subst}[x^{\frac{m+1}{n}-1} F[x], x, x^n] \partial_x x^n$

Note: If $\frac{m+1}{n} \in \mathbb{Z}^-$, resulting integrand is not integrable.

Rule: If $\frac{m+1}{n} \in \mathbb{Z}^+ \wedge p \in \mathbb{Z}$, then

$$\int x^m (a + b \tan[c + d x^n])^p dx \rightarrow \frac{1}{n} \text{Subst}\left[\int x^{\frac{m+1}{n}-1} (a + b \tan[c + d x])^p dx, x, x^n\right]$$

Program code:

```
Int[x^m_.*(a_._+b_._*Tan[c_._+d_._*x_^n_])^p_.,x_Symbol]:=  
  1/n*Subst[Int[x^(Simplify[(m+1)/n]-1)*(a+b*Tan[c+d*x])^p,x],x,x^n] /;  
  FreeQ[{a,b,c,d,m,n,p},x] && IGtQ[Simplify[(m+1)/n],0] && IntegerQ[p]
```

```
Int[x^m_.*(a_._+b_._*Cot[c_._+d_._*x_^n_])^p_.,x_Symbol]:=  
  1/n*Subst[Int[x^(Simplify[(m+1)/n]-1)*(a+b*Cot[c+d*x])^p,x],x,x^n] /;  
  FreeQ[{a,b,c,d,m,n,p},x] && IGtQ[Simplify[(m+1)/n],0] && IntegerQ[p]
```

2: $\int x^m \tan[c + d x^n]^2 dx$

Note: Although this rule reduces the degree of the tangent factor, the resulting integral is not integrable unless $\frac{m+1}{n} \in \mathbb{Z}^+$.

Rule:

$$\int x^m \tan[c + d x^n]^2 dx \rightarrow \frac{x^{m-n+1} \tan[c + d x^n]}{d n} - \int x^m dx - \frac{m - n + 1}{d n} \int x^{m-n} \tan[c + d x^n] dx$$

Program code:

```
Int[x^m.*Tan[c._+d._*x^n_]^2,x_Symbol] :=
  x^(m-n+1)*Tan[c+d*x^n]/(d*n) - Int[x^m,x] - (m-n+1)/(d*n)*Int[x^(m-n)*Tan[c+d*x^n],x] /;
FreeQ[{c,d,m,n},x]
```

```
Int[x^m.*Cot[c._+d._*x^n_]^2,x_Symbol] :=
  -x^(m-n+1)*Cot[c+d*x^n]/(d*n) - Int[x^m,x] + (m-n+1)/(d*n)*Int[x^(m-n)*Cot[c+d*x^n],x] /;
FreeQ[{c,d,m,n},x]
```

x. $\int x^m \tan[a + b x^n]^p dx$ when $0 < n < m + 1$

1: $\int x^m \tan[a + b x^n]^p dx$ when $0 < n < m + 1 \wedge p > 1$

Note: Although this rule reduces the degree of the tangent factor, the resulting integrals are not integrable unless $\frac{m+1}{n} \in \mathbb{Z}^+ \wedge p \in \mathbb{Z}$.

Rule: If $0 < n < m + 1 \wedge p > 1$, then

$$\int x^m \tan[a + b x^n]^p dx \rightarrow \frac{x^{m-n+1} \tan[a + b x^n]^{p-1}}{b n (p - 1)} - \frac{m - n + 1}{b n (p - 1)} \int x^{m-n} \tan[a + b x^n]^{p-1} dx - \int x^m \tan[a + b x^n]^{p-2} dx$$

Program code:

```
(* Int[x^m.*Tan[a._+b._*x^n_]^p_,x_Symbol] :=
  x^(m-n+1)*Tan[a+b*x^n]^(p-1)/(b*n*(p-1)) -
  (m-n+1)/(b*n*(p-1))*Int[x^(m-n)*Tan[a+b*x^n]^(p-1),x] -
  Int[x^m*Tan[a+b*x^n]^(p-2),x] /;
FreeQ[{a,b},x] && LtQ[0,n,m+1] && GtQ[p,1] *)
```

```
(* Int[x^m.*Cot[a_.+b_.*x^n_]^p_,x_Symbol] :=  
-x^(m-n+1)*Cot[a+b*x^n]^p/(b*n*(p-1)) +  
(m-n+1)/(b*n*(p-1))*Int[x^(m-n)*Cot[a+b*x^n]^p,x] -  
Int[x^m*Cot[a+b*x^n]^p,x] /;  
FreeQ[{a,b},x] && LtQ[0,n,m+1] && GtQ[p,1] *)
```

2: $\int x^m \tan[a + b x^n]^p dx$ when $0 < n < m + 1 \wedge p < -1$

Note: Although this rule reduces the degree of the tangent factor, the resulting integrals are not integrable unless $\frac{m+1}{n} \in \mathbb{Z}^+ \wedge p \in \mathbb{Z}$.

Rule: If $0 < n < m + 1 \wedge p < -1$, then

$$\int x^m \tan[a + b x^n]^p dx \rightarrow \frac{x^{m-n+1} \tan[a + b x^n]^{p+1}}{b n (p+1)} - \frac{m - n + 1}{b n (p+1)} \int x^{m-n} \tan[a + b x^n]^{p+1} dx - \int x^m \tan[a + b x^n]^{p+2} dx$$

Program code:

```
(* Int[x^m.*Tan[a_.+b_.*x^n_]^p_,x_Symbol] :=  
x^(m-n+1)*Tan[a+b*x^n]^(p+1)/(b*n*(p+1)) -  
(m-n+1)/(b*n*(p+1))*Int[x^(m-n)*Tan[a+b*x^n]^(p+1),x] -  
Int[x^m*Tan[a+b*x^n]^(p+2),x] /;  
FreeQ[{a,b},x] && LtQ[0,n,m+1] && LtQ[p,-1] *)
```

```
(* Int[x^m.*Cot[a_.+b_.*x^n_]^p_,x_Symbol] :=  
-x^(m-n+1)*Cot[a+b*x^n]^(p+1)/(b*n*(p+1)) +  
(m-n+1)/(b*n*(p+1))*Int[x^(m-n)*Cot[a+b*x^n]^(p+1),x] -  
Int[x^m*Cot[a+b*x^n]^(p+2),x] /;  
FreeQ[{a,b},x] && LtQ[0,n,m+1] && LtQ[p,-1] *)
```

X: $\int x^m (a + b \tan[c + d x^n])^p dx$

— Rule:

$$\int x^m (a + b \tan[c + d x^n])^p dx \rightarrow \int x^m (a + b \tan[c + d x^n])^p dx$$

— Program code:

```
Int[x^m.*(a.+b.*Tan[c.+d.*x^n])^p.,x_Symbol] :=
  Unintegrable[x^m*(a+b*Tan[c+d*x^n])^p,x] /;
FreeQ[{a,b,c,d,m,n,p},x]
```

```
Int[x^m.*(a.+b.*Cot[c.+d.*x^n])^p.,x_Symbol] :=
  Unintegrable[x^m*(a+b*Cot[c+d*x^n])^p,x] /;
FreeQ[{a,b,c,d,m,n,p},x]
```

2: $\int (e x)^m (a + b \tan[c + d x^n])^p dx$

Derivation: Piecewise constant extraction

Basis: $a_x \frac{(e x)^m}{x^m} = 0$

Rule:

$$\int (e x)^m (a + b \tan[c + d x^n])^p dx \rightarrow \frac{e^{\text{IntPart}[m]} (e x)^{\text{FracPart}[m]}}{x^{\text{FracPart}[m]}} \int x^m (a + b \tan[c + d x^n])^p dx$$

Program code:

```
Int[(e_*x_)^m_.*(a_._+b_._*Tan[c_._+d_._*x_._^n_])^p_.,x_Symbol]:=  
  e^IntPart[m]* (e*x)^FracPart[m]/x^FracPart[m]*Int[x^m*(a+b*Tan[c+d*x^n])^p,x] /;  
FreeQ[{a,b,c,d,e,m,n,p},x]
```

```
Int[(e_*x_)^m_.*(a_._+b_._*Cot[c_._+d_._*x_._^n_])^p_.,x_Symbol]:=  
  e^IntPart[m]* (e*x)^FracPart[m]/x^FracPart[m]*Int[x^m*(a+b*Cot[c+d*x^n])^p,x] /;  
FreeQ[{a,b,c,d,e,m,n,p},x]
```

N: $\int (e x)^m (a + b \tan[u])^p dx$ when $u = c + d x^n$

Derivation: Algebraic normalization

– Rule: If $u = c + d x^n$, then

$$\int (e x)^m (a + b \tan[u])^p dx \rightarrow \int (e x)^m (a + b \tan[c + d x^n])^p dx$$

– Program code:

```
Int[(e_*x_)^m_.*(a_.+b_.*Tan[u_])^p_,x_Symbol] :=
  Int[(e*x)^m*(a+b*Tan[ExpandToSum[u,x]])^p,x] /;
  FreeQ[{a,b,e,m,p},x] && BinomialQ[u,x] && Not[BinomialMatchQ[u,x]]
```

```
Int[(e_*x_)^m_.*(a_.+b_.*Cot[u_])^p_,x_Symbol] :=
  Int[(e*x)^m*(a+b*Cot[ExpandToSum[u,x]])^p,x] /;
  FreeQ[{a,b,e,m,p},x] && BinomialQ[u,x] && Not[BinomialMatchQ[u,x]]
```

Rules for integrands of the form $x^m \sec[a + b x^n]^p \tan[a + b x^n]$

1: $\int x^m \sec[a + b x^n]^p \tan[a + b x^n] dx$ when $n \in \mathbb{Z} \wedge m - n \geq 0$

Derivation: Integration by parts

Note: Dummy exponent $q = 1$ required in program code so InputForm of integrand is recognized.

Rule: If $n \in \mathbb{Z} \wedge m - n \geq 0$, then

$$\int x^m \sec[a + b x^n]^p \tan[a + b x^n] dx \rightarrow \frac{x^{m-n+1} \sec[a + b x^n]^p}{b n p} - \frac{m - n + 1}{b n p} \int x^{m-n} \sec[a + b x^n]^p dx$$

Program code:

```
Int[x^m.*Sec[a.+b.*x^n.]^p.*Tan[a.+b.*x^n.]^q.,x_Symbol] :=
  x^(m-n+1)*Sec[a+b*x^n]^p/(b*n*p) -
  (m-n+1)/(b*n*p)*Int[x^(m-n)*Sec[a+b*x^n]^p,x] /;
FreeQ[{a,b,p},x] && IntegerQ[n] && GeQ[m,n] && EqQ[q,1]
```

```
Int[x^m.*Csc[a.+b.*x^n.]^p.*Cot[a.+b.*x^n.]^q.,x_Symbol] :=
  -x^(m-n+1)*Csc[a+b*x^n]^p/(b*n*p) +
  (m-n+1)/(b*n*p)*Int[x^(m-n)*Csc[a+b*x^n]^p,x] /;
FreeQ[{a,b,p},x] && IntegerQ[n] && GeQ[m,n] && EqQ[q,1]
```