

## Rules for integrands of the form $(c x)^m P_q[x] (a + b x^2)^p$

1:  $\int x^m P_q[x^2] (a + b x^2)^p dx \text{ when } \frac{m-1}{2} \in \mathbb{Z}$

Derivation: Integration by substitution

Basis: If  $\frac{m-1}{2} \in \mathbb{Z}$ , then  $x^m F[x^2] = \frac{1}{2} \text{Subst}[x^{\frac{m-1}{2}} F[x], x, x^2] \partial_x x^2$

Rule 1.1.2.y.1: If  $\frac{m-1}{2} \in \mathbb{Z}$ , then

$$\int x^m P_q[x^2] (a + b x^2)^p dx \rightarrow \frac{1}{2} \text{Subst}\left[\int x^{\frac{m-1}{2}} P_q[x] (a + b x)^p dx, x, x^2\right]$$

Program code:

```
Int[x_^m_.*Pq_*(a_+b_.*x_^2)^p_.,x_Symbol] :=  
  1/2*Subst[Int[x^((m-1)/2)*SubstFor[x^2,Pq,x]*(a+b*x)^p,x],x,x^2] /;  
 FreeQ[{a,b,p},x] && PolyQ[Pq,x^2] && IntegerQ[(m-1)/2]
```

2:  $\int (c x)^m P_q[x] (a + b x^2)^p dx \text{ when } P_q[x, 0] = 0$

Derivation: Algebraic simplification

Rule 1.1.2.y.2: If  $P_q[x, 0] = 0$ , then

$$\int (c x)^m P_q[x] (a + b x^2)^p dx \rightarrow \frac{1}{c} \int (c x)^{m+1} \text{PolynomialQuotient}[P_q[x], x, x] (a + b x^2)^p dx$$

Program code:

```
Int[(c_.*x_)^m_.*Pq_*(a_+b_.*x_^2)^p_.,x_Symbol] :=  
  1/c*Int[(c*x)^(m+1)*PolynomialQuotient[Pq,x,x]*(a+b*x^2)^p,x] /;  
 FreeQ[{a,b,c,m,p},x] && PolyQ[Pq,x] && EqQ[coeff[Pq,x,0],0]
```

3:  $\int (c x)^m (a + b x^2)^p (f + h x^2) dx$  when  $a h (m + 1) - b f (m + 2 p + 3) = 0 \wedge m \neq -1$

Derivation: Special case of one step of the Ostrogradskiy-Hermite integration method

– Rule 1.1.2.y.3: If  $a h (m + 1) - b f (m + 2 p + 3) = 0 \wedge m \neq -1$ , then

$$\int (c x)^m (a + b x^2)^p (f + h x^2) dx \rightarrow \frac{f (c x)^{m+1} (a + b x^2)^{p+1}}{a c (m + 1)}$$

Program code:

```
Int[(c_.*x_)^m_.*P2_*(a_+b_.*x_^2)^p_.,x_Symbol]:=  
With[{f=Coeff[P2,x,0],g=Coeff[P2,x,1],h=Coeff[P2,x,2]},  
h*(c*x)^(m+1)*(a+b*x^2)^(p+1)/(b*c*(m+2*p+3)) /;  
EqQ[g,0] && EqQ[a*h*(m+1)-b*f*(m+2*p+3),0]] /;  
FreeQ[{a,b,c,m,p},x] && PolyQ[P2,x,2] && NeQ[m,-1]
```

4:  $\int (c x)^m P_q[x] (a + b x^2)^p dx$  when  $p + 2 \in \mathbb{Z}^+$

Derivation: Algebraic expansion

– Rule 1.1.2.y.4: If  $p + 2 \in \mathbb{Z}^+$ , then

$$\int (c x)^m P_q[x] (a + b x^2)^p dx \rightarrow \int \text{ExpandIntegrand}[(c x)^m P_q[x] (a + b x^2)^p, x] dx$$

Program code:

```
Int[(c_.*x_)^m_.*Pq_*(a_+b_.*x_^2)^p_.,x_Symbol]:=  
Int[ExpandIntegrand[(c*x)^m*Pq*(a+b*x^2)^p,x],x] /;  
FreeQ[{a,b,c,m},x] && PolyQ[Pq,x] && IGtQ[p,-2]
```

5:  $\int x^m P_q[x^2] (a + b x^2)^p dx$  when  $\frac{m}{2} \in \mathbb{Z} \wedge \frac{m+1}{2} + p \in \mathbb{Z}^- \wedge m + 2 q + 2 p + 1 < 0$

Derivation: Algebraic expansion and binomial recurrence 3b

Basis:  $\int x^m (a + b x^2)^p dx = \frac{x^{m+1} (a + b x^2)^{p+1}}{a (m+1)} - \frac{b (m+2 (p+1)+1)}{a (m+1)} \int x^{m+2} (a + b x^2)^p dx$

Note: Interestingly this rule eliminates the constant term of  $P_q[x^2]$  rather than the highest degree term.

Rule 1.1.2.y.5: If  $\frac{m}{2} \in \mathbb{Z} \wedge \frac{m+1}{2} + p \in \mathbb{Z}^- \wedge m + 2 q + 2 p + 1 < 0$ , let  $A \rightarrow P_q[x^2, 0]$  and

$Q_{q-1}[x^2] \rightarrow \text{PolynomialQuotient}[P_q[x^2] - A, x^2, x]$ , then

$$\int x^m P_q[x^2] (a + b x^2)^p dx \rightarrow$$

$$A \int x^m (a + b x^2)^p dx + \int x^{m+2} Q_{q-1}[x^2] (a + b x^2)^p dx \rightarrow$$

$$\frac{A x^{m+1} (a + b x^2)^{p+1}}{a (m+1)} + \frac{1}{a (m+1)} \int x^{m+2} (a + b x^2)^p (a (m+1) Q_{q-1}[x^2] - A b (m+2 (p+1)+1)) dx$$

Program code:

```
Int[x^m_*Pq_*(a+b_*x^2)^p_,x_Symbol] :=
With[{A=Coeff[Pq,x,0],Q=PolynomialQuotient[Pq-Coeff[Pq,x,0],x^2,x]}, 
A*x^(m+1)*(a+b*x^2)^(p+1)/(a*(m+1)) + 1/(a*(m+1))*Int[x^(m+2)*(a+b*x^2)^p*(a*(m+1)*Q-A*b*(m+2*(p+1)+1)),x] ];
FreeQ[{a,b},x] && PolyQ[Pq,x^2] && IntegerQ[m/2] && ILtQ[(m+1)/2+p,0] && LtQ[m+Expon[Pq,x]+2*p+1,0]
```

6.  $\int (c x)^m Pq[x] (a + b x^2)^p dx \text{ when } p < -1$

1:  $\int (c x)^m Pq[x] (a + b x^2)^p dx \text{ when } p < -1 \wedge m > 0$

Derivation: Algebraic expansion and quadratic recurrence 2a

Rule 1.1.2.y.6.1: If  $p < -1 \wedge m > 0$ ,

let  $Q_{q-2}[x] \rightarrow \text{PolynomialQuotient}[Pq[x], a+b x^2, x]$  and  $f + g x \rightarrow \text{PolynomialRemainder}[Pq[x], a+b x^2, x]$ , then

$$\int (c x)^m Pq[x] (a + b x^2)^p dx \rightarrow$$

$$\int (c x)^m (f + g x) (a + b x^2)^p dx + \int (c x)^{m-1} (c x) Q_{q-2}[x] (a + b x^2)^{p+1} dx \rightarrow$$

$$\frac{(c x)^m (a + b x^2)^{p+1} (a g - b f x)}{2 a b (p + 1)} + \frac{c}{2 a b (p + 1)} \int (c x)^{m-1} (a + b x^2)^{p+1} (2 a b (p + 1) x Q_{q-2}[x] - a g m + b f (m + 2 p + 3) x) dx$$

Program code:

```
Int[(c_.*x_)^m.*Pq_*(a_+b_.*x_^2)^p_,x_Symbol]:=  
With[{Q=PolynomialQuotient[Pq,a+b*x^2,x],  
f=Coeff[PolynomialRemainder[Pq,a+b*x^2,x],x,0],  
g=Coeff[PolynomialRemainder[Pq,a+b*x^2,x],x,1]},  
(c*x)^m*(a+b*x^2)^(p+1)*(a*g-b*f*x)/(2*a*b*(p+1)) +  
c/(2*a*b*(p+1))*Int[(c*x)^(m-1)*(a+b*x^2)^(p+1)*ExpandToSum[2*a*b*(p+1)*x*Q-a*g*m+b*f*(m+2*p+3)*x,x],x]];  
FreeQ[{a,b,c},x] && PolyQ[Pq,x] && LtQ[p,-1] && GtQ[m,0]
```

2.  $\int (c x)^m P_q[x] (a + b x^2)^p dx$  when  $p < -1 \wedge m \geq 0$

1:  $\int (c x)^m P_q[x] (a + b x^2)^p dx$  when  $p < -1 \wedge m \in \mathbb{Z}^-$

## Derivation: Algebraic expansion and trinomial recurrence 2b

Rule 1.1.2.y.6.2.1: If  $p < -1 \wedge m \in \mathbb{Z}^-$ ,

let  $Q_{m+q-2}[x] \rightarrow \text{PolynomialQuotient}[(c x)^m P_q[x], a + b x^2, x]$  and  
 $f + g x \rightarrow \text{PolynomialRemainder}[(c x)^m P_q[x], a + b x^2, x]$ , then

$$\int (c x)^m P_q[x] (a + b x^2)^p dx \rightarrow$$

$$\int (f + g x) (a + b x^2)^p dx + \int Q_{m+q-2}[x] (a + b x^2)^{p+1} dx \rightarrow$$

$$\frac{(a g - b f x) (a + b x^2)^{p+1}}{2 a b (p + 1)} + \frac{1}{2 a (p + 1)} \int (c x)^m (a + b x^2)^{p+1} (2 a (p + 1) (c x)^{-m} Q_{m+q-2}[x] + f (2 p + 3) (c x)^{-m}) dx$$

## Program code:

```
Int[(c_.*x_)^m_.*Pq_*(a_+b_.*x_^2)^p_,x_Symbol]:=  
With[{Q=PolynomialQuotient[(c*x)^m*Pq,a+b*x^2,x],  
f=Coeff[PolynomialRemainder[(c*x)^m*Pq,a+b*x^2,x],x,0],  
g=Coeff[PolynomialRemainder[(c*x)^m*Pq,a+b*x^2,x],x,1]},  
(a*g-b*f*x)*(a+b*x^2)^(p+1)/(2*a*b*(p+1)) +  
1/(2*a*(p+1))*Int[(c*x)^m*(a+b*x^2)^(p+1)*ExpandToSum[2*a*(p+1)*(c*x)^(-m)*Q+f*(2*p+3)*(c*x)^(-m),x],x]] /;  
FreeQ[{a,b,c},x] && PolyQ[Pq,x] && LtQ[p,-1] && LtQ[m,0]
```

2:  $\int (c x)^m Pq[x] (a + b x^2)^p dx \text{ when } p < -1 \wedge m \geq 0$

### Derivation: Algebraic expansion and quadratic recurrence 2b

Rule 1.1.2.y.6.2.2: If  $p < -1 \wedge m \geq 0$ ,

let  $Q_{q-2}[x] \rightarrow \text{PolynomialQuotient}[Pq[x], a+b x^2, x]$  and  $f+g x \rightarrow \text{PolynomialRemainder}[Pq[x], a+b x^2, x]$ , then

$$\int (c x)^m Pq[x] (a + b x^2)^p dx \rightarrow$$

$$\int (c x)^m (f + g x) (a + b x^2)^p dx + \int (c x)^m Q_{q-2}[x] (a + b x^2)^{p+1} dx \rightarrow$$

$$-\frac{(c x)^{m+1} (f + g x) (a + b x^2)^{p+1}}{2 a c (p + 1)} + \frac{1}{2 a (p + 1)} \int (c x)^m (a + b x^2)^{p+1} (2 a (p + 1) Q_{q-2}[x] + f (m + 2 p + 3) + g (m + 2 p + 4) x) dx$$

### Program code:

```
Int[(c_.*x_)^m_.*Pq_*(a_+b_.*x_^2)^p_,x_Symbol]:=  
With[{Q=PolynomialQuotient[Pq,a+b*x^2,x],  
f=Coeff[PolynomialRemainder[Pq,a+b*x^2,x],x,0],  
g=Coeff[PolynomialRemainder[Pq,a+b*x^2,x],x,1]},  
-(c*x)^(m+1)*(f+g*x)*(a+b*x^2)^(p+1)/(2*a*c*(p+1)) +  
1/(2*a*(p+1))*Int[(c*x)^m*(a+b*x^2)^(p+1)*ExpandToSum[2*a*(p+1)*Q+f*(m+2*p+3)+g*(m+2*p+4)*x,x]] /;  
FreeQ[{a,b,c,m},x] && PolyQ[Pq,x] && LtQ[p,-1] && Not[GtQ[m,0]]
```

7:  $\int (c x)^m Pq[x] (a + b x^2)^p dx$  when  $m < -1$

Derivation: Algebraic expansion and quadratic recurrence 3b

Note: If  $q = 1$ , no need to reduce integrand since  $\int (c x)^m Pq[x] (a + b x^2)^p dx$  can be expressed as a two term sum of hyperbolic functions.

Rule 1.1.2.y.7: If  $m < -1$ ,

let  $Q_{q-1}[x] \rightarrow \text{PolynomialQuotient}[Pq[x], c x, x]$  and  $R \rightarrow \text{PolynomialRemainder}[Pq[x], c x, x]$ , then

$$\begin{aligned} & \int (c x)^m Pq[x] (a + b x^2)^p dx \rightarrow \\ & \int (c x)^{m+1} Q_{q-1}[x] (a + b x^2)^p dx + R \int (c x)^m (a + b x^2)^p dx \rightarrow \\ & \frac{R (c x)^{m+1} (a + b x^2)^{p+1}}{a c (m+1)} + \frac{1}{a c (m+1)} \int (c x)^{m+1} (a + b x^2)^p (a c (m+1) Q_{q-1}[x] - b R (m+2 p+3) x) dx \end{aligned}$$

Program code:

```
Int[(c.*x.)^m_*Pq_*(a+b.*x.^2)^p_,x_Symbol]:=  
With[{Q=PolynomialQuotient[Pq,c*x,x], R=PolynomialRemainder[Pq,c*x,x]},  
R*(c*x)^(m+1)*(a+b*x^2)^(p+1)/(a*c*(m+1)) +  
1/(a*c*(m+1))*Int[(c*x)^(m+1)*(a+b*x^2)^p*ExpandToSum[a*c*(m+1)*Q-b*R*(m+2*p+3)*x,x],x]/;  
FreeQ[{a,b,c,p},x] && PolyQ[Pq,x] && LtQ[m,-1] && (IntegerQ[2*p] || NeQ[Expon[Pq,x],1])
```

8:  $\int (c x)^m Pq[x] (a + b x^2)^p dx$  when  $m + q + 2 p + 1 = 0$

Derivation: Algebraic expansion

Basis:  $(c x)^m Pq[x] = \frac{Pq[x, q]}{c^q} (c x)^{m+q} + \frac{(c x)^m (c^q Pq[x] - Pq[x, q] (c x)^q)}{c^q}$

Rule 1.1.2.y.8: If  $m + q + 2 p + 1 = 0$ , then

$$\begin{aligned} & \int (c x)^m Pq[x] (a + b x^2)^p dx \rightarrow \\ & \frac{Pq[x, q]}{c^q} \int (c x)^{m+q} (a + b x^2)^p dx + \frac{1}{c^q} \int (c x)^m (a + b x^2)^p (c^q Pq[x] - Pq[x, q] (c x)^q) dx \end{aligned}$$

Program code:

```
Int[(c_.*x_)^m_.*Pq_*(a_+b_.*x_^2)^p_,x_Symbol]:=  
With[{q=Expon[Pq,x]},  
Coeff[Pq,x,q]/c^q*Int[(c*x)^(m+q)*(a+b*x^2)^p,x]+  
1/c^q*Int[(c*x)^m*(a+b*x^2)^p*ExpandToSum[c^q*Pq-Coeff[Pq,x,q]*(c*x)^q,x],x]/;  
EqQ[q,1]||EqQ[m+q+2*p+1,0]]/;  
FreeQ[{a,b,c,m,p},x] && PolyQ[Pq,x] && Not[IGtQ[m,0] && ILtQ[p+1/2,0]]
```

9:  $\int (c x)^m Pq[x] (a + b x^2)^p dx$  when  $q > 1 \wedge m + q + 2 p + 1 \neq 0 \wedge (m \notin \mathbb{Z}^+ \vee p + \frac{1}{2} + 1 \in \mathbb{Z}^+)$

Derivation: Algebraic expansion and quadratic recurrence 3a with  $A = d$ ,  $B = e$  and  $m = m - 1$

Rule 1.1.2.y.9: If  $q > 1 \wedge m + q + 2 p + 1 \neq 0 \wedge (m \notin \mathbb{Z}^+ \vee p + \frac{1}{2} + 1 \in \mathbb{Z}^+)$ , let  $f \rightarrow Pq[x, q]$ , then

$$\int (c x)^m Pq[x] (a + b x^2)^p dx \rightarrow$$

$$\int (c x)^m \left( Pq[x] - \frac{f}{c^q} (c x)^q \right) (a + b x^2)^p dx + \frac{f}{c^q} \int (c x)^{m+q} (a + b x^2)^p dx \rightarrow$$

$$\frac{f (c x)^{m+q-1} (a + b x^2)^{p+1}}{b c^{q-1} (m + q + 2 p + 1)} + \frac{1}{b (m + q + 2 p + 1)} \int (c x)^m (a + b x^2)^p (b (m + q + 2 p + 1) P_q[x] - b f (m + q + 2 p + 1) x^q - a f (m + q - 1) x^{q-2}) dx$$

## Program code:

```

Int[(c_.*x_)^m_.*Pq_*(a_+b_.*x_^2)^p_,x_Symbol]:=

With[{q=Expon[Pq,x],f=Coeff[Pq,x,Expon[Pq,x]]},
f*(c*x)^(m+q-1)*(a+b*x^2)^(p+1)/(b*c^(q-1)*(m+q+2*p+1)) +
1/(b*(m+q+2*p+1))*Int[(c*x)^m*(a+b*x^2)^p*ExpandToSum[b*(m+q+2*p+1)*Pq-b*f*(m+q+2*p+1)*x^q-a*f*(m+q-1)*x^(q-2),x],x] /;

GtQ[q,1] && NeQ[m+q+2*p+1,0]] /;

FreeQ[{a,b,c,m,p},x] && PolyQ[Pq,x] && (Not[IGtQ[m,0]] || IGtQ[p+1/2,-1])

```