

Rules for integrands of the form $P_q[x] (a + b x^2)^p$

1: $\int P_q[x] (a + b x^2)^p dx$ when $p + 2 \in \mathbb{Z}^+$

Derivation: Algebraic expansion

Rule 1.1.2.x.1: If $p + 2 \in \mathbb{Z}^+$, then

$$\int P_q[x] (a + b x^2)^p dx \rightarrow \int \text{ExpandIntegrand}[P_q[x] (a + b x^2)^p, x] dx$$

Program code:

```
Int [Pq_* (a_+b_.*x_^2)^p_, x_Symbol] :=  
  Int [ExpandIntegrand [Pq*(a+b*x^2)^p,x], x] /;  
FreeQ[{a,b},x] && PolyQ[Pq,x] && IGtQ[p,-2]
```

2: $\int P_q[x] (a + b x^2)^p dx$ when $P_q[x, 0] = 0$

Derivation: Algebraic simplification

Rule 1.1.2.x.2: If $P_q[x, 0] = 0$, then

$$\int P_q[x] (a + b x^2)^p dx \rightarrow \int x \text{PolynomialQuotient}[P_q[x], x, x] (a + b x^2)^p dx$$

Program code:

```
Int [Pq_* (a_+b_.*x_^2)^p_, x_Symbol] :=  
  Int [x*PolynomialQuotient [Pq,x,x]*(a+b*x^2)^p,x] /;  
FreeQ[{a,b,p},x] && PolyQ[Pq,x] && EqQ[Coeff[Pq,x,0],0] && Not[MatchQ[Pq,x^m_.*u_./; IntegerQ[m]]]
```

$$3: \int P_q[x] (a + b x^2)^p dx \text{ when } \text{PolynomialRemainder}[P_q[x], a + b x^2, x] = 0$$

Derivation: Algebraic expansion

Rule: If $\text{PolynomialRemainder}[P_q[x], a + b x^2, x] = 0$, then

$$\int P_q[x] (a + b x^2)^p dx \rightarrow \int \text{PolynomialQuotient}[P_q[x], a + b x^2, x] (a + b x^2)^{p+1} dx$$

Program code:

```
Int [Px_*(a_+b_.*x_^2)^p_.,x_Symbol] :=
  Int [PolynomialQuotient [Px,a+b*x^2,x] *(a+b*x^2)^(p+1),x] /;
FreeQ[{a,b,p},x] && PolyQ[Px,x] && EqQ[PolynomialRemainder [Px,a+b*x^2,x],0]
```

$$4. \int P_q[x] (a + b x^2)^p dx \text{ when } p < -1$$

$$1: \int P_q[x^2] (a + b x^2)^p dx \text{ when } p + \frac{1}{2} \in \mathbb{Z}^- \wedge 2q + 2p + 1 < 0$$

Derivation: Algebraic expansion and binomial recurrence 3b

$$\text{Basis: } \int (a + b x^2)^p dx = \frac{x (a + b x^2)^{p+1}}{a} - \frac{b (2p+3)}{a} \int x^2 (a + b x^2)^p dx$$

Note: Interestingly this rule eliminates the constant term of $P_q[x^2]$ rather than the highest degree term.

Rule 1.1.2.x.4.1: If $p + \frac{1}{2} \in \mathbb{Z}^- \wedge 2q + 2p + 1 < 0$, let $A \rightarrow P_q[x^2, 0]$ and $Q_{q-1}[x^2] \rightarrow \text{PolynomialQuotient}[P_q[x^2] - A, x^2, x]$, then

$$\int P_q[x^2] (a + b x^2)^p dx \rightarrow$$

$$A \int (a + b x^2)^p dx + \int x^2 Q_{q-1}[x^2] (a + b x^2)^p dx \rightarrow$$

$$\frac{Ax(a+bx^2)^{p+1}}{a} + \frac{1}{a} \int x^2 (a+bx^2)^p (aQ_{q-1}[x^2] - Ab(2p+3)) dx$$

Program code:

```
Int[Pq_*(a_+b_.*x_^2)^p_,x_Symbol] :=
  With[{A=Coeff[Pq,x,0],Q=PolynomialQuotient[Pq-Coeff[Pq,x,0],x^2,x]},
    A*x*(a+b*x^2)^(p+1)/a + 1/a*Int[x^2*(a+b*x^2)^p*(a*Q-A*b*(2*p+3)),x] /;
  FreeQ[{a,b},x] && PolyQ[Pq,x^2] && ILtQ[p+1/2,0] && LtQ[Expon[Pq,x]+2*p+1,0]
```

2: $\int P_q[x] (a+bx^2)^p dx$ when $p < -1$

Derivation: Algebraic expansion and quadratic recurrence 2a

Rule 1.1.2.x.4.2: If $p < -1$,

let $Q_{q-2}[x] \rightarrow \text{PolynomialQuotient}[P_q[x], a+bx^2, x]$ and $f+gx \rightarrow \text{PolynomialRemainder}[P_q[x], a+bx^2, x]$, then

$$\int P_q[x] (a+bx^2)^p dx \rightarrow$$

$$\int (f+gx) (a+bx^2)^p dx + \int Q_{q-2}[x] (a+bx^2)^{p+1} dx \rightarrow$$

$$\frac{(ag-bfx)(a+bx^2)^{p+1}}{2ab(p+1)} + \frac{1}{2a(p+1)} \int (a+bx^2)^{p+1} (2a(p+1)Q_{q-2}[x] + f(2p+3)) dx$$

Program code:

```
Int[Pq_*(a_+b_.*x_^2)^p_,x_Symbol] :=
  With[{Q=PolynomialQuotient[Pq,a+b*x^2,x],
    f=Coeff[PolynomialRemainder[Pq,a+b*x^2,x],x,0],
    g=Coeff[PolynomialRemainder[Pq,a+b*x^2,x],x,1]},
    (a*g-b*f*x)*(a+b*x^2)^(p+1)/(2*a*b*(p+1)) +
    1/(2*a*(p+1))*Int[(a+b*x^2)^(p+1)*ExpandToSum[2*a*(p+1)*Q+f*(2*p+3),x],x] /;
  FreeQ[{a,b},x] && PolyQ[Pq,x] && LtQ[p,-1]
```

5: $\int P_q[x] (a + bx^2)^p dx$ when $p \neq -1$

Reference: G&R 2.160.3

Derivation: Trinomial recurrence 3a with $A = 0$, $B = 1$ and $m = m - n$

Reference: G&R 2.104

Note: This special case of the Ostrogradskiy-Hermite integration method reduces the degree of the polynomial in the resulting integrand.

Rule 1.1.2.x.5: If $p \neq -1$, let $e \rightarrow P_q[x, q]$, then

$$\int P_q[x] (a + bx^2)^p dx \rightarrow$$

$$\int (P_q[x] - ex^q) (a + bx^2)^p dx + e \int x^q (a + bx^2)^p dx \rightarrow$$

$$\frac{ex^{q-1} (a + bx^2)^{p+1}}{b(q+2p+1)} + \frac{1}{b(q+2p+1)} \int (a + bx^2)^p (b(q+2p+1)P_q[x] - ae(q-1)x^{q-2} - be(q+2p+1)x^q) dx$$

Program code:

```
Int [Pq_ * (a_+b_.*x_^2)^p_,x_Symbol] :=
  With [ {q=Expon [Pq,x],e=Coeff [Pq,x,Expon [Pq,x]] },
    e*x^(q-1) * (a+b*x^2)^(p+1) / (b*(q+2*p+1)) +
    1 / (b*(q+2*p+1)) * Int [ (a+b*x^2)^p * ExpandToSum [ b*(q+2*p+1) * Pq - a*e*(q-1) * x^(q-2) - b*e*(q+2*p+1) * x^q, x ] ] /;
  FreeQ [ {a,b,p}, x ] && PolyQ [ Pq, x ] && Not [ LeQ [ p, -1 ] ]
```