

Rules for integrands of the form $(g \operatorname{Sec}[e + f x])^p (a + b \operatorname{Sec}[e + f x])^m (c + d \operatorname{Sec}[e + f x])^n$

1. $\int (g \operatorname{Sec}[e + f x])^p (a + b \operatorname{Sec}[e + f x])^m (c + d \operatorname{Sec}[e + f x])^n dx$ when $b c + a d = 0 \wedge a^2 - b^2 = 0$

1. $\int \operatorname{Sec}[e + f x] (a + b \operatorname{Sec}[e + f x])^m (c + d \operatorname{Sec}[e + f x])^n dx$ when $b c + a d = 0 \wedge a^2 - b^2 = 0$

1. $\int \operatorname{Sec}[e + f x] (a + b \operatorname{Sec}[e + f x])^m (c + d \operatorname{Sec}[e + f x])^n dx$ when $b c + a d = 0 \wedge a^2 - b^2 = 0 \wedge m + n \in \mathbb{Z}^-$

1: $\int \operatorname{Sec}[e + f x] (a + b \operatorname{Sec}[e + f x])^m (c + d \operatorname{Sec}[e + f x])^n dx$ when $b c + a d = 0 \wedge a^2 - b^2 = 0 \wedge m + n + 1 = 0 \wedge m \neq -\frac{1}{2}$

Rule: If $b c + a d = 0 \wedge a^2 - b^2 = 0 \wedge m + n + 1 = 0 \wedge m \neq -\frac{1}{2}$, then

$$\int \operatorname{Sec}[e + f x] (a + b \operatorname{Sec}[e + f x])^m (c + d \operatorname{Sec}[e + f x])^n dx \rightarrow -\frac{b \operatorname{Tan}[e + f x] (a + b \operatorname{Sec}[e + f x])^m (c + d \operatorname{Sec}[e + f x])^n}{a f (2m + 1)}$$

Program code:

```
Int[csc[e_+f_*x_]*(a+b*csc[e_+f_*x_])^m*(c+d*csc[e_+f_*x_])^n_,x_Symbol]:=  
  b*Cot[e+f*x]*(a+b*Csc[e+f*x])^m*(c+d*Csc[e+f*x])^n/(a*f*(2*m+1)) /;  
FreeQ[{a,b,c,d,e,f,m,n},x] && EqQ[b*c+a*d,0] && EqQ[a^2-b^2,0] && EqQ[m+n+1,0] && NeQ[2*m+1,0]
```

2: $\int \sec[e + f x] (a + b \sec[e + f x])^m (c + d \sec[e + f x])^n dx$ when $b c + a d = 0 \wedge a^2 - b^2 = 0 \wedge m + n + 1 \in \mathbb{Z}^- \wedge m \neq -\frac{1}{2}$

Note: If $n + \frac{1}{2} \in \mathbb{Z}^+ \wedge n + \frac{1}{2} < -(m+n)$, then it is better to drive n to $\frac{1}{2}$ in $n - \frac{1}{2}$ steps.

Rule: If $b c + a d = 0 \wedge a^2 - b^2 = 0 \wedge m + n + 1 \in \mathbb{Z}^- \wedge m \neq -\frac{1}{2}$, then

$$\int \sec[e + f x] (a + b \sec[e + f x])^m (c + d \sec[e + f x])^n dx \rightarrow$$

$$-\frac{b \tan[e + f x] (a + b \sec[e + f x])^m (c + d \sec[e + f x])^n}{a f (2m+1)} + \frac{(m+n+1)}{a (2m+1)} \int \sec[e + f x] (a + b \sec[e + f x])^{m+1} (c + d \sec[e + f x])^n dx$$

Program code:

```
Int[csc[e_.+f_.*x_]*(a_+b_.*csc[e_.+f_.*x_])^m*(c_+d_.*csc[e_.+f_.*x_])^n_,x_Symbol]:=  
b*Cot[e+f*x]*(a+b*Csc[e+f*x])^m*(c+d*Csc[e+f*x])^n/(a*f*(2*m+1)) +  
(m+n+1)/(a*(2*m+1))*Int[Csc[e+f*x]*(a+b*Csc[e+f*x])^(m+1)*(c+d*Csc[e+f*x])^n,x] /;  
FreeQ[{a,b,c,d,e,f,m,n},x] && EqQ[b*c+a*d,0] && EqQ[a^2-b^2,0] && ILtQ[m+n+1,0] && NeQ[2*m+1,0] && Not[LtQ[n,0]] &&  
Not[IGtQ[n+1/2,0] && LtQ[n+1/2,-(m+n)]]
```

2. $\int \sec[e+f x] (a+b \sec[e+f x])^m (c+d \sec[e+f x])^n dx$ when $b c + a d = 0 \wedge a^2 - b^2 = 0 \wedge m + \frac{1}{2} \in \mathbb{Z}^+$

1. $\int \sec[e+f x] (a+b \sec[e+f x])^m \sqrt{c+d \sec[e+f x]} dx$ when $b c + a d = 0 \wedge a^2 - b^2 = 0$

1: $\int \frac{\sec[e+f x] \sqrt{c+d \sec[e+f x]}}{\sqrt{a+b \sec[e+f x]}} dx$ when $b c + a d = 0 \wedge a^2 - b^2 = 0$

Rule: If $b c + a d = 0 \wedge a^2 - b^2 = 0$, then

$$\int \frac{\sec[e+f x] \sqrt{c+d \sec[e+f x]}}{\sqrt{a+b \sec[e+f x]}} dx \rightarrow -\frac{a c \log[1 + \frac{b}{a} \sec[e+f x]] \tan[e+f x]}{b f \sqrt{a+b \sec[e+f x]} \sqrt{c+d \sec[e+f x]}}$$

Program code:

```
Int[csc[e_+f_*x_]*Sqrt[c_+d_.*csc[e_+f_*x_]]/Sqrt[a_+b_.*csc[e_+f_*x_]],x_Symbol]:=  
a*c*Log[1+b/a*Csc[e+f*x]]*Cot[e+f*x]/(b*f*Sqrt[a+b*Csc[e+f*x]]*Sqrt[c+d*Csc[e+f*x]]) /;  
FreeQ[{a,b,c,d,e,f},x] && EqQ[b*c+a*d,0] && EqQ[a^2-b^2,0]
```

2: $\int \sec[e+f x] (a+b \sec[e+f x])^m \sqrt{c+d \sec[e+f x]} dx$ when $b c + a d = 0 \wedge a^2 - b^2 = 0 \wedge m \neq -\frac{1}{2}$

Rule: If $b c + a d = 0 \wedge a^2 - b^2 = 0 \wedge m \neq -\frac{1}{2}$, then

$$\int \sec[e+f x] (a+b \sec[e+f x])^m \sqrt{c+d \sec[e+f x]} dx \rightarrow -\frac{2 a c \tan[e+f x] (a+b \sec[e+f x])^m}{b f (2 m + 1) \sqrt{c+d \sec[e+f x]}}$$

Program code:

```
Int[csc[e_+f_*x_]*(a_+b_.*csc[e_+f_*x_])^m.*Sqrt[c_+d_.*csc[e_+f_*x_]],x_Symbol]:=  
2*a*c*Cot[e+f*x]*(a+b*Csc[e+f*x])^m/(b*f*(2*m+1)*Sqrt[c+d*Csc[e+f*x]]) /;  
FreeQ[{a,b,c,d,e,f,m},x] && EqQ[b*c+a*d,0] && EqQ[a^2-b^2,0] && NeQ[m,-1/2]
```

2. $\int \sec[e + f x] (a + b \sec[e + f x])^m (c + d \sec[e + f x])^n dx$ when $b c + a d = 0 \wedge a^2 - b^2 = 0 \wedge m - \frac{1}{2} \in \mathbb{Z}^+$

1: $\int \sec[e + f x] (a + b \sec[e + f x])^m (c + d \sec[e + f x])^n dx$ when $b c + a d = 0 \wedge a^2 - b^2 = 0 \wedge n - \frac{1}{2} \in \mathbb{Z}^+ \wedge m < -\frac{1}{2}$

Rule: If $b c + a d = 0 \wedge a^2 - b^2 = 0 \wedge n - \frac{1}{2} \in \mathbb{Z}^+ \wedge m < -\frac{1}{2}$, then

$$\int \sec[e + f x] (a + b \sec[e + f x])^m (c + d \sec[e + f x])^n dx \rightarrow$$

$$-\frac{2 a c \tan[e + f x] (a + b \sec[e + f x])^m (c + d \sec[e + f x])^{n-1}}{b f (2 m + 1)} - \frac{d (2 n - 1)}{b (2 m + 1)} \int \sec[e + f x] (a + b \sec[e + f x])^{m+1} (c + d \sec[e + f x])^{n-1} dx$$

Program code:

```
Int[csc[e_.*f_.*x_]*(a_+b_.*csc[e_.*f_.*x_])^m*(c_+d_.*csc[e_.*f_.*x_])^n_,x_Symbol]:=  
2*a*c*Cot[e+f*x]*(a+b*Csc[e+f*x])^m*(c+d*Csc[e+f*x])^(n-1)/(b*f*(2*m+1)) -  
d*(2*n-1)/(b*(2*m+1))*Int[Csc[e+f*x]*(a+b*Csc[e+f*x])^(m+1)*(c+d*Csc[e+f*x])^(n-1),x] /;  
FreeQ[{a,b,c,d,e,f},x] && EqQ[b*c+a*d,0] && EqQ[a^2-b^2,0] && IGtQ[n-1/2,0] && LtQ[m,-1/2]
```

2: $\int \sec[e + f x] (a + b \sec[e + f x])^m (c + d \sec[e + f x])^n dx$ when $b c + a d = 0 \wedge a^2 - b^2 = 0 \wedge n - \frac{1}{2} \in \mathbb{Z}^+ \wedge m \neq -\frac{1}{2}$

Rule: If $b c + a d = 0 \wedge a^2 - b^2 = 0 \wedge n - \frac{1}{2} \in \mathbb{Z}^+ \wedge m \neq -\frac{1}{2}$, then

$$\int \sec[e + f x] (a + b \sec[e + f x])^m (c + d \sec[e + f x])^n dx \rightarrow$$

$$\frac{d \tan[e + f x] (a + b \sec[e + f x])^m (c + d \sec[e + f x])^{n-1}}{f(m+n)} + \frac{c(2n-1)}{m+n} \int \sec[e + f x] (a + b \sec[e + f x])^m (c + d \sec[e + f x])^{n-1} dx$$

Program code:

```
Int[csc[e_.*f_.*x_]*(a_+b_.*csc[e_.*f_.*x_])^m_.* (c_+d_.*csc[e_.*f_.*x_])^n_,x_Symbol]:=  
-d*Cot[e+f*x]* (a+b*Csc[e+f*x])^m*(c+d*Csc[e+f*x])^(n-1)/(f*(m+n)) +  
c*(2*n-1)/(m+n)*Int[Csc[e+f*x]* (a+b*Csc[e+f*x])^m*(c+d*Csc[e+f*x])^(n-1),x] /;  
FreeQ[{a,b,c,d,e,f,m},x] && EqQ[b*c+a*d,0] && EqQ[a^2-b^2,0] && IGtQ[n-1/2,0] && Not[LTQ[m,-1/2]] && Not[IGtQ[m-1/2,0] && LTQ[m,n]]
```

3. $\int \sec[e+f x] (a+b \sec[e+f x])^m (c+d \sec[e+f x])^n dx$ when $b c + a d = 0 \wedge a^2 - b^2 = 0 \wedge n \in \mathbb{Z}^+ \wedge m < 0$

1: $\int \frac{\sec[e+f x] (c+d \sec[e+f x])^n}{\sqrt{a+b \sec[e+f x]}} dx$ when $b c + a d = 0 \wedge a^2 - b^2 = 0 \wedge n \in \mathbb{Z}^+$

Rule: If $b c + a d = 0 \wedge a^2 - b^2 = 0 \wedge n \in \mathbb{Z}^+$, then

$$\int \frac{\sec[e+f x] (c+d \sec[e+f x])^n}{\sqrt{a+b \sec[e+f x]}} dx \rightarrow$$

$$\frac{2 d \tan[e+f x] (c+d \sec[e+f x])^{n-1}}{f (2 n - 1) \sqrt{a+b \sec[e+f x]}} + \frac{2 c (2 n - 1)}{2 n - 1} \int \frac{\sec[e+f x] (c+d \sec[e+f x])^{n-1}}{\sqrt{a+b \sec[e+f x]}} dx$$

Program code:

```
Int[csc[e_.+f_.*x_]*(c_.+d_.*csc[e_.+f_.*x_])^n_./Sqrt[a_+b_.*csc[e_.+f_.*x_]],x_Symbol]:=
-2*d*Cot[e+f*x]*(c+d*Csc[e+f*x])^(n-1)/(f*(2*n-1)*Sqrt[a+b*Csc[e+f*x]])+
2*c*(2*n-1)/(2*n-1)*Int[Csc[e+f*x]*(c+d*Csc[e+f*x])^(n-1)/Sqrt[a+b*Csc[e+f*x]],x] /;
FreeQ[{a,b,c,d,e,f},x] && EqQ[b*c+a*d,0] && EqQ[a^2-b^2,0] && IGtQ[n,0]
```

2: $\int \sec[e + f x] (a + b \sec[e + f x])^m (c + d \sec[e + f x])^n dx$ when $b c + a d = 0 \wedge a^2 - b^2 = 0 \wedge n \in \mathbb{Z}^+ \wedge m < -\frac{1}{2}$

Rule: If $b c + a d = 0 \wedge a^2 - b^2 = 0 \wedge n \in \mathbb{Z}^+ \wedge m < -\frac{1}{2}$, then

$$\int \sec[e + f x] (a + b \sec[e + f x])^m (c + d \sec[e + f x])^n dx \rightarrow$$

$$-\frac{2 a c \tan[e + f x] (a + b \sec[e + f x])^m (c + d \sec[e + f x])^{n-1}}{b f (2 m + 1)} - \frac{d (2 n - 1)}{b (2 m + 1)} \int \sec[e + f x] (a + b \sec[e + f x])^{m+1} (c + d \sec[e + f x])^{n-1} dx$$

Program code:

```
Int[csc[e_.*f_.*x_]*(a_+b_.*csc[e_.*f_.*x_])^m*(c_+d_.*csc[e_.*f_.*x_])^n_,x_Symbol]:=  
  2*a*c*Cot[e+f*x]*(a+b*Csc[e+f*x])^m*(c+d*Csc[e+f*x])^(n-1)/(b*f*(2*m+1)) -  
  d*(2*n-1)/(b*(2*m+1))*Int[Csc[e+f*x]*(a+b*Csc[e+f*x])^(m+1)*(c+d*Csc[e+f*x])^(n-1),x];  
FreeQ[{a,b,c,d,e,f},x] && EqQ[b*c+a*d,0] && EqQ[a^2-b^2,0] && IGtQ[n,0] && LtQ[m,-1/2] && IntegerQ[2*m]
```

4: $\int \sec[e + f x] (a + b \sec[e + f x])^m (c + d \sec[e + f x])^n dx$ when $b c + a d = 0 \wedge a^2 - b^2 = 0 \wedge m \in \mathbb{Z} \wedge n \in \mathbb{Z} \wedge n - m \geq 0 \wedge m n > 0$

Derivation: Algebraic simplification

Basis: If $b c + a d = 0 \wedge a^2 - b^2 = 0$, then $(a + b \sec[z]) (c + d \sec[z]) = -a c \tan[z]^2$

Rule: If $b c + a d = 0 \wedge a^2 - b^2 = 0 \wedge m \in \mathbb{Z} \wedge n \in \mathbb{Z} \wedge n - m \geq 0 \wedge m n > 0$, then

$$\int \sec[e + f x] (a + b \sec[e + f x])^m (c + d \sec[e + f x])^n dx \rightarrow (-a c)^m \int (g \sec[e + f x])^p \tan[e + f x]^{2m} (c + d \sec[e + f x])^{n-m} dx$$

Program code:

```
Int[csc[e_+f_*x_]*(a_+b_*csc[e_+f_*x_])^m_*(c_+d_*csc[e_+f_*x_])^n_,x_Symbol]:=(-a*c)^m*Int[ExpandTrig[csc[e+f*x]*cot[e+f*x]^(2*m),(c+d*csc[e+f*x])^(n-m),x],x]/;
FreeQ[{a,b,c,d,e,f,n},x] && EqQ[b*c+a*d,0] && EqQ[a^2-b^2,0] && IntegersQ[m,n] && GeQ[n-m,0] && GtQ[m*n,0]
```

5: $\int \sec[e+f x] (a+b \sec[e+f x])^m (c+d \sec[e+f x])^n dx$ when $b c + a d = 0 \wedge a^2 - b^2 = 0 \wedge m - \frac{1}{2} \in \mathbb{Z}^-$

Derivation: Algebraic expansion and piecewise constant extraction

Basis: If $b c + a d = 0 \wedge a^2 - b^2 = 0 \wedge m + \frac{1}{2} \in \mathbb{Z}$, then $(a+b \sec[z])^m (c+d \sec[z])^m = \frac{(-a c)^{\frac{m+1}{2}} \tan[z]^{2m+1}}{\sqrt{a+b \sec[z]} \sqrt{c+d \sec[z]}}$

Basis: If $b c + a d = 0 \wedge a^2 - b^2 = 0$, then $\partial_x \frac{\tan[e+f x]}{\sqrt{a+b \sec[e+f x]} \sqrt{c+d \sec[e+f x]}} = 0$

Rule: If $b c + a d = 0 \wedge a^2 - b^2 = 0 \wedge m + \frac{1}{2} \in \mathbb{Z}$, then

$$\int \sec[e+f x] (a+b \sec[e+f x])^m (c+d \sec[e+f x])^n dx \rightarrow \frac{(-a c)^{\frac{m+1}{2}} \tan[e+f x]}{\sqrt{a+b \sec[e+f x]} \sqrt{c+d \sec[e+f x]}} \int \sec[e+f x] \tan[e+f x]^{2m} dx$$

Program code:

```
Int[csc[e_.+f_.*x_]*(a_+b_.*csc[e_.+f_.*x_])^m*(c_+d_.*csc[e_.+f_.*x_])^n,x_Symbol]:= (-a*c)^(m+1/2)*Cot[e+f*x]/(Sqrt[a+b*Csc[e+f*x]]*Sqrt[c+d*Csc[e+f*x]])*Int[Csc[e+f*x]*Cot[e+f*x]^(2*m),x]; FreeQ[{a,b,c,d,e,f},x] && EqQ[b*c+a*d,0] && EqQ[a^2-b^2,0] && IntegerQ[m+1/2]
```

6: $\int \sec[e+f x] (a+b \sec[e+f x])^m (c+d \sec[e+f x])^n dx$ when $b c + a d = 0 \wedge a^2 - b^2 = 0 \wedge ((m | n - \frac{1}{2}) \in \mathbb{Z}^- \vee (m - \frac{1}{2} | n - \frac{1}{2}) \in \mathbb{Z}^-)$

Rule: If $b c + a d = 0 \wedge a^2 - b^2 = 0 \wedge (m \in \mathbb{Z}^- \vee (m - \frac{1}{2} | n - \frac{1}{2}) \in \mathbb{Z}^-)$, then

$$\int \sec[e+f x] (a+b \sec[e+f x])^m (c+d \sec[e+f x])^n dx \rightarrow$$

$$-\frac{b \tan[e+f x] (a+b \sec[e+f x])^m (c+d \sec[e+f x])^n}{a f (2m+1)} + \frac{(m+n+1)}{a (2m+1)} \int \sec[e+f x] (a+b \sec[e+f x])^{m+1} (c+d \sec[e+f x])^n dx$$

Program code:

```
Int[csc[e_..+f_..*x_]*(a_+b_.*csc[e_..+f_..*x_])^m*(c_+d_.*csc[e_..+f_..*x_])^n,x_Symbol]:=  
b*Cot[e+f*x]*(a+b*Csc[e+f*x])^m*(c+d*Csc[e+f*x])^n/(a*f*(2*m+1)) +  
(m+n+1)/(a*(2*m+1))*Int[Csc[e+f*x]*(a+b*Csc[e+f*x])^(m+1)*(c+d*Csc[e+f*x])^n,x]/;  
FreeQ[{a,b,c,d,e,f,n},x] && EqQ[b*c+a*d,0] && EqQ[a^2-b^2,0] && (ILtQ[m,0] && ILtQ[n-1/2,0] || ILtQ[m-1/2,0] && ILtQ[n-1/2,0] && LtQ[m,n])
```

7: $\int \sec[e+f x] (a+b \sec[e+f x])^m (c+d \sec[e+f x])^n dx$ when $b c + a d = 0 \wedge a^2 - b^2 = 0$

Derivation: Piecewise constant extraction and integration by substitution

Basis: If $b c + a d = 0 \wedge a^2 - b^2 = 0$, then $\partial_x \frac{\tan[e+f x]}{\sqrt{a+b \sec[e+f x]} \sqrt{c+d \sec[e+f x]}} = 0$

Basis: If $b c + a d = 0 \wedge a^2 - b^2 = 0$, then $-\frac{a c \tan[e+f x]}{\sqrt{a+b \sec[e+f x]} \sqrt{c+d \sec[e+f x]}} - \frac{\tan[e+f x]}{\sqrt{a+b \sec[e+f x]} \sqrt{c+d \sec[e+f x]}} = 1$

Basis: $\tan[e+f x] F[\sec[e+f x]] = \frac{1}{f} \text{Subst}\left[\frac{F[x]}{x}, x, \sec[e+f x]\right] \partial_x \sec[e+f x]$

Rule: If $b c + a d = 0 \wedge a^2 - b^2 = 0$, then

$$\int \sec[e+f x] (a+b \sec[e+f x])^m (c+d \sec[e+f x])^n dx \rightarrow$$

$$\begin{aligned}
& -\frac{a c \tan[e+f x]}{\sqrt{a+b \sec[e+f x]} \sqrt{c+d \sec[e+f x]}} \int \tan[e+f x] \sec[e+f x] (a+b \sec[e+f x])^{\frac{m-1}{2}} (c+d \sec[e+f x])^{\frac{n-1}{2}} dx \rightarrow \\
& -\frac{a c \tan[e+f x]}{f \sqrt{a+b \sec[e+f x]} \sqrt{c+d \sec[e+f x]}} \text{Subst} \left[\int (a+b x)^{\frac{m-1}{2}} (c+d x)^{\frac{n-1}{2}} dx, x, \sec[e+f x] \right]
\end{aligned}$$

Program code:

```

Int[csc[e_+f_*x_]*(a_+b_.*csc[e_+f_*x_])^m_.*(c_+d_.*csc[e_+f_*x_])^n_,x_Symbol]:= 
  a*c*Cot[e+f*x]/(f*Sqrt[a+b*Csc[e+f*x]]*Sqrt[c+d*Csc[e+f*x]])*Subst[Int[(a+b*x)^(m-1/2)*(c+d*x)^(n-1/2),x],x,Csc[e+f*x]] /;
FreeQ[{a,b,c,d,e,f,m,n},x] && EqQ[b*c+a*d,0] && EqQ[a^2-b^2,0]

```

2: $\int (g \sec[e+f x])^p (a+b \sec[e+f x])^m (c+d \sec[e+f x])^n dx$ when $b c + a d = 0 \wedge a^2 - b^2 = 0 \wedge m \in \mathbb{Z} \wedge n \in \mathbb{Z} \wedge n - m \geq 0 \wedge m n > 0$

Derivation: Algebraic simplification

Basis: If $b c + a d = 0 \wedge a^2 - b^2 = 0$, then $(a+b \sec[z]) (c+d \sec[z]) = -a c \tan[z]^2$

Rule: If $b c + a d = 0 \wedge a^2 - b^2 = 0 \wedge m \in \mathbb{Z} \wedge n \in \mathbb{Z} \wedge n - m \geq 0 \wedge m n > 0$, then

$$\int (g \sec[e+f x])^p (a+b \sec[e+f x])^m (c+d \sec[e+f x])^n dx \rightarrow (-a c)^m \int (g \sec[e+f x])^p \tan[e+f x]^{2m} (c+d \sec[e+f x])^{n-m} dx$$

Program code:

```

Int[(g_.*csc[e_+f_*x_])^p_.*(a_+b_.*csc[e_+f_*x_])^m_.*(c_+d_.*csc[e_+f_*x_])^n_,x_Symbol]:= 
  (-a*c)^m*Int[ExpandTrig[(g*csc[e+f*x])^p*cot[e+f*x]^(2*m),(c+d*csc[e+f*x])^(n-m),x],x] /;
FreeQ[{a,b,c,d,e,f,g,n,p},x] && EqQ[b*c+a*d,0] && EqQ[a^2-b^2,0] && IntegersQ[m,n] && GeQ[n-m,0] && GtQ[m*n,0]

```

3: $\int (g \sec[e+f x])^p (a+b \sec[e+f x])^m (c+d \sec[e+f x])^n dx$ when $b c + a d = 0 \wedge a^2 - b^2 = 0 \wedge m - \frac{1}{2} \in \mathbb{Z}^-$

Derivation: Algebraic expansion and piecewise constant extraction

Basis: If $b c + a d = 0 \wedge a^2 - b^2 = 0 \wedge m + \frac{1}{2} \in \mathbb{Z}$, then $(a+b \sec[z])^m (c+d \sec[z])^m = \frac{(-a c)^{\frac{m+1}{2}} \tan[z]^{2m+1}}{\sqrt{a+b \sec[z]} \sqrt{c+d \sec[z]}}$

Basis: If $b c + a d = 0 \wedge a^2 - b^2 = 0$, then $\partial_x \frac{\tan[e+f x]}{\sqrt{a+b \sec[e+f x]} \sqrt{c+d \sec[e+f x]}} = 0$

Rule: If $b c + a d = 0 \wedge a^2 - b^2 = 0 \wedge m + \frac{1}{2} \in \mathbb{Z}$, then

$$\int (g \sec[e+f x])^p (a+b \sec[e+f x])^m (c+d \sec[e+f x])^n dx \rightarrow \frac{(-a c)^{\frac{m+1}{2}} \tan[e+f x]}{\sqrt{a+b \sec[e+f x]} \sqrt{c+d \sec[e+f x]}} \int (g \sec[e+f x])^p \tan[e+f x]^{2m} dx$$

Program code:

```
Int[(g_.*csc[e_.*f_.*x_])^p_.*(a_+b_.*csc[e_.*f_.*x_])^m_.*(c_+d_.*csc[e_.*f_.*x_])^n_,x_Symbol]:= (-a*c)^(m+1/2)*Cot[e+f*x]/(Sqrt[a+b*Csc[e+f*x]]*Sqrt[c+d*Csc[e+f*x]])*Int[(g*Csc[e+f*x])^p*Cot[e+f*x]^(2*m),x]; FreeQ[{a,b,c,d,e,f,g,p},x] && EqQ[b*c+a*d,0] && EqQ[a^2-b^2,0] && IntegerQ[m+1/2]
```

4: $\int (g \sec[e+f x])^p (a+b \sec[e+f x])^m (c+d \sec[e+f x])^n dx$ when $b c + a d = 0 \wedge a^2 - b^2 = 0$

Derivation: Piecewise constant extraction and integration by substitution

Basis: If $b c + a d = 0 \wedge a^2 - b^2 = 0$, then $\partial_x \frac{\tan[e+f x]}{\sqrt{a+b \sec[e+f x]} \sqrt{c+d \sec[e+f x]}} = 0$

Basis: If $b c + a d = 0 \wedge a^2 - b^2 = 0$, then $-\frac{a c \tan[e+f x]}{\sqrt{a+b \sec[e+f x]} \sqrt{c+d \sec[e+f x]}} - \frac{\tan[e+f x]}{\sqrt{a+b \sec[e+f x]} \sqrt{c+d \sec[e+f x]}} = 1$

Basis: $\tan[e+f x] F[\sec[e+f x]] = \frac{1}{f} \text{Subst}\left[\frac{F[x]}{x}, x, \sec[e+f x]\right] \partial_x \sec[e+f x]$

Rule: If $b c + a d = 0 \wedge a^2 - b^2 = 0$, then

$$\begin{aligned} & \int (g \sec[e+f x])^p (a+b \sec[e+f x])^m (c+d \sec[e+f x])^n dx \rightarrow \\ & -\frac{a c \tan[e+f x]}{\sqrt{a+b \sec[e+f x]} \sqrt{c+d \sec[e+f x]}} \int \tan[e+f x] (g \sec[e+f x])^p (a+b \sec[e+f x])^{m-\frac{1}{2}} (c+d \sec[e+f x])^{n-\frac{1}{2}} dx \rightarrow \\ & -\frac{a c g \tan[e+f x]}{f \sqrt{a+b \sec[e+f x]} \sqrt{c+d \sec[e+f x]}} \text{Subst}\left[\int (g x)^{p-1} (a+b x)^{m-\frac{1}{2}} (c+d x)^{n-\frac{1}{2}} dx, x, \sec[e+f x]\right] \end{aligned}$$

Program code:

```
Int[(g_.*csc[e_.*f_.*x_])^p_.*(a_+b_.*csc[e_.*f_.*x_])^m_.*(c_+d_.*csc[e_.*f_.*x_])^n_,x_Symbol]:=  
  a*c*g*Cot[e+f*x]/(f*Sqrt[a+b*Csc[e+f*x]]*Sqrt[c+d*Csc[e+f*x]])*  
  Subst[Int[(g*x)^(p-1)*(a+b*x)^(m-1/2)*(c+d*x)^(n-1/2),x],x,Csc[e+f*x]]/;  
  FreeQ[{a,b,c,d,e,f,g,m,n,p},x] && EqQ[b*c+a*d,0] && EqQ[a^2-b^2,0]
```

2. $\int \frac{(g \sec[e+f x])^p (a+b \sec[e+f x])^m}{c+d \sec[e+f x]} dx$ when $b c - a d \neq 0$

$$1. \int \frac{(g \sec[e+f x])^p \sqrt{a+b \sec[e+f x]}}{c+d \sec[e+f x]} dx \text{ when } b c - a d \neq 0$$

$$1. \int \frac{\sqrt{g \sec[e+f x]} \sqrt{a+b \sec[e+f x]}}{c+d \sec[e+f x]} dx \text{ when } b c - a d \neq 0$$

$$1: \int \frac{\sqrt{g \sec[e+f x]} \sqrt{a+b \sec[e+f x]}}{c+d \sec[e+f x]} dx \text{ when } b c - a d \neq 0 \wedge a^2 - b^2 = 0$$

Derivation: Integration by substitution

Basis: If $a^2 - b^2 = 0$, then $\frac{\sqrt{g \sec[e+f x]} \sqrt{a+b \sec[e+f x]}}{c+d \sec[e+f x]} = \frac{2 b g}{f} \text{Subst}\left[\frac{1}{b c + a d - c g x^2}, x, \frac{b \tan[e+f x]}{\sqrt{g \sec[e+f x]} \sqrt{a+b \sec[e+f x]}}\right] \partial_x \frac{b \tan[e+f x]}{\sqrt{g \sec[e+f x]} \sqrt{a+b \sec[e+f x]}}$

Rule: If $b c - a d \neq 0 \wedge a^2 - b^2 = 0$, then

$$\int \frac{\sqrt{g \sec[e+f x]} \sqrt{a+b \sec[e+f x]}}{c+d \sec[e+f x]} dx \rightarrow \frac{2 b g}{f} \text{Subst}\left[\int \frac{1}{b c + a d - c g x^2} dx, x, \frac{b \tan[e+f x]}{\sqrt{g \sec[e+f x]} \sqrt{a+b \sec[e+f x]}}\right]$$

Program code:

```
Int[Sqrt[g_.*csc[e_.*f_.*x_]]*Sqrt[a_+b_.*csc[e_.*f_.*x_]]/(c_+d_.*csc[e_.*f_.*x_]),x_Symbol]:=  
-2*b*g/f*Subst[Int[1/(b*c+a*d-c*g*x^2),x],x,b*Cot[e+f*x]/(Sqrt[g*Csc[e+f*x]]*Sqrt[a+b*Csc[e+f*x]])] /;  
FreeQ[{a,b,c,d,e,f,g},x] && NeQ[b*c-a*d,0] && EqQ[a^2-b^2,0]
```

$$2: \int \frac{\sqrt{g \sec[e+f x]} \sqrt{a+b \sec[e+f x]}}{c+d \sec[e+f x]} dx \text{ when } b c - a d \neq 0 \wedge a^2 - b^2 \neq 0$$

Derivation: Algebraic expansion

Basis: $\frac{\sqrt{a+b z}}{c+d z} = \frac{a}{c \sqrt{a+b z}} + \frac{(b c - a d) g z}{c g \sqrt{a+b z} (c+d z)}$

Rule: If $b c - a d \neq 0 \wedge a^2 - b^2 \neq 0$, then

$$\int \frac{\sqrt{g \sec[e+f x]} \sqrt{a+b \sec[e+f x]}}{c+d \sec[e+f x]} dx \rightarrow \frac{a}{c} \int \frac{\sqrt{g \sec[e+f x]}}{\sqrt{a+b \sec[e+f x]}} dx + \frac{b c - a d}{c g} \int \frac{(g \sec[e+f x])^{3/2}}{\sqrt{a+b \sec[e+f x]} (c+d \sec[e+f x])} dx$$

Program code:

```
Int[Sqrt[g_.*csc[e_+f_.*x_]]*Sqrt[a_+b_.*csc[e_+f_.*x_]]/(c_+d_.*csc[e_+f_.*x_]),x_Symbol]:=  
a/c*Int[Sqrt[g*Csc[e+f*x]]/Sqrt[a+b*Csc[e+f*x]],x]+  
(b*c-a*d)/(c*g)*Int[(g*Csc[e+f*x])^(3/2)/(Sqrt[a+b*Csc[e+f*x]]*(c+d*Csc[e+f*x])),x]/;  
FreeQ[{a,b,c,d,e,f,g},x] && NeQ[b*c-a*d,0] && NeQ[a^2-b^2,0]
```

2. $\int \frac{\sec[e+f x] \sqrt{a+b \sec[e+f x]}}{c+d \sec[e+f x]} dx$ when $b c - a d \neq 0$

1: $\int \frac{\sec[e+f x] \sqrt{a+b \sec[e+f x]}}{c+d \sec[e+f x]} dx$ when $b c - a d \neq 0 \wedge a^2 - b^2 = 0$

Derivation: Integration by substitution

Basis: If $a^2 - b^2 = 0$, then $\frac{\sec[e+f x] \sqrt{a+b \sec[e+f x]}}{c+d \sec[e+f x]} = \frac{2b}{f} \text{Subst}\left[\frac{1}{b c + a d + d x^2}, x, \frac{b \tan[e+f x]}{\sqrt{a+b \sec[e+f x]}}\right] \partial_x \frac{b \tan[e+f x]}{\sqrt{a+b \sec[e+f x]}}$

Rule: If $b c - a d \neq 0 \wedge a^2 - b^2 = 0$, then

$$\int \frac{\sec[e+f x] \sqrt{a+b \sec[e+f x]}}{c+d \sec[e+f x]} dx \rightarrow \frac{2b}{f} \text{Subst}\left[\int \frac{1}{b c + a d + d x^2} dx, x, \frac{b \tan[e+f x]}{\sqrt{a+b \sec[e+f x]}}\right]$$

Program code:

```
Int[csc[e_+f_.*x_]*Sqrt[a_+b_.*csc[e_+f_.*x_]]/(c_+d_.*csc[e_+f_.*x_]),x_Symbol]:=  
-2*b/f*Subst[Int[1/(b*c+a*d+d*x^2),x],x,b*Cot[e+f*x]/Sqrt[a+b*Csc[e+f*x]]]/;  
FreeQ[{a,b,c,d,e,f},x] && NeQ[b*c-a*d,0] && EqQ[a^2-b^2,0]
```

2. $\int \frac{\sec[e+f x] \sqrt{a+b \sec[e+f x]}}{c+d \sec[e+f x]} dx$ when $b c - a d \neq 0 \wedge a^2 - b^2 \neq 0$

1: $\int \frac{\sec[e+f x] \sqrt{a+b \sec[e+f x]}}{c+d \sec[e+f x]} dx$ when $b c - a d \neq 0 \wedge a^2 - b^2 \neq 0 \wedge c^2 - d^2 = 0$

Rule: If $b c - a d \neq 0 \wedge a^2 - b^2 \neq 0 \wedge c^2 - d^2 = 0$, then

$$\int \frac{\sec[e+f x] \sqrt{a+b \sec[e+f x]}}{c+d \sec[e+f x]} dx \rightarrow \frac{\sqrt{a+b \sec[e+f x]} \sqrt{\frac{c}{c+d \sec[e+f x]}}}{d f \sqrt{\frac{c d (a+b \sec[e+f x])}{(b c+a d) (c+d \sec[e+f x])}}} \text{EllipticE}\left[\text{ArcSin}\left[\frac{c \tan[e+f x]}{c+d \sec[e+f x]}\right], -\frac{b c-a d}{b c+a d}\right]$$

Program code:

```
Int[csc[e_.+f_.*x_] *Sqrt[a_+b_.*csc[e_.+f_.*x_]]/(c_+d_.*csc[e_.+f_.*x_]),x_Symbol]:=  
-Sqrt[a+b*Csc[e+f*x]]*Sqrt[c/(c+d*Csc[e+f*x])]/(d*f*Sqrt[c*d*(a+b*Csc[e+f*x])/((b*c+a*d)*(c+d*Csc[e+f*x]))])*  
EllipticE[ArcSin[c*Cot[e+f*x]/(c+d*Csc[e+f*x])],-(b*c-a*d)/(b*c+a*d)]/;  
FreeQ[{a,b,c,d,e,f},x] && NeQ[b*c-a*d,0] && NeQ[a^2-b^2,0] && EqQ[c^2-d^2,0]
```

2: $\int \frac{\sec[e+f x] \sqrt{a+b \sec[e+f x]}}{c+d \sec[e+f x]} dx$ when $b c - a d \neq 0 \wedge a^2 - b^2 \neq 0 \wedge c^2 - d^2 \neq 0$

Derivation: Algebraic expansion

Basis: $\frac{\sqrt{a+b z}}{c+d z} = \frac{b}{d \sqrt{a+b z}} - \frac{b c - a d}{d \sqrt{a+b z} (c+d z)}$

Rule: If $b c - a d \neq 0 \wedge a^2 - b^2 \neq 0 \wedge c^2 - d^2 \neq 0$, then

$$\int \frac{\sec[e+f x] \sqrt{a+b \sec[e+f x]}}{c+d \sec[e+f x]} dx \rightarrow \frac{b}{d} \int \frac{\sec[e+f x]}{\sqrt{a+b \sec[e+f x]}} dx - \frac{b c - a d}{d} \int \frac{\sec[e+f x]}{\sqrt{a+b \sec[e+f x]} (c+d \sec[e+f x])} dx$$

Program code:

```
Int[csc[e_.+f_.*x_] * Sqrt[a_+b_.*csc[e_.+f_.*x_]]/(c_+d_.*csc[e_.+f_.*x_]), x_Symbol] :=
  b/d*Int[Csc[e+f*x]/Sqrt[a+b*Csc[e+f*x]], x] -
  (b*c-a*d)/d*Int[Csc[e+f*x]/(Sqrt[a+b*Csc[e+f*x]]*(c+d*Csc[e+f*x])), x] /;
FreeQ[{a,b,c,d,e,f}, x] && NeQ[b*c-a*d, 0] && NeQ[a^2-b^2, 0] && NeQ[c^2-d^2, 0]
```

3. $\int \frac{(g \sec[e+f x])^{3/2} \sqrt{a+b \sec[e+f x]}}{c+d \sec[e+f x]} dx$ when $b c - a d \neq 0$

1: $\int \frac{(g \sec[e+f x])^{3/2} \sqrt{a+b \sec[e+f x]}}{c+d \sec[e+f x]} dx$ when $b c - a d \neq 0 \wedge a^2 - b^2 = 0$

Derivation: Algebraic expansion

Basis: $\frac{(g z)^{3/2}}{c+d z} = \frac{g \sqrt{g z}}{d} - \frac{c g \sqrt{g z}}{d (c+d z)}$

Rule: If $b c - a d \neq 0 \wedge a^2 - b^2 = 0$, then

$$\int \frac{(g \sec[e+f x])^{3/2} \sqrt{a+b \sec[e+f x]}}{c+d \sec[e+f x]} dx \rightarrow \frac{g}{d} \int \sqrt{g \sec[e+f x]} \sqrt{a+b \sec[e+f x]} dx - \frac{c g}{d} \int \frac{\sqrt{g \sec[e+f x]} \sqrt{a+b \sec[e+f x]}}{c+d \sec[e+f x]} dx$$

Program code:

```
Int[(g_.*csc[e_._+f_._*x_])^(3/2)*Sqrt[a_._+b_._*csc[e_._+f_._*x_]]/(c_._+d_._*csc[e_._+f_._*x_]),x_Symbol]:=  
g/d*Int[Sqrt[g*Csc[e+f*x]]*Sqrt[a+b*Csc[e+f*x]],x]-  
c*g/d*Int[Sqrt[g*Csc[e+f*x]]*Sqrt[a+b*Csc[e+f*x]]/(c+d*Csc[e+f*x]),x]/;  
FreeQ[{a,b,c,d,e,f,g},x] && NeQ[b*c-a*d,0] && EqQ[a^2-b^2,0]
```

2: $\int \frac{(g \sec[e+f x])^{3/2} \sqrt{a+b \sec[e+f x]}}{c+d \sec[e+f x]} dx$ when $b c - a d \neq 0 \wedge a^2 - b^2 \neq 0$

- Derivation: Algebraic expansion

- Basis: $\frac{\sqrt{a+b z}}{c+d z} = \frac{b}{d \sqrt{a+b z}} - \frac{b c - a d}{d \sqrt{a+b z} (c+d z)}$

- Rule: If $b c - a d \neq 0 \wedge a^2 - b^2 \neq 0$, then

$$\int \frac{(g \sec[e+f x])^{3/2} \sqrt{a+b \sec[e+f x]}}{c+d \sec[e+f x]} dx \rightarrow \frac{b}{d} \int \frac{(g \sec[e+f x])^{3/2}}{\sqrt{a+b \sec[e+f x]}} dx - \frac{b c - a d}{d} \int \frac{(g \sec[e+f x])^{3/2}}{\sqrt{a+b \sec[e+f x]} (c+d \sec[e+f x])} dx$$

Program code:

```
Int[(g_.*csc[e_._+f_._*x_])^(3/2)*Sqrt[a_._+b_._*csc[e_._+f_._*x_]]/(c_._+d_._*csc[e_._+f_._*x_]),x_Symbol]:=  
b/d*Int[(g*Csc[e+f*x])^(3/2)/Sqrt[a+b*Csc[e+f*x]],x]-  
(b*c-a*d)/d*Int[(g*Csc[e+f*x])^(3/2)/(Sqrt[a+b*Csc[e+f*x]]*(c+d*Csc[e+f*x])),x]/;  
FreeQ[{a,b,c,d,e,f,g},x] && NeQ[b*c-a*d,0] && NeQ[a^2-b^2,0]
```

2. $\int \frac{(g \sec[e+f x])^p}{\sqrt{a+b \sec[e+f x]} (c+d \sec[e+f x])} dx$ when $b c - a d \neq 0$

1. $\int \frac{\sec[e+f x]}{\sqrt{a+b \sec[e+f x]} (c+d \sec[e+f x])} dx$ when $b c - a d \neq 0$

1: $\int \frac{\sec[e+f x]}{\sqrt{a+b \sec[e+f x]} (c+d \sec[e+f x])} dx$ when $b c - a d \neq 0 \wedge (a^2 - b^2 = 0 \vee c^2 - d^2 = 0)$

Derivation: Algebraic expansion

Basis: $\frac{1}{\sqrt{a+b z} (c+d z)} = \frac{b}{(b c-a d) \sqrt{a+b z}} - \frac{d \sqrt{a+b z}}{(b c-a d) (c+d z)}$

Rule: If $b c - a d \neq 0 \wedge (a^2 - b^2 = 0 \vee c^2 - d^2 = 0)$, then

$$\int \frac{\sec[e+f x]}{\sqrt{a+b \sec[e+f x]} (c+d \sec[e+f x])} dx \rightarrow \frac{b}{b c - a d} \int \frac{\sec[e+f x]}{\sqrt{a+b \sec[e+f x]}} dx - \frac{d}{b c - a d} \int \frac{\sec[e+f x] \sqrt{a+b \sec[e+f x]}}{c+d \sec[e+f x]} dx$$

Program code:

```
Int[csc[e_.+f_.*x_]/(Sqrt[a+b_.*csc[e_.+f_.*x_]]*(c+d_.*csc[e_.+f_.*x_])),x_Symbol] :=
  b/(b*c-a*d)*Int[Csc[e+f*x]/Sqrt[a+b*Csc[e+f*x]],x] -
  d/(b*c-a*d)*Int[Csc[e+f*x]*Sqrt[a+b*Csc[e+f*x]]/(c+d*Csc[e+f*x]),x] /;
FreeQ[{a,b,c,d,e,f},x] && NeQ[b*c-a*d,0] && (EqQ[a^2-b^2,0] || EqQ[c^2-d^2,0])
```

2: $\int \frac{\sec[e+f x]}{\sqrt{a+b \sec[e+f x]} (c+d \sec[e+f x])} dx$ when $b c - a d \neq 0 \wedge a^2 - b^2 \neq 0 \wedge c^2 - d^2 \neq 0$

Rule: If $b c - a d \neq 0 \wedge a^2 - b^2 \neq 0 \wedge c^2 - d^2 \neq 0$, then

$$\int \frac{\sec[e+f x]}{\sqrt{a+b \sec[e+f x]} (c+d \sec[e+f x])} dx \rightarrow$$

$$\frac{2 \tan[e+f x]}{f (c+d) \sqrt{a+b \sec[e+f x]} \sqrt{-\tan[e+f x]^2}} \sqrt{\frac{a+b \sec[e+f x]}{a+b}} \text{EllipticPi}\left[\frac{2 d}{c+d}, \text{ArcSin}\left[\frac{\sqrt{1-\sec[e+f x]}}{\sqrt{2}}\right], \frac{2 b}{a+b}\right]$$

Program code:

```

Int[csc[e_+f_.*x_]/(Sqrt[a_+b_.*csc[e_+f_.*x_]]*(c_+d_.*csc[e_+f_.*x_])),x_Symbol] :=
-2*Cot[e+f*x]/(f*(c+d)*Sqrt[a+b*Csc[e+f*x]]*Sqrt[-Cot[e+f*x]^2])*Sqrt[(a+b*Csc[e+f*x])/(a+b)]*
EllipticPi[2*d/(c+d),ArcSin[Sqrt[1-Csc[e+f*x]]/Sqrt[2]],2*b/(a+b)] /;
FreeQ[{a,b,c,d,e,f},x] && NeQ[b*c-a*d,0] && NeQ[a^2-b^2,0] && NeQ[c^2-d^2,0]

```

2. $\int \frac{(g \sec[e+f x])^{3/2}}{\sqrt{a+b \sec[e+f x]} (c+d \sec[e+f x])} dx$ when $b c - a d \neq 0$

1: $\int \frac{(g \sec[e+f x])^{3/2}}{\sqrt{a+b \sec[e+f x]} (c+d \sec[e+f x])} dx$ when $b c - a d \neq 0 \wedge a^2 - b^2 = 0$

Derivation: Algebraic expansion

Basis: $\frac{g z}{\sqrt{a+b z}} = -\frac{a g}{(b c - a d) \sqrt{a+b z}} + \frac{c g \sqrt{a+b z}}{(b c - a d) (c+d z)}$

Rule: If $b c - a d \neq 0 \wedge a^2 - b^2 = 0$, then

$$\int \frac{(g \sec[e+f x])^{3/2}}{\sqrt{a+b \sec[e+f x]} (c+d \sec[e+f x])} dx \rightarrow -\frac{a g}{b c - a d} \int \frac{\sqrt{g \sec[e+f x]}}{\sqrt{a+b \sec[e+f x]}} dx + \frac{c g}{b c - a d} \int \frac{\sqrt{g \sec[e+f x]} \sqrt{a+b \sec[e+f x]}}{c+d \sec[e+f x]} dx$$

Program code:

```

Int[(g_.*csc[e_+f_.*x_])^(3/2)/(Sqrt[a_+b_.*csc[e_+f_.*x_]]*(c_+d_.*csc[e_+f_.*x_])),x_Symbol] :=
-a*g/(b*c-a*d)*Int[Sqrt[g*Csc[e+f*x]]/Sqrt[a+b*Csc[e+f*x]],x] +
c*g/(b*c-a*d)*Int[Sqrt[g*Csc[e+f*x]]*Sqrt[a+b*Csc[e+f*x]]/(c+d*Csc[e+f*x]),x] /;
FreeQ[{a,b,c,d,e,f,g},x] && NeQ[b*c-a*d,0] && EqQ[a^2-b^2,0]

```

2: $\int \frac{(g \sec[e+f x])^{3/2}}{\sqrt{a+b \sec[e+f x]} (c+d \sec[e+f x])} dx$ when $b c - a d \neq 0 \wedge a^2 - b^2 \neq 0$

Derivation: Piecewise constant extraction

Basis: $\partial_x \frac{\sqrt{g \sec[e+f x]} \sqrt{b+a \cos[e+f x]}}{\sqrt{a+b \sec[e+f x]}} = 0$

Rule: If $b c - a d \neq 0 \wedge a^2 - b^2 \neq 0$, then

$$\int \frac{(g \sec[e+f x])^{3/2}}{\sqrt{a+b \sec[e+f x]} (c+d \sec[e+f x])} dx \rightarrow \frac{g \sqrt{g \sec[e+f x]} \sqrt{b+a \cos[e+f x]}}{\sqrt{a+b \sec[e+f x]}} \int \frac{1}{\sqrt{b+a \cos[e+f x]} (d+c \cos[e+f x])} dx$$

Program code:

```
Int[(g_.*csc[e_+f_*x_])^(3/2)/(Sqrt[a_+b_.*csc[e_+f_*x_]]*(c_+d_.*csc[e_+f_*x_])),x_Symbol]:=  
g*Sqrt[g*Csc[e+f*x]]*Sqrt[b+a*Sin[e+f*x]]/Sqrt[a+b*Csc[e+f*x]]*Int[1/(Sqrt[b+a*Sin[e+f*x]]*(d+c*Sin[e+f*x])),x]/;  
FreeQ[{a,b,c,d,e,f,g},x] && NeQ[b*c-a*d,0] && NeQ[a^2-b^2,0]
```

3. $\int \frac{\sec[e+f x]^2}{\sqrt{a+b \sec[e+f x]} (c+d \sec[e+f x])} dx$ when $b c - a d \neq 0$

1: $\int \frac{\sec[e+f x]^2}{\sqrt{a+b \sec[e+f x]} (c+d \sec[e+f x])} dx$ when $b c - a d \neq 0 \wedge (a^2 - b^2 = 0 \vee c^2 - d^2 = 0)$

Derivation: Algebraic expansion

Basis: $\frac{z^2}{\sqrt{a+b z} (c+d z)} = -\frac{a z}{(b c-a d) \sqrt{a+b z}} + \frac{c z \sqrt{a+b z}}{(b c-a d) (c+d z)}$

Rule: If $b c - a d \neq 0 \wedge (a^2 - b^2 = 0 \vee c^2 - d^2 = 0)$, then

$$\int \frac{\sec^2[e+f x]}{\sqrt{a+b \sec[e+f x]} (c+d \sec[e+f x])} dx \rightarrow -\frac{a}{b c - a d} \int \frac{\sec[e+f x]}{\sqrt{a+b \sec[e+f x]}} dx + \frac{c}{b c - a d} \int \frac{\sec[e+f x] \sqrt{a+b \sec[e+f x]}}{c+d \sec[e+f x]} dx$$

Program code:

```

Int[csc[e_+f_*x_]^2/(Sqrt[a_+b_.*csc[e_+f_*x_]]*(c_+d_.*csc[e_+f_*x_])),x_Symbol]:=  

-a/(b*c-a*d)*Int[Csc[e+f*x]/Sqrt[a+b*Csc[e+f*x]],x]+  

c/(b*c-a*d)*Int[Csc[e+f*x]*Sqrt[a+b*Csc[e+f*x]]/(c+d*Csc[e+f*x]),x]/;  

FreeQ[{a,b,c,d,e,f},x] && NeQ[b*c-a*d,0] && (EqQ[a^2-b^2,0] || EqQ[c^2-d^2,0])

```

2: $\int \frac{\sec^2[e+f x]}{\sqrt{a+b \sec[e+f x]} (c+d \sec[e+f x])} dx$ when $b c - a d \neq 0 \wedge a^2 - b^2 \neq 0 \wedge c^2 - d^2 \neq 0$

Derivation: Algebraic expansion

Basis: $\frac{z^2}{\sqrt{a+b z} (c+d z)} = \frac{z}{d \sqrt{a+b z}} - \frac{c z}{d \sqrt{a+b z} (c+d z)}$

Rule: If $b c - a d \neq 0 \wedge a^2 - b^2 \neq 0 \wedge c^2 - d^2 \neq 0$, then

$$\int \frac{\sec^2[e+f x]}{\sqrt{a+b \sec[e+f x]} (c+d \sec[e+f x])} dx \rightarrow \frac{1}{d} \int \frac{\sec[e+f x]}{\sqrt{a+b \sec[e+f x]}} dx - \frac{c}{d} \int \frac{\sec[e+f x]}{\sqrt{a+b \sec[e+f x]} (c+d \sec[e+f x])} dx$$

Program code:

```

Int[csc[e_+f_*x_]^2/(Sqrt[a_+b_.*csc[e_+f_*x_]]*(c_+d_.*csc[e_+f_*x_])),x_Symbol]:=  

1/d*Int[Csc[e+f*x]/Sqrt[a+b*Csc[e+f*x]],x]-  

c/d*Int[Csc[e+f*x]/(Sqrt[a+b*Csc[e+f*x]]*(c+d*Csc[e+f*x])),x]/;  

FreeQ[{a,b,c,d,e,f},x] && NeQ[b*c-a*d,0] && NeQ[a^2-b^2,0] && NeQ[c^2-d^2,0]

```

4. $\int \frac{(g \sec[e+f x])^{5/2}}{\sqrt{a+b \sec[e+f x]} (c+d \sec[e+f x])} dx$ when $b c - a d \neq 0$

1: $\int \frac{(g \sec[e+f x])^{5/2}}{\sqrt{a+b \sec[e+f x]} (c+d \sec[e+f x])} dx$ when $b c - a d \neq 0 \wedge a^2 - b^2 = 0$

Derivation: Algebraic expansion

Basis: $\frac{g^2 z^2}{\sqrt{a+b z} (c+d z)} = -\frac{c^2 g^2 \sqrt{a+b z}}{d (b c - a d) (c+d z)} + \frac{g^2 (a c + (b c - a d) z)}{d (b c - a d) \sqrt{a+b z}}$

Rule: If $b c - a d \neq 0 \wedge a^2 - b^2 = 0$, then

$$\int \frac{(g \sec[e+f x])^{5/2}}{\sqrt{a+b \sec[e+f x]} (c+d \sec[e+f x])} dx \rightarrow -\frac{c^2 g^2}{d (b c - a d)} \int \frac{\sqrt{g \sec[e+f x]} \sqrt{a+b \sec[e+f x]}}{c+d \sec[e+f x]} dx + \\ \frac{g^2}{d (b c - a d)} \int \frac{\sqrt{g \sec[e+f x]} (a c + (b c - a d) \sec[e+f x])}{\sqrt{a+b \sec[e+f x]}} dx$$

Program code:

```
Int[(g_.*csc[e_.*f_.*x_])^(5/2)/(Sqrt[a_+b_.*csc[e_.*f_.*x_]]*(c_+d_.*csc[e_.*f_.*x_])),x_Symbol]:=\\
-c^2*g^2/(d*(b*c-a*d))*Int[Sqrt[g*Csc[e+f*x]]*Sqrt[a+b*Csc[e+f*x]]/(c+d*Csc[e+f*x]),x]+\\
g^2/(d*(b*c-a*d))*Int[Sqrt[g*Csc[e+f*x]]*(a*c+(b*c-a*d)*Csc[e+f*x])/Sqrt[a+b*Csc[e+f*x]],x];\\
FreeQ[{a,b,c,d,e,f,g},x] && NeQ[b*c-a*d,0] && EqQ[a^2-b^2,0]
```

2: $\int \frac{(g \sec[e+f x])^{5/2}}{\sqrt{a+b \sec[e+f x]} (c+d \sec[e+f x])} dx$ when $b c - a d \neq 0 \wedge a^2 - b^2 \neq 0$

Derivation: Algebraic expansion

Basis: $\frac{g z}{c+d z} = \frac{g}{d} - \frac{c g}{d (c+d z)}$

Rule: If $b c - a d \neq 0 \wedge a^2 - b^2 \neq 0$, then

$$\int \frac{(g \sec[e+f x])^{5/2}}{\sqrt{a+b \sec[e+f x]} (c+d \sec[e+f x])} dx \rightarrow \frac{g}{d} \int \frac{(g \sec[e+f x])^{3/2}}{\sqrt{a+b \sec[e+f x]}} dx - \frac{c g}{d} \int \frac{(g \sec[e+f x])^{3/2}}{\sqrt{a+b \sec[e+f x]} (c+d \sec[e+f x])} dx$$

Program code:

```
Int[(g_.*csc[e_.*f_.*x_])^(5/2)/(Sqrt[a_+b_.*csc[e_.*f_.*x_]]*(c_+d_.*csc[e_.*f_.*x_])),x_Symbol]:=  
g/d*Int[(g*Csc[e+f*x])^(3/2)/Sqrt[a+b*Csc[e+f*x]],x]-  
c*g/d*Int[(g*Csc[e+f*x])^(3/2)/(Sqrt[a+b*Csc[e+f*x]]*(c+d*Csc[e+f*x])),x];  
FreeQ[{a,b,c,d,e,f,g},x] && NeQ[b*c-a*d,0] && NeQ[a^2-b^2,0]
```

3. $\int \frac{\sec[e+f x]^p (a+b \sec[e+f x])^m}{\sqrt{c+d \sec[e+f x]}} dx$ when $b c - a d \neq 0 \wedge m^2 = \frac{1}{4}$

1. $\int \frac{\sec[e+f x] \sqrt{a+b \sec[e+f x]}}{\sqrt{c+d \sec[e+f x]}} dx$ when $b c - a d \neq 0$

1: $\int \frac{\sec[e+f x] \sqrt{a+b \sec[e+f x]}}{\sqrt{c+d \sec[e+f x]}} dx$ when $b c - a d \neq 0 \wedge a^2 - b^2 = 0 \wedge c^2 - d^2 \neq 0$

Derivation: Integration by substitution

Basis: If $a^2 - b^2 = 0 \wedge c^2 - d^2 \neq 0$, then

$$\frac{\sec[e+f x] \sqrt{a+b \sec[e+f x]}}{\sqrt{c+d \sec[e+f x]}} = \frac{2b}{f} \text{Subst}\left[\frac{1}{1-b d x^2}, x, \frac{\tan[e+f x]}{\sqrt{a+b \sec[e+f x]} \sqrt{c+d \sec[e+f x]}}\right] \partial_x \frac{\tan[e+f x]}{\sqrt{a+b \sec[e+f x]} \sqrt{c+d \sec[e+f x]}}$$

Rule: If $b c - a d \neq 0 \wedge a^2 - b^2 = 0 \wedge c^2 - d^2 \neq 0$, then

$$\int \frac{\sec[e+f x] \sqrt{a+b \sec[e+f x]}}{\sqrt{c+d \sec[e+f x]}} dx \rightarrow \frac{2b}{f} \text{Subst}\left[\int \frac{1}{1-b d x^2} dx, x, \frac{\tan[e+f x]}{\sqrt{a+b \sec[e+f x]} \sqrt{c+d \sec[e+f x]}}\right]$$

Program code:

```
Int[csc[e_+f_*x_]*Sqrt[a_+b_.*csc[e_+f_.*x_]]/Sqrt[c_+d_.*csc[e_+f_.*x_]],x_Symbol] :=
-2*b/f*Subst[Int[1/(1-b*d*x^2),x],x,Cot[e+f*x]/(Sqrt[a+b*Csc[e+f*x]]*Sqrt[c+d*Csc[e+f*x]])] /;
FreeQ[{a,b,c,d,e,f},x] && NeQ[b*c-a*d,0] && EqQ[a^2-b^2,0] && NeQ[c^2-d^2,0]
```

2: $\int \frac{\sec[e+f x] \sqrt{a+b \sec[e+f x]}}{\sqrt{c+d \sec[e+f x]}} dx$ when $b c - a d \neq 0 \wedge a^2 - b^2 \neq 0 \wedge c^2 - d^2 = 0$

Derivation: Algebraic expansion

Basis: $\frac{\sqrt{a+b z}}{\sqrt{c+d z}} = -\frac{b c - a d}{d \sqrt{a+b z} \sqrt{c+d z}} + \frac{b \sqrt{c+d z}}{d \sqrt{a+b z}}$

Rule: If $b c - a d \neq 0 \wedge a^2 - b^2 \neq 0 \wedge c^2 - d^2 = 0$, then

$$\int \frac{\sec[e+f x] \sqrt{a+b \sec[e+f x]}}{\sqrt{c+d \sec[e+f x]}} dx \rightarrow -\frac{b c - a d}{d} \int \frac{\sec[e+f x]}{\sqrt{a+b \sec[e+f x]} \sqrt{c+d \sec[e+f x]}} dx + \frac{b}{d} \int \frac{\sec[e+f x] \sqrt{c+d \sec[e+f x]}}{\sqrt{a+b \sec[e+f x]}} dx$$

Program code:

```
Int[csc[e_+f_*x_]*Sqrt[a_+b_.*csc[e_+f_.*x_]]/Sqrt[c_+d_.*csc[e_+f_.*x_]],x_Symbol] :=
-(b*c-a*d)/d*Int[Csc[e+f*x]/(Sqrt[a+b*Csc[e+f*x]]*Sqrt[c+d*Csc[e+f*x]]),x] +
b/d*Int[Csc[e+f*x]*Sqrt[c+d*Csc[e+f*x]]/Sqrt[a+b*Csc[e+f*x]],x] /;
FreeQ[{a,b,c,d,e,f},x] && NeQ[b*c-a*d,0] && NeQ[a^2-b^2,0] && EqQ[c^2-d^2,0]
```

$$3: \int \frac{\sec[e+f x] \sqrt{a+b \sec[e+f x]}}{\sqrt{c+d \sec[e+f x]}} dx \text{ when } b c - a d \neq 0 \wedge a^2 - b^2 \neq 0 \wedge c^2 - d^2 \neq 0$$

Rule: If $b c - a d \neq 0 \wedge a^2 - b^2 \neq 0 \wedge c^2 - d^2 \neq 0$, then

$$\begin{aligned} & \int \frac{\sec[e+f x] \sqrt{a+b \sec[e+f x]}}{\sqrt{c+d \sec[e+f x]}} dx \rightarrow \\ & \frac{2(a+b \sec[e+f x])}{d f \sqrt{\frac{a+b}{c+d}} \tan[e+f x]} \sqrt{-\frac{(b c - a d) (1 - \sec[e+f x])}{(c + d) (a + b \sec[e+f x])}} \\ & \sqrt{\frac{(b c - a d) (1 + \sec[e+f x])}{(c - d) (a + b \sec[e+f x])}} \operatorname{EllipticPi}\left[\frac{b (c + d)}{d (a + b)}, \operatorname{ArcSin}\left[\sqrt{\frac{a + b}{c + d}} \frac{\sqrt{c + d \sec[e+f x]}}{\sqrt{a + b \sec[e+f x]}}\right], \frac{(a - b) (c + d)}{(a + b) (c - d)}\right] \end{aligned}$$

Program code:

```
Int[csc[e_+f_*x_]*Sqrt[a_+b_.*csc[e_+f_*x_]]/Sqrt[c_+d_.*csc[e_+f_*x_]],x_Symbol]:=  
-2*(a+b*Csc[e+f*x])/((d*f*Sqrt[(a+b)/(c+d)]*Cot[e+f*x])*  
Sqrt[-(b*c-a*d)*(1-Csc[e+f*x])/((c+d)*(a+b*Csc[e+f*x]))]*Sqrt[(b*c-a*d)*(1+Csc[e+f*x])/((c-d)*(a+b*Csc[e+f*x]))]*  
EllipticPi[b*(c+d)/(d*(a+b)),ArcSin[Sqrt[(a+b)/(c+d)]*Sqrt[c+d*Csc[e+f*x]]/Sqrt[a+b*Csc[e+f*x]]],(a-b)*(c+d)/((a+b)*(c-d))];  
FreeQ[{a,b,c,d,e,f},x] && NeQ[b*c-a*d,0] && NeQ[a^2-b^2,0] && NeQ[c^2-d^2,0]
```

$$2. \int \frac{\sec[e+f x]}{\sqrt{a+b \sec[e+f x]} \sqrt{c+d \sec[e+f x]}} dx \text{ when } b c - a d \neq 0$$

$$1: \int \frac{\sec[e+f x]}{\sqrt{a+b \sec[e+f x]} \sqrt{c+d \sec[e+f x]}} dx \text{ when } b c - a d \neq 0 \wedge a^2 - b^2 = 0 \wedge c^2 - d^2 \neq 0$$

Derivation: Integration by substitution

Basis: If $a^2 - b^2 = 0 \wedge c^2 - d^2 \neq 0$, then

$$\frac{\sec[e+f x]}{\sqrt{a+b \sec[e+f x]} \sqrt{c+d \sec[e+f x]}} = \\ \frac{2a}{bf} \text{Subst} \left[\frac{1}{2 + (a c - b d) x^2}, x, \frac{\tan[e+f x]}{\sqrt{a+b \sec[e+f x]} \sqrt{c+d \sec[e+f x]}} \right] \partial_x \frac{\tan[e+f x]}{\sqrt{a+b \sec[e+f x]} \sqrt{c+d \sec[e+f x]}}$$

Rule: If $b c - a d \neq 0 \wedge a^2 - b^2 = 0 \wedge c^2 - d^2 \neq 0$, then

$$\int \frac{\sec[e+f x]}{\sqrt{a+b \sec[e+f x]} \sqrt{c+d \sec[e+f x]}} dx \rightarrow \frac{2a}{bf} \text{Subst} \left[\int \frac{1}{2 + (a c - b d) x^2} dx, x, \frac{\tan[e+f x]}{\sqrt{a+b \sec[e+f x]} \sqrt{c+d \sec[e+f x]}} \right]$$

Program code:

```
Int[csc[e_.+f_.*x_]/(Sqrt[a_+b_.*csc[e_.+f_.*x_]]*Sqrt[c_+d_.*csc[e_.+f_.*x_]]),x_Symbol] :=  
-2*a/(b*f)*Subst[Int[1/(2+(a*c-b*d)*x^2),x],x,Cot[e+f*x]/(Sqrt[a+b*Csc[e+f*x]]*Sqrt[c+d*Csc[e+f*x]])] /;  
FreeQ[{a,b,c,d,e,f},x] && NeQ[b*c-a*d,0] && EqQ[a^2-b^2,0] && NeQ[c^2-d^2,0]
```

2: $\int \frac{\sec[e+f x]}{\sqrt{a+b \sec[e+f x]} \sqrt{c+d \sec[e+f x]}} dx$ when $b c - a d \neq 0 \wedge a^2 - b^2 \neq 0 \wedge c^2 - d^2 \neq 0$

Rule: If $b c - a d \neq 0 \wedge a^2 - b^2 \neq 0 \wedge c^2 - d^2 \neq 0$, then

$$\int \frac{\sec[e+f x]}{\sqrt{a+b \sec[e+f x]} \sqrt{c+d \sec[e+f x]}} dx \rightarrow$$

$$\frac{2 (c+d \sec[e+f x])}{f (b c - a d) \sqrt{\frac{c+d}{a+b}} \tan[e+f x]} \sqrt{\frac{(b c - a d) (1 - \sec[e+f x])}{(a+b) (c+d \sec[e+f x])}}$$

$$\sqrt{-\frac{(b c - a d) (1 + \sec[e+f x])}{(a-b) (c+d \sec[e+f x])}} \text{EllipticF}[\text{ArcSin}\left[\sqrt{\frac{c+d}{a+b}} \frac{\sqrt{a+b \sec[e+f x]}}{\sqrt{c+d \sec[e+f x]}}\right], \frac{(a+b) (c-d)}{(a-b) (c+d)}]$$

Program code:

```
Int[csc[e_.+f_.*x_]/(Sqrt[a_.+b_.*csc[e_.+f_.*x_]]*Sqrt[c_.+d_.*csc[e_.+f_.*x_]]),x_Symbol]:=
-2*(c+d*Csc[e+f*x])/((f*(b*c-a*d)*Rt[(c+d)/(a+b),2]*Cot[e+f*x])*_
Sqrt[(b*c-a*d)*(1-Csc[e+f*x])/((a+b)*(c+d*Csc[e+f*x]))]*Sqrt[-(b*c-a*d)*(1+Csc[e+f*x])/((a-b)*(c+d*Csc[e+f*x]))]*_
EllipticF[ArcSin[Rt[(c+d)/(a+b),2]*(Sqrt[a+b*Csc[e+f*x]]/Sqrt[c+d*Csc[e+f*x]])],(a+b)*(c-d)/((a-b)*(c+d))]];
FreeQ[{a,b,c,d,e,f},x] && NeQ[b*c-a*d,0] && NeQ[a^2-b^2,0] && NeQ[c^2-d^2,0]
```

3: $\int \frac{\sec^2[e+f x]}{\sqrt{a+b \sec[e+f x]} \sqrt{c+d \sec[e+f x]}} dx$ when $b c - a d \neq 0$

Derivation: Algebraic expansion

Basis: $\frac{z}{\sqrt{a+b z}} = -\frac{a}{b \sqrt{a+b z}} + \frac{\sqrt{a+b z}}{b}$

Rule: If $b c - a d \neq 0$, then

$$\int \frac{\sec^2[e+f x]}{\sqrt{a+b \sec[e+f x]} \sqrt{c+d \sec[e+f x]}} dx \rightarrow -\frac{a}{b} \int \frac{\sec[e+f x]}{\sqrt{a+b \sec[e+f x]} \sqrt{c+d \sec[e+f x]}} dx + \frac{1}{b} \int \frac{\sec[e+f x] \sqrt{a+b \sec[e+f x]}}{\sqrt{c+d \sec[e+f x]}} dx$$

Program code:

```
Int[csc[e_.+f_.*x_]^2/(Sqrt[a_.+b_.*csc[e_.+f_.*x_]]*Sqrt[c_.+d_.*csc[e_.+f_.*x_]]),x_Symbol]:=  
-a/b*Int[Csc[e+f*x]/(Sqrt[a+b*Csc[e+f*x]]*Sqrt[c+d*Csc[e+f*x]]),x]+  
1/b*Int[Csc[e+f*x]*Sqrt[a+b*Csc[e+f*x]]/Sqrt[c+d*Csc[e+f*x]],x];  
FreeQ[{a,b,c,d,e,f},x] && NeQ[b*c-a*d,0]
```

4: $\int \frac{\sec[e+f x] \sqrt{a+b \sec[e+f x]}}{(c+d \sec[e+f x])^{3/2}} dx$ when $b c - a d \neq 0 \wedge a^2 - b^2 \neq 0 \wedge c^2 - d^2 \neq 0$

Derivation: Algebraic expansion

Basis: $\frac{\sqrt{a+b z}}{c+d z} = \frac{a-b}{(c-d) \sqrt{a+b z}} + \frac{(b c-a d) (1+z)}{(c-d) \sqrt{a+b z} (c+d z)}$

Rule: If $b c - a d \neq 0 \wedge a^2 - b^2 \neq 0 \wedge c^2 - d^2 \neq 0$, then

$$\int \frac{\sec[e+f x] \sqrt{a+b \sec[e+f x]}}{(c+d \sec[e+f x])^{3/2}} dx \rightarrow$$

$$\frac{a - b}{c - d} \int \frac{\sec[e + f x]}{\sqrt{a + b \sec[e + f x]} \sqrt{c + d \sec[e + f x]}} dx + \frac{b c - a d}{c - d} \int \frac{\sec[e + f x] (1 + \sec[e + f x])}{\sqrt{a + b \sec[e + f x]} (c + d \sec[e + f x])^{3/2}} dx$$

Program code:

```

Int[csc[e_+f_*x_]*Sqrt[a_+b_*csc[e_+f_*x_]]/(c_+d_*csc[e_+f_*x_])^(3/2),x_Symbol]:=  

(a-b)/(c-d)*Int[Csc[e+f*x]/(Sqrt[a+b*Csc[e+f*x]]*Sqrt[c+d*Csc[e+f*x]]),x]+  

(b*c-a*d)/(c-d)*Int[Csc[e+f*x]*(1+Csc[e+f*x])/((Sqrt[a+b*Csc[e+f*x]]*(c+d*Csc[e+f*x])^(3/2)),x)];  

FreeQ[{a,b,c,d,e,f},x] && NeQ[b*c-a*d,0] && NeQ[a^2-b^2,0] && NeQ[c^2-d^2,0]

```

5: $\int (g \sec[e+f x])^p (a+b \sec[e+f x])^m (c+d \sec[e+f x])^n dx$ when $b c - a d \neq 0 \wedge a^2 - b^2 = 0 \wedge c^2 - d^2 \neq 0 \wedge (p = 1 \vee m - \frac{1}{2} \in \mathbb{Z})$

Derivation: Piecewise constant extraction and integration by substitution

Basis: If $a^2 - b^2 = 0$, then $\partial_x \frac{\tan[e+f x]}{\sqrt{a+b \sec[e+f x]} \sqrt{a-b \sec[e+f x]}} = 0$

Basis: If $a^2 - b^2 = 0$, then $-\frac{a^2 \tan[e+f x]}{\sqrt{a+b \sec[e+f x]} \sqrt{a-b \sec[e+f x]}} \frac{\tan[e+f x]}{\sqrt{a+b \sec[e+f x]} \sqrt{a-b \sec[e+f x]}} = 1$

Basis: $\tan[e+f x] F[\sec[e+f x]] = \frac{1}{f} \text{Subst}\left[\frac{F[x]}{x}, x, \sec[e+f x]\right] \partial_x \sec[e+f x]$

Rule: If $b c - a d \neq 0 \wedge a^2 - b^2 = 0 \wedge c^2 - d^2 \neq 0 \wedge (p = 1 \vee m - \frac{1}{2} \in \mathbb{Z})$, then

$$\int (g \sec[e+f x])^p (a+b \sec[e+f x])^m (c+d \sec[e+f x])^n dx \rightarrow$$

$$-\frac{a^2 \tan[e+f x]}{\sqrt{a+b \sec[e+f x]} \sqrt{a-b \sec[e+f x]}} \int \frac{\tan[e+f x] (g \sec[e+f x])^p (a+b \sec[e+f x])^{m-\frac{1}{2}} (c+d \sec[e+f x])^n}{\sqrt{a-b \sec[e+f x]}} dx \rightarrow$$

$$-\frac{a^2 g \tan[e+f x]}{f \sqrt{a+b \sec[e+f x]} \sqrt{a-b \sec[e+f x]}} \text{Subst}\left[\int \frac{(g x)^{p-1} (a+b x)^{m-\frac{1}{2}} (c+d x)^n}{\sqrt{a-b x}} dx, x, \sec[e+f x]\right]$$

Program code:

```
Int[(g_.*csc[e_.*f_.*x_])^p_.*(a_+b_.*csc[e_.*f_.*x_])^m_.*(c_+d_.*csc[e_.*f_.*x_])^n_,x_Symbol]:=  
a^2*g*Cot[e+f*x]/(f*Sqrt[a+b*Csc[e+f*x]]*Sqrt[a-b*Csc[e+f*x]])*  
Subst[Int[(g*x)^(p-1)*(a+b*x)^(m-1/2)*(c+d*x)^n/Sqrt[a-b*x],x],x,Csc[e+f*x]] /;  
FreeQ[{a,b,c,d,e,f,g,m,n,p},x] && NeQ[b*c-a*d,0] && EqQ[a^2-b^2,0] && NeQ[c^2-d^2,0] && (EqQ[p,1] || IntegerQ[m-1/2])
```

6: $\int (g \sec[e + f x])^p (a + b \sec[e + f x])^m (c + d \sec[e + f x])^n dx$ when $b c - a d \neq 0 \wedge m \in \mathbb{Z} \wedge n \in \mathbb{Z}$

Derivation: Algebraic normalization

Basis: If $m \in \mathbb{Z} \wedge n \in \mathbb{Z}$, then $(a + b \sec[z])^m (c + d \sec[z])^n = \sec[z]^{m+n} (b + a \cos[z])^m (d + c \cos[z])^n$

Rule: If $b c - a d \neq 0 \wedge m \in \mathbb{Z} \wedge n \in \mathbb{Z}$, then

$$\int (g \sec[e + f x])^p (a + b \sec[e + f x])^m (c + d \sec[e + f x])^n dx \rightarrow \frac{1}{g^{m+n}} \int (g \sec[e + f x])^{m+n+p} (b + a \cos[e + f x])^m (d + c \cos[e + f x])^n dx$$

Program code:

```
Int[(g_.*csc[e_._+f_._*x_])^p_.*(a_._+b_._*csc[e_._+f_._*x_])^m_.*(c_._+d_._*csc[e_._+f_._*x_])^n_,x_Symbol]:=  
1/g^(m+n)*Int[(g*Csc[e+f*x])^(m+n+p)*(b+a*Sin[e+f*x])^m*(d+c*Sin[e+f*x])^n,x]/;  
FreeQ[{a,b,c,d,e,f,g,p},x] && NeQ[b*c-a*d,0] && IntegerQ[m] && IntegerQ[n]
```

7. $\int (g \sec[e + f x])^p (a + b \sec[e + f x])^m (c + d \sec[e + f x])^n dx$ when $b c - a d \neq 0 \wedge m + n + p = 0$

1: $\int (g \sec[e + f x])^p (a + b \sec[e + f x])^m (c + d \sec[e + f x])^n dx$ when $b c - a d \neq 0 \wedge m + n + p = 0 \wedge m \in \mathbb{Z}$

Derivation: Algebraic normalization and piecewise constant extraction

Basis: $a + b \sec[e + f x] = \sec[e + f x] (b + a \cos[e + f x])$

Basis: If $m + n + p = 0$, then $\partial_x \frac{(g \sec[e + f x])^{m+p} (c + d \sec[e + f x])^n}{(d + c \cos[e + f x])^n} = 0$

Rule: If $b c - a d \neq 0 \wedge m + n + p = 0 \wedge m \in \mathbb{Z}$, then

$$\int (g \sec[e + f x])^p (a + b \sec[e + f x])^m (c + d \sec[e + f x])^n dx \rightarrow \frac{1}{g^m} \int (g \sec[e + f x])^{m+p} (b + a \cos[e + f x])^m (c + d \sec[e + f x])^n dx$$

$$\rightarrow \frac{(g \sec[e+f x])^{m+p} (c+d \sec[e+f x])^n}{g^m (d+c \cos[e+f x])^n} \int (b+a \cos[e+f x])^m (d+c \cos[e+f x])^n dx$$

Program code:

```
Int[(g_.*csc[e_.*f_.*x_])^p_.*(a_+b_.*csc[e_.*f_.*x_])^m_.*(c_+d_.*csc[e_.*f_.*x_])^n_,x_Symbol]:=  
    (g*Csc[e+f*x])^(m+p)*(c+d*Csc[e+f*x])^n/(g^m*(d+c*Sin[e+f*x])^n)*Int[(b+a*Sin[e+f*x])^m*(d+c*Sin[e+f*x])^n,x]/;  
FreeQ[{a,b,c,d,e,f,g,n,p},x] && NeQ[b*c-a*d,0] && EqQ[m+n+p,0] && IntegerQ[m]
```

2: $\int (g \sec[e+f x])^p (a+b \sec[e+f x])^m (c+d \sec[e+f x])^n dx$ when $b c - a d \neq 0 \wedge m + n + p = 0 \wedge m \notin \mathbb{Z}$

Derivation: Piecewise constant extraction

Basis: If $m + n + p = 0$, then $\partial_x \frac{(g \sec[e+f x])^p (a+b \sec[e+f x])^m (c+d \sec[e+f x])^n}{(b+a \cos[e+f x])^m (d+c \cos[e+f x])^n} = 0$

Rule: If $b c - a d \neq 0 \wedge m + n + p = 0 \wedge m \notin \mathbb{Z}$, then

$$\int (g \sec[e+f x])^p (a+b \sec[e+f x])^m (c+d \sec[e+f x])^n dx \rightarrow \frac{(g \sec[e+f x])^p (a+b \sec[e+f x])^m (c+d \sec[e+f x])^n}{(b+a \cos[e+f x])^m (d+c \cos[e+f x])^n} \int (b+a \cos[e+f x])^m (d+c \cos[e+f x])^n dx$$

Program code:

```
Int[(g_.*csc[e_.*f_.*x_])^p_.*(a_+b_.*csc[e_.*f_.*x_])^m_.*(c_+d_.*csc[e_.*f_.*x_])^n_,x_Symbol]:=  
    (g*Csc[e+f*x])^p*(a+b*Csc[e+f*x])^m*(c+d*Csc[e+f*x])^n/((b+a*Sin[e+f*x])^m*(d+c*Sin[e+f*x])^n)*  
    Int[(b+a*Sin[e+f*x])^m*(d+c*Sin[e+f*x])^n,x]/;  
FreeQ[{a,b,c,d,e,f,g,m,n,p},x] && NeQ[b*c-a*d,0] && EqQ[m+n+p,0] && Not[IntegerQ[m]]
```

8: $\int \sec[e+f x]^p (a+b \sec[e+f x])^m (c+d \sec[e+f x])^n dx$ when $b c - a d \neq 0 \wedge m - \frac{1}{2} \in \mathbb{Z} \wedge n - \frac{1}{2} \in \mathbb{Z} \wedge p \in \mathbb{Z}$

Derivation: Piecewise constant extraction

Basis: $\partial_x \frac{\sqrt{d+c \cos[e+f x]}}{\sqrt{b+a \cos[e+f x]}} \frac{\sqrt{a+b \sec[e+f x]}}{\sqrt{c+d \sec[e+f x]}} = 0$

Note: The restriction $m + n + p \in \{-1, -2\}$ can be lifted if and when the cosine integration rules are extended to handle integrands of the form $\cos[e+f x]^p (a+b \cos[e+f x])^m (c+d \cos[e+f x])^n$ for arbitrary p .

Rule: If $b c - a d \neq 0 \wedge m - \frac{1}{2} \in \mathbb{Z} \wedge n - \frac{1}{2} \in \mathbb{Z} \wedge p \in \mathbb{Z}$, then

$$\int \sec[e+f x]^p (a+b \sec[e+f x])^m (c+d \sec[e+f x])^n dx \rightarrow \frac{\sqrt{d+c \cos[e+f x]}}{\sqrt{b+a \cos[e+f x]}} \frac{\sqrt{a+b \sec[e+f x]}}{\sqrt{c+d \sec[e+f x]}} \int \frac{(b+a \cos[e+f x])^m (d+c \cos[e+f x])^n}{\cos[e+f x]^{m+n+p}} dx$$

Program code:

```
Int[csc[e_..+f_..*x_]^p.*(a_+b_.*csc[e_..+f_..*x_])^m.*(c_+d_.*csc[e_..+f_..*x_])^n_,x_Symbol]:=  
  Sqrt[d+c*Sin[e+f*x]]*Sqrt[a+b*Csc[e+f*x]]/(Sqrt[b+a*Sin[e+f*x]]*Sqrt[c+d*Csc[e+f*x]])*  
  Int[(b+a*Sin[e+f*x])^m*(d+c*Sin[e+f*x])^n/Sin[e+f*x]^(m+n+p),x];;  
FreeQ[{a,b,c,d,e,f,m,n},x] && NeQ[b*c-a*d,0] && IntegerQ[m-1/2] && IntegerQ[n-1/2] && IntegerQ[p] && LeQ[-2,m+n+p,-1]
```

9: $\int (g \sec[e+f x])^p (a+b \sec[e+f x])^m (c+d \sec[e+f x])^n dx$ when $b c - a d \neq 0 \wedge ((m+n) \in \mathbb{Z} \vee (m+p) \in \mathbb{Z} \vee (n+p) \in \mathbb{Z})$

Derivation: Algebraic expansion

Rule: If $b c - a d \neq 0 \wedge ((m+n) \in \mathbb{Z} \vee (m+p) \in \mathbb{Z} \vee (n+p) \in \mathbb{Z})$, then

$$\int (g \sec[e+f x])^p (a+b \sec[e+f x])^m (c+d \sec[e+f x])^n dx \rightarrow$$

$$\int \text{ExpandTrig}[(g \sec[e+f x])^p (a+b \sec[e+f x])^m (c+d \sec[e+f x])^n, x] dx$$

Program code:

```
Int[(g_.*csc[e_.*f_.*x_])^p_.*(a_+b_.*csc[e_.*f_.*x_])^m_*(c_+d_.*csc[e_.*f_.*x_])^n_,x_Symbol]:=  
Int[ExpandTrig[(g*csc[e+f*x])^p*(a+b*csc[e+f*x])^m*(c+d*csc[e+f*x])^n,x],x]/;  
FreeQ[{a,b,c,d,e,f,g,m,n,p},x] && NeQ[b*c-a*d,0] && (IntegersQ[m,n] || IntegersQ[m,p] || IntegersQ[n,p])
```

x: $\int (g \sec[e+f x])^p (a+b \sec[e+f x])^m (c+d \sec[e+f x])^n dx$

Rule:

$$\int (g \sec[e+f x])^p (a+b \sec[e+f x])^m (c+d \sec[e+f x])^n dx \rightarrow \int (g \sec[e+f x])^p (a+b \sec[e+f x])^m (c+d \sec[e+f x])^n dx$$

Program code:

```
Int[(g_.*csc[e_.*f_.*x_])^p_.*(a_+b_.*csc[e_.*f_.*x_])^m_*(c_+d_.*csc[e_.*f_.*x_])^n_,x_Symbol]:=  
Unintegrable[(g*Csc[e+f*x])^p*(a+b*Csc[e+f*x])^m*(c+d*Csc[e+f*x])^n,x]/;  
FreeQ[{a,b,c,d,e,f,g,m,n,p},x]
```

Rules for integrands of the form $(g \sec[e+f x])^p (a+b \sec[e+f x])^m (c+d \sec[e+f x])^n (A+B \sec[e+f x])$

1: $\int \frac{\sec[e+f x] (A+B \sec[e+f x])}{\sqrt{a+b \sec[e+f x]} (c+d \sec[e+f x])^{3/2}} dx$ when $b c - a d \neq 0 \wedge a^2 - b^2 \neq 0 \wedge c^2 - d^2 \neq 0 \wedge A == B$

Rule: If $b c - a d \neq 0 \wedge a^2 - b^2 \neq 0 \wedge c^2 - d^2 \neq 0 \wedge A == B$, then

$$\int \frac{\sec[e+f x] (A+B \sec[e+f x])}{\sqrt{a+b \sec[e+f x]} (c+d \sec[e+f x])^{3/2}} dx \rightarrow$$

$$\frac{2 A \left(1 + \sec[e + f x]\right) \sqrt{\frac{(b c - a d) (1 - \sec[e + f x])}{(a + b) (c + d \sec[e + f x])}}}{f (b c - a d) \sqrt{\frac{c + d}{a + b}} \tan[e + f x] \sqrt{-\frac{(b c - a d) (1 + \sec[e + f x])}{(a - b) (c + d \sec[e + f x])}}} \text{EllipticE}\left[\text{ArcSin}\left[\sqrt{\frac{c + d}{a + b}} \frac{\sqrt{a + b \sec[e + f x]}}{\sqrt{c + d \sec[e + f x]}}\right], \frac{(a + b) (c - d)}{(a - b) (c + d)}\right]$$

Program code:

```
Int[sec[e_+f_.*x_]*(A_+B_.*sec[e_+f_.*x_])/({Sqrt[a_+b_.*sec[e_+f_.*x_]]*(c_+d_.*sec[e_+f_.*x_])^(3/2)},x_Symbol]:=2*A*(1+Sec[e+f*x])*Sqrt[(b*c-a*d)*(1-Sec[e+f*x])/((a+b)*(c+d*Sec[e+f*x]))]/(f*(b*c-a*d)*Rt[(c+d)/(a+b),2]*Tan[e+f*x]*Sqrt[-(b*c-a*d)*(1+Sec[e+f*x])/((a-b)*(c+d*Sec[e+f*x]))])*EllipticE[ArcSin[Rt[(c+d)/(a+b),2]*Sqrt[a+b*Sec[e+f*x]]/Sqrt[c+d*Sec[e+f*x]]],(a+b)*(c-d)/((a-b)*(c+d))];FreeQ[{a,b,c,d,e,f,A,B},x] && NeQ[b*c-a*d,0] && NeQ[a^2-b^2,0] && NeQ[c^2-d^2,0] && EqQ[A,B]
```

```
Int[csc[e_+f_.*x_]*(A_+B_.*csc[e_+f_.*x_])/({Sqrt[a_+b_.*csc[e_+f_.*x_]]*(c_+d_.*csc[e_+f_.*x_])^(3/2)},x_Symbol):=-2*A*(1+Csc[e+f*x])*Sqrt[(b*c-a*d)*(1-Csc[e+f*x])/((a+b)*(c+d*Csc[e+f*x]))]/(f*(b*c-a*d)*Rt[(c+d)/(a+b),2]*Cot[e+f*x]*Sqrt[-(b*c-a*d)*(1+Csc[e+f*x])/((a-b)*(c+d*Csc[e+f*x]))])*EllipticE[ArcSin[Rt[(c+d)/(a+b),2]*Sqrt[a+b*Csc[e+f*x]]/Sqrt[c+d*Csc[e+f*x]]],(a+b)*(c-d)/((a-b)*(c+d))];FreeQ[{a,b,c,d,e,f,A,B},x] && NeQ[b*c-a*d,0] && NeQ[a^2-b^2,0] && NeQ[c^2-d^2,0] && EqQ[A,B]
```