

Rules for integrands of the form $(g \tan[e + f x])^p (a + b \tan[e + f x])^m (c + d \tan[e + f x])^n$

x: $\int (g \tan[e + f x])^p (a + b \tan[e + f x])^m (c + d \tan[e + f x])^n dx$

Rule:

$$\int (g \tan[e + f x])^p (a + b \tan[e + f x])^m (c + d \tan[e + f x])^n dx \rightarrow \int (g \tan[e + f x])^p (a + b \tan[e + f x])^m (c + d \tan[e + f x])^n dx$$

Program code:

```
Int[(g_.*tan[e_.+f_.*x_])^p_.*(a_.+b_.*tan[e_.+f_.*x_])^m_.*(c_.+d_.*tan[e_.+f_.*x_])^n_,x_Symbol] :=
  Unintegrable[(g*Tan[e+f*x])^p*(a+b*Tan[e+f*x])^m*(c+d*Tan[e+f*x])^n,x] /;
  FreeQ[{a,b,c,d,e,f,g,m,n,p},x]
```

Rules for integrands of the form $(g \tan[e + f x]^q)^p (a + b \tan[e + f x])^m (c + d \tan[e + f x])^n$

1: $\int (g \cot[e + f x])^p (a + b \tan[e + f x])^m (c + d \tan[e + f x])^n dx$ when $p \notin \mathbb{Z} \wedge m \in \mathbb{Z} \wedge n \in \mathbb{Z}$

Derivation: Algebraic normalization

Basis: If $m \in \mathbb{Z} \wedge n \in \mathbb{Z}$, then $(a + b \tan[z])^m (c + d \tan[z])^n = \frac{g^{m+n} (b+a \cot[z])^m (d+c \cot[z])^n}{(g \cot[z])^{m+n}}$

Rule: If $p \notin \mathbb{Z} \wedge m \in \mathbb{Z} \wedge n \in \mathbb{Z}$, then

$$\int (g \cot[e + f x])^p (a + b \tan[e + f x])^m (c + d \tan[e + f x])^n dx \rightarrow g^{m+n} \int (g \cot[e + f x])^{p-m-n} (b + a \cot[e + f x])^m (d + c \cot[e + f x])^n dx$$

Program code:

```
Int[(g_.*cot[e_.+f_.*x_])^p_.*(a_.+b_.*tan[e_.+f_.*x_])^m_.*(c_.+d_.*tan[e_.+f_.*x_])^n_,x_Symbol] :=
  g^(m+n)*Int[(g*Cot[e+f*x])^(p-m-n)*(b+a*Cot[e+f*x])^m*(d+c*Cot[e+f*x])^n,x] /;
  FreeQ[{a,b,c,d,e,f,g,p},x] && Not[IntegerQ[p]] && IntegerQ[m] && IntegerQ[n]
```

```
Int [(g_.*tan[e_.+f_.*x_])^p_*(a_.+b_.*cot[e_.+f_.*x_])^m_.*(c_.+d_.*cot[e_.+f_.*x_])^n_.,x_Symbol] :=
  g^(m+n)*Int [(g*Tan[e+f*x])^(p-m-n)*(b+a*Tan[e+f*x])^m*(d+c*Tan[e+f*x])^n,x] /;
FreeQ[{a,b,c,d,e,f,g,p},x] && Not[IntegerQ[p]] && IntegerQ[m] && IntegerQ[n]
```

2: $\int (g \tan(e+fx))^p (a+b \tan(e+fx))^m (c+d \tan(e+fx))^n dx$ when $p \notin \mathbb{Z} \wedge \neg (m \in \mathbb{Z} \wedge n \in \mathbb{Z})$

Derivation: Piecewise constant extraction

Basis: $\partial_x \frac{(g \tan(e+fx))^p}{(g \tan(e+fx))^{pq}} == 0$

– Rule: If $p \notin \mathbb{Z} \wedge \neg (m \in \mathbb{Z} \wedge n \in \mathbb{Z})$, then

$$\int (g \tan(e+fx))^p (a+b \tan(e+fx))^m (c+d \tan(e+fx))^n dx \rightarrow \frac{(g \tan(e+fx))^p}{(g \tan(e+fx))^{pq}} \int (g \tan(e+fx))^{pq} (a+b \tan(e+fx))^m (c+d \tan(e+fx))^n dx$$

– Program code:

```
Int [(g_.*tan[e_.+f_.*x_]^q_)^p_*(a_.+b_.*tan[e_.+f_.*x_]^m_.*(c_.+d_.*tan[e_.+f_.*x_]^n_.,x_Symbol] :=
  (g*Tan[e+f*x]^q)^p/(g*Tan[e+f*x])^(p*q)*Int [(g*Tan[e+f*x])^(p*q)*(a+b*Tan[e+f*x])^m*(c+d*Tan[e+f*x])^n,x] /;
FreeQ[{a,b,c,d,e,f,g,m,n,p,q},x] && Not[IntegerQ[p]] && Not[IntegerQ[m]] && IntegerQ[n]
```

Rules for integrands of the form $(g \tan[e + f x])^p (a + b \tan[e + f x])^m (c + d \cot[e + f x])^n$

1: $\int (g \tan[e + f x])^p (a + b \tan[e + f x])^m (c + d \cot[e + f x])^n dx$ when $n \in \mathbb{Z}$

Derivation: Algebraic normalization

Basis: $c + d \cot[z] == \frac{d+c \tan[z]}{\tan[z]}$

Rule: If $n \in \mathbb{Z}$, then

$$\int (g \tan[e + f x])^p (a + b \tan[e + f x])^m (c + d \cot[e + f x])^n dx \rightarrow g^n \int (g \tan[e + f x])^{p-n} (a + b \tan[e + f x])^m (d + c \tan[e + f x])^n dx$$

Program code:

```
Int[(g_.*tan[e_.+f_.*x_])^p_.*(a_+b_.*tan[e_.+f_.*x_])^m_.*(c_+d_.*cot[e_.+f_.*x_])^n_.,x_Symbol] :=
  g^n*Int[(g*Tan[e+f*x])^(p-n)*(a+b*Tan[e+f*x])^m*(d+c*Tan[e+f*x])^n,x] /;
FreeQ[{a,b,c,d,e,f,g,m,p},x] && IntegerQ[n]
```

$$2. \int (g \tan[e+fx])^p (a+b \tan[e+fx])^m (c+d \cot[e+fx])^n dx \text{ when } n \notin \mathbb{Z}$$

$$1. \int (g \tan[e+fx])^p (a+b \tan[e+fx])^m (c+d \cot[e+fx])^n dx \text{ when } n \notin \mathbb{Z} \wedge m \in \mathbb{Z}$$

$$1: \int \tan[e+fx]^p (a+b \tan[e+fx])^m (c+d \cot[e+fx])^n dx \text{ when } n \notin \mathbb{Z} \wedge m \in \mathbb{Z} \wedge p \in \mathbb{Z}$$

Derivation: Algebraic normalization

$$\text{Basis: } a + b \tan[z] == \frac{b+a \cot[z]}{\cot[z]}$$

Rule: If $n \notin \mathbb{Z} \wedge m \in \mathbb{Z} \wedge p \in \mathbb{Z}$, then

$$\int \tan[e+fx]^p (a+b \tan[e+fx])^m (c+d \cot[e+fx])^n dx \rightarrow \int \frac{(b+a \cot[e+fx])^m (c+d \cot[e+fx])^n}{\cot[e+fx]^{m+p}} dx$$

Program code:

```
Int[tan[e_+f_*x_]^p_.*(a_+b_*tan[e_+f_*x_]^m_.*(c_+d_*cot[e_+f_*x_]^n_,x_Symbol)] :=
  Int[(b+a*Cot[e+f*x])^m*(c+d*Cot[e+f*x])^n/Cot[e+f*x]^(m+p),x] /;
  FreeQ[{a,b,c,d,e,f,n},x] && Not[IntegerQ[n]] && IntegerQ[m] && IntegerQ[p]
```

$$2: \int (g \tan[e+fx])^p (a+b \tan[e+fx])^m (c+d \cot[e+fx])^n dx \text{ when } n \notin \mathbb{Z} \wedge m \in \mathbb{Z} \wedge p \notin \mathbb{Z}$$

Derivation: Algebraic normalization and piecewise constant extraction

$$\text{Basis: } a + b \tan[z] == \frac{b+a \cot[z]}{\cot[z]}$$

$$\text{Basis: } \partial_x (\cot[e+fx]^p (g \tan[e+fx])^p) == 0$$

Rule: If $n \notin \mathbb{Z} \wedge m \in \mathbb{Z} \wedge p \notin \mathbb{Z}$, then

$$\int (g \tan[e+fx])^p (a+b \tan[e+fx])^m (c+d \cot[e+fx])^n dx \rightarrow \cot[e+fx]^p (g \tan[e+fx])^p \int \frac{(b+a \cot[e+fx])^m (c+d \cot[e+fx])^n}{\cot[e+fx]^{m+p}} dx$$

Program code:

```
Int [(g_*tan[e_+f_*x_])^p_*(a_+b_*tan[e_+f_*x_])^m_*(c_+d_*cot[e_+f_*x_])^n_,x_Symbol] :=
  Cot[e+f*x]^p*(g*Tan[e+f*x])^p*Int [(b+a*Cot[e+f*x])^m*(c+d*Cot[e+f*x])^n/Cot[e+f*x]^(m+p),x] /;
FreeQ[{a,b,c,d,e,f,g,n,p},x] && Not[IntegerQ[n]] && IntegerQ[m] && Not[IntegerQ[p]]
```

2: $\int (g \tan[e+fx])^p (a+b \tan[e+fx])^m (c+d \cot[e+fx])^n dx$ when $n \notin \mathbb{Z} \wedge m \notin \mathbb{Z}$

Derivation: Piecewise constant extraction

Basis: $\partial_x \frac{(c+d \cot[e+fx])^n (g \tan[e+fx])^n}{(d+c \tan[e+fx])^n} == 0$

Rule: If $n \notin \mathbb{Z} \wedge m \notin \mathbb{Z}$, then

$$\int (g \tan[e+fx])^p (a+b \tan[e+fx])^m (c+d \cot[e+fx])^n dx \rightarrow \frac{(g \tan[e+fx])^n (c+d \cot[e+fx])^n}{(d+c \tan[e+fx])^n} \int (g \tan[e+fx])^{p-n} (a+b \tan[e+fx])^m (d+c \tan[e+fx])^n dx$$

Program code:

```
Int [(g_*tan[e_+f_*x_])^p_*(a_+b_*tan[e_+f_*x_])^m_*(c_+d_*cot[e_+f_*x_])^n_,x_Symbol] :=
  (g*Tan[e+f*x])^n*(c+d*Cot[e+f*x])^n/(d+c*Tan[e+f*x])^n*Int [(g*Tan[e+f*x])^(p-n)*(a+b*Tan[e+f*x])^m*(d+c*Tan[e+f*x])^n,x] /;
FreeQ[{a,b,c,d,e,f,g,m,n,p},x] && Not[IntegerQ[n]] && Not[IntegerQ[m]]
```