

## Rules for integrands of the form $\text{Trig}[d + e x]^m (a + b \text{Tan}[d + e x]^n + c \text{Tan}[d + e x]^{2n})^p$

$$1. \int (a + b \text{Tan}[d + e x]^n + c \text{Tan}[d + e x]^{2n})^p dx$$

$$1. \int (a + b \text{Tan}[d + e x]^n + c \text{Tan}[d + e x]^{2n})^p dx \text{ when } b^2 - 4ac = 0$$

$$1: \int (a + b \text{Tan}[d + e x]^n + c \text{Tan}[d + e x]^{2n})^p dx \text{ when } b^2 - 4ac = 0 \wedge p \in \mathbb{Z}$$

Derivation: Algebraic simplification

$$\text{Basis: If } b^2 - 4ac = 0, \text{ then } a + bz + cz^2 = \frac{(b+2cz)^2}{4c}$$

Rule: If  $b^2 - 4ac = 0 \wedge p \in \mathbb{Z}$ , then

$$\int (a + b \text{Tan}[d + e x]^n + c \text{Tan}[d + e x]^{2n})^p dx \rightarrow \frac{1}{4^p c^p} \int (b + 2c \text{Tan}[d + e x]^n)^{2p} dx$$

Program code:

```
Int[(a+b_.tan[d_.+e_.x_]^n_.+c_.tan[d_.+e_.x_]^2n_.)^p_.,x_Symbol] :=
  1/(4^p*c^p)*Int[(b+2*c*Tan[d+e*x]^n)^(2*p),x] /;
FreeQ[{a,b,c,d,e,n},x] && EqQ[n2,2*n] && EqQ[b^2-4*a*c,0] && IntegerQ[p]
```

```
Int[(a+b_.cot[d_.+e_.x_]^n_.+c_.cot[d_.+e_.x_]^2n_.)^p_.,x_Symbol] :=
  1/(4^p*c^p)*Int[(b+2*c*Cot[d+e*x]^n)^(2*p),x] /;
FreeQ[{a,b,c,d,e,n},x] && EqQ[n2,2*n] && EqQ[b^2-4*a*c,0] && IntegerQ[p]
```

$$2: \int (a + b \text{Tan}[d + e x]^n + c \text{Tan}[d + e x]^{2n})^p dx \text{ when } b^2 - 4ac = 0 \wedge p \notin \mathbb{Z}$$

Derivation: Piecewise constant extraction

$$\text{Basis: If } b^2 - 4ac = 0, \text{ then } \partial_x \frac{(a+bF[x]+cF[x]^2)^p}{(b+2cF[x])^{2p}} = 0$$

Rule: If  $b^2 - 4ac = 0 \wedge p \notin \mathbb{Z}$ , then

$$\int (a+b \tan [d+e x]^n+c \tan [d+e x]^{2 n})^p dx \rightarrow \frac{(a+b \tan [d+e x]^n+c \tan [d+e x]^{2 n})^p}{(b+2 c \tan [d+e x]^n)^{2 p}} \int (b+2 c \tan [d+e x]^n)^{2 p} dx$$

### Program code:

```
Int [(a_+b_.*tan[d_+e_.*x_]^n_+c_.*tan[d_+e_.*x_]^n2_)^p_,x_Symbol] :=
(a+b*Tan[d+e*x]^n+c*Tan[d+e*x]^(2*n))^p/(b+2*c*Tan[d+e*x]^n)^(2*p)*Int[(b+2*c*Tan[d+e*x]^n)^(2*p),x] /;
FreeQ[{a,b,c,d,e,n},x] && EqQ[n2,2*n] && EqQ[b^2-4*a*c,0] && Not[IntegerQ[p]]
```

```
Int [(a_+b_.*cot[d_+e_.*x_]^n_+c_.*cot[d_+e_.*x_]^n2_)^p_,x_Symbol] :=
(a+b*Cot[d+e*x]^n+c*Cot[d+e*x]^(2*n))^p/(b+2*c*Cot[d+e*x]^n)^(2*p)*Int[(b+2*c*Cot[d+e*x]^n)^(2*p),x] /;
FreeQ[{a,b,c,d,e,n},x] && EqQ[n2,2*n] && EqQ[b^2-4*a*c,0] && Not[IntegerQ[p]]
```

2.  $\int (a+b \tan [d+e x]^n+c \tan [d+e x]^{2 n})^p dx$  when  $b^2-4 a c \neq 0$

1:  $\int \frac{1}{a+b \tan [d+e x]^n+c \tan [d+e x]^{2 n}} dx$  when  $b^2-4 a c \neq 0$

### Derivation: Algebraic expansion

Basis: If  $q = \sqrt{b^2-4 a c}$ , then  $\frac{1}{a+b z+c z^2} = \frac{2 c}{q(b-q+2 c z)} - \frac{2 c}{q(b+q+2 c z)}$

Rule: If  $b^2-4 a c \neq 0$ , let  $q = \sqrt{b^2-4 a c}$ , then

$$\int \frac{1}{a+b \tan [d+e x]^n+c \tan [d+e x]^{2 n}} dx \rightarrow \frac{2 c}{q} \int \frac{1}{b-q+2 c \tan [d+e x]^n} dx - \frac{2 c}{q} \int \frac{1}{b+q+2 c \tan [d+e x]^n} dx$$

### Program code:

```
Int [1/(a_+b_.*tan[d_+e_.*x_]^n_+c_.*tan[d_+e_.*x_]^n2_),x_Symbol] :=
Module[{q=Rt[b^2-4*a*c,2]},
2*c/q*Int[1/(b-q+2*c*Tan[d+e*x]^n),x] -
2*c/q*Int[1/(b+q+2*c*Tan[d+e*x]^n),x] /;
FreeQ[{a,b,c,d,e,n},x] && EqQ[n2,2*n] && NeQ[b^2-4*a*c,0]
```

```
Int[1/(a_.+b_.*cot[d_.+e_.*x_]^n_.+c_.*cot[d_.+e_.*x_]^n2_.),x_Symbol] :=
Module[{q=Rt[b^2-4*a*c,2]},
2*c/q*Int[1/(b-q+2*c*Cot[d+e*x]^n),x] -
2*c/q*Int[1/(b+q+2*c*Cot[d+e*x]^n),x] /;
FreeQ[{a,b,c,d,e,n},x] && EqQ[n2,2*n] && NeQ[b^2-4*a*c,0]
```

$$2. \int \sin[d+e x]^m (a+b (f \tan[d+e x])^n+c (f \tan[d+e x])^{2n})^p dx$$

$$1: \int \sin[d+e x]^m (a+b (f \tan[d+e x])^n+c (f \tan[d+e x])^{2n})^p dx \text{ when } \frac{m}{2} \in \mathbb{Z}$$

Derivation: Integration by substitution

$$\text{Basis: } \sin[z]^2 = \frac{\tan[z]^2}{1+\tan[z]^2}$$

Basis: If  $\frac{m}{2} \in \mathbb{Z}$ , then

$$\sin[d+e x]^m F[f \tan[d+e x]] = \frac{f}{e} \text{Subst}\left[\frac{x^m F[x]}{(f^2+x^2)^{\frac{m}{2}+1}}, x, f \tan[d+e x]\right] \partial_x (f \tan[d+e x])$$

Rule: If  $\frac{m}{2} \in \mathbb{Z}$ , then

$$\int \sin[d+e x]^m (a+b (f \tan[d+e x])^n+c (f \tan[d+e x])^{2n})^p dx \rightarrow \frac{f}{e} \text{Subst}\left[\int \frac{x^m (a+b x^n+c x^{2n})^p}{(f^2+x^2)^{\frac{m}{2}+1}} dx, x, f \tan[d+e x]\right]$$

Program code:

```
Int[sin[d_.+e_.*x_]^m*(a_.+b_.*(f_.*tan[d_.+e_.*x_]^n_.+c_.*(f_.*tan[d_.+e_.*x_]^n2_.)^p_,x_Symbol] :=
f/e*Subst[Int[x^m*(a+b*x^n+c*x^(2*n))^p/(f^2+x^2)^(m/2+1),x],x,f*Tan[d+e*x] /;
FreeQ[{a,b,c,d,e,f,n,p},x] && EqQ[n2,2*n] && IntegerQ[m/2]
```

```
Int[cos[d_.+e_.*x_]^m*(a_.+b_.*(f_.*cot[d_.+e_.*x_]^n_.+c_.*(f_.*cot[d_.+e_.*x_]^n2_.)^p_,x_Symbol] :=
-f/e*Subst[Int[x^m*(a+b*x^n+c*x^(2*n))^p/(f^2+x^2)^(m/2+1),x],x,f*Cot[d+e*x] /;
FreeQ[{a,b,c,d,e,f,n,p},x] && EqQ[n2,2*n] && IntegerQ[m/2]
```

2:  $\int \text{Sin}[d+e x]^m (a+b \text{Tan}[d+e x]^n+c \text{Tan}[d+e x]^{2 n})^p dx$  when  $\frac{m-1}{2} \in \mathbb{Z} \wedge \frac{n}{2} \in \mathbb{Z} \wedge p \in \mathbb{Z}$

Derivation: Integration by substitution

Basis:  $\text{Tan}[z]^2 \equiv \frac{1-\text{Cos}[z]^2}{\text{Cos}[z]^2}$

Basis: If  $\frac{m-1}{2} \in \mathbb{Z} \wedge \frac{n}{2} \in \mathbb{Z}$ , then

$\text{Sin}[d+e x]^m F[\text{Tan}[d+e x]^n] \equiv -\frac{1}{d} \text{Subst}\left[\left(1-x^2\right)^{\frac{m-1}{2}} F\left[\frac{\left(1-x^2\right)^{\frac{n}{2}}}{x^n}\right], x, \text{Cos}[d+e x]\right] \partial_x \text{Cos}[d+e x]$

Rule: If  $\frac{m-1}{2} \in \mathbb{Z} \wedge \frac{n}{2} \in \mathbb{Z} \wedge p \in \mathbb{Z}$ , then

$$\int \text{Sin}[d+e x]^m (a+b \text{Tan}[d+e x]^n+c \text{Tan}[d+e x]^{2 n})^p dx \rightarrow -\frac{1}{d} \text{Subst}\left[\int \frac{\left(1-x^2\right)^{\frac{m-1}{2}}\left(a x^{2 n}+b x^n\left(1-x^2\right)^{n / 2}+c\left(1-x^2\right)^n\right)^p}{x^{2 n p}} dx, x, \text{Cos}[d+e x]\right]$$

Program code:

```
Int[sin[d_+e_*x_]^m_.*(a_+b_*tan[d_+e_*x_]^n_+c_*tan[d_+e_*x_]^(2n_)^p_,x_Symbol] :=
Module[{g=FreeFactors[Cos[d+e*x],x]},
-g/e*Subst[Int[(1-g^2*x^2)^((m-1)/2)*ExpandToSum[a*(g*x)^(2*n)+b*(g*x)^n*(1-g^2*x^2)^(n/2)+c*(1-g^2*x^2)^n,x]^p/(g*x)^(2*n*p),x],x,Cos[d+e*x],
FreeQ[{a,b,c,d,e},x] && EqQ[n2,2*n] && IntegerQ[(m-1)/2] && IntegerQ[n/2] && IntegerQ[p]
```

```
Int[cos[d_+e_*x_]^m_.*(a_+b_*cot[d_+e_*x_]^n_+c_*tan[d_+e_*x_]^(2n_)^p_,x_Symbol] :=
Module[{g=FreeFactors[Sin[d+e*x],x]},
g/e*Subst[Int[(1-g^2*x^2)^((m-1)/2)*ExpandToSum[a*(g*x)^(2*n)+b*(g*x)^n*(1-g^2*x^2)^(n/2)+c*(1-g^2*x^2)^n,x]^p/(g*x)^(2*n*p),x],x,Sin[d+e*x],
FreeQ[{a,b,c,d,e},x] && EqQ[n2,2*n] && IntegerQ[(m-1)/2] && IntegerQ[n/2] && IntegerQ[p]
```

$$3. \int \cos [d+e x]^m (a+b \tan [d+e x]^n+c \tan [d+e x]^{2 n})^p dx$$

$$1: \int \cos [d+e x]^m (a+b \tan [d+e x]^n+c \tan [d+e x]^{2 n})^p dx \text{ when } \frac{m}{2} \in \mathbb{Z}$$

Derivation: Integration by substitution

$$\text{Basis: } \cos [z]^2 = \frac{1}{1+\tan [z]^2}$$

Basis: If  $\frac{m}{2} \in \mathbb{Z}$ , then

$$\cos [d+e x]^m F [f \tan [d+e x]] = \frac{f^{m+1}}{e} \text{Subst} \left[ \frac{F[x]}{(f^2+x^2)^{\frac{m}{2}+1}}, x, f \tan [d+e x] \right] \partial_x (f \tan [d+e x])$$

Rule: If  $\frac{m}{2} \in \mathbb{Z}$ , then

$$\int \cos [d+e x]^m (a+b \tan [d+e x]^n+c \tan [d+e x]^{2 n})^p dx \rightarrow \frac{f^{m+1}}{e} \text{Subst} \left[ \int \frac{(a+b x^n+c x^{2 n})^p}{(f^2+x^2)^{\frac{m}{2}+1}} dx, x, f \tan [d+e x] \right]$$

Program code:

```
Int [cos [d_+e_.*x_] ^m_*(a_+b_.*(f_.*tan [d_+e_.*x_] ) ^n_+c_.*(f_.*tan [d_+e_.*x_] ) ^2n_) ^p_.,x_Symbol] :=
  f^(m+1)/e*Subst [Int [(a+b*x^n+c*x^(2*n))^p/(f^2+x^2)^(m/2+1),x],x,f*Tan [d+e*x]] /;
FreeQ[{a,b,c,d,e,f,n,p},x] && EqQ[n2,2*n] && IntegerQ[m/2]
```

```
Int [sin [d_+e_.*x_] ^m_*(a_+b_.*(f_.*cot [d_+e_.*x_] ) ^n_+c_.*(f_.*cot [d_+e_.*x_] ) ^2n_) ^p_.,x_Symbol] :=
  -f^(m+1)/e*Subst [Int [(a+b*x^n+c*x^(2*n))^p/(f^2+x^2)^(m/2+1),x],x,f*Cot [d+e*x]] /;
FreeQ[{a,b,c,d,e,f,n,p},x] && EqQ[n2,2*n] && IntegerQ[m/2]
```

$$2: \int \cos [d+e x]^m (a+b \tan [d+e x]^n+c \tan [d+e x]^{2n})^p dx \text{ when } \frac{m-1}{2} \in \mathbb{Z} \wedge \frac{n}{2} \in \mathbb{Z} \wedge p \in \mathbb{Z}$$

Derivation: Integration by substitution

$$\text{Basis: } \tan [z]^2 \equiv \frac{\sin [z]^2}{1-\sin [z]^2}$$

Basis: If  $\frac{m-1}{2} \in \mathbb{Z} \wedge \frac{n}{2} \in \mathbb{Z}$ , then

$$\cos [d+e x]^m F[\tan [d+e x]^n] \equiv \frac{1}{e} \text{Subst} \left[ (1-x^2)^{\frac{m-1}{2}} F \left[ \frac{x^n}{(1-x^2)^{\frac{n}{2}}} \right], x, \sin [d+e x] \right] \partial_x \sin [d+e x]$$

Rule: If  $\frac{m-1}{2} \in \mathbb{Z} \wedge \frac{n}{2} \in \mathbb{Z} \wedge p \in \mathbb{Z}$ , then

$$\int \cos [d+e x]^m (a+b \tan [d+e x]^n+c \tan [d+e x]^{2n})^p dx \rightarrow \frac{1}{e} \text{Subst} \left[ \int (1-x^2)^{(m-2n p-1)/2} (c x^{2n}+b x^n (1-x^2)^{n/2}+a (1-x^2)^n)^p dx, x, \sin [d+e x] \right]$$

Program code:

```
Int[cos[d_+e_.*x_]^m_*(a_+b_.*tan[d_+e_.*x_]^n_+c_.*tan[d_+e_.*x_]^2n_)^p_,x_Symbol] :=
Module[{g=FreeFactors[Sin[d+e*x],x]},
g/e*Subst[Int[(1-g^2*x^2)^(m-2*n*p-1)/2]*ExpandToSum[c*x^(2*n)+b*x^n*(1-x^2)^(n/2)+a*(1-x^2)^n,x]^p,x],x,Sin[d+e*x]/g] /;
FreeQ[{a,b,c,d,e},x] && EqQ[n2,2*n] && IntegerQ[(m-1)/2] && IntegerQ[n/2] && IntegerQ[p]
```

```
Int[sin[d_+e_.*x_]^m_*(a_+b_.*cot[d_+e_.*x_]^n_+c_.*cot[d_+e_.*x_]^2n_)^p_,x_Symbol] :=
Module[{g=FreeFactors[Cos[d+e*x],x]},
-g/e*Subst[Int[(1-g^2*x^2)^(m-2*n*p-1)/2]*ExpandToSum[c*x^(2*n)+b*x^n*(1-x^2)^(n/2)+a*(1-x^2)^n,x]^p,x],x,Cos[d+e*x]/g] /;
FreeQ[{a,b,c,d,e},x] && EqQ[n2,2*n] && IntegerQ[(m-1)/2] && IntegerQ[n/2] && IntegerQ[p]
```

$$4. \int \tan [d+e x]^m (a+b \tan [d+e x]^n+c \tan [d+e x]^{2 n})^p dx$$

$$1. \int \tan [d+e x]^m (a+b \tan [d+e x]^n+c \tan [d+e x]^{2 n})^p dx \text{ when } b^2-4 a c == 0$$

$$1: \int \tan [d+e x]^m (a+b \tan [d+e x]^n+c \tan [d+e x]^{2 n})^p dx \text{ when } b^2-4 a c == 0 \wedge p \in \mathbb{Z}$$

Derivation: Algebraic simplification

$$\text{Basis: If } b^2-4 a c == 0, \text{ then } a+b z+c z^2 == \frac{(b+2 c z)^2}{4 c}$$

Rule: If  $b^2-4 a c == 0 \wedge p \in \mathbb{Z}$ , then

$$\int \tan [d+e x]^m (a+b \tan [d+e x]^n+c \tan [d+e x]^{2 n})^p dx \rightarrow \frac{1}{4^p c^p} \int \tan [d+e x]^m (b+2 c \tan [d+e x]^n)^{2 p} dx$$

Program code:

```
Int[tan[d_+e_.*x_]^m_.*(a_+b_.*tan[d_+e_.*x_]^n_+c_.*tan[d_+e_.*x_]^(2*n_)^p_,x_Symbol] :=
  1/(4^p*c^p)*Int[Tan[d+e*x]^m*(b+2*c*Tan[d+e*x]^n)^(2*p),x] /;
FreeQ[{a,b,c,d,e,m,n},x] && EqQ[n2,2*n] && EqQ[b^2-4*a*c,0] && IntegerQ[p]
```

```
Int[cot[d_+e_.*x_]^m_.*(a_+b_.*cot[d_+e_.*x_]^n_+c_.*cot[d_+e_.*x_]^(2*n_)^p_,x_Symbol] :=
  1/(4^p*c^p)*Int[Cot[d+e*x]^m*(b+2*c*Cot[d+e*x]^n)^(2*p),x] /;
FreeQ[{a,b,c,d,e,m,n},x] && EqQ[n2,2*n] && EqQ[b^2-4*a*c,0] && IntegerQ[p]
```

$$2: \int \tan [d+e x]^m (a+b \tan [d+e x]^n+c \tan [d+e x]^{2 n})^p dx \text{ when } b^2-4 a c == 0 \wedge p \notin \mathbb{Z}$$

Derivation: Piecewise constant extraction

$$\text{Basis: If } b^2-4 a c == 0, \text{ then } \partial_x \frac{(a+b F[x]+c F[x]^2)^p}{(b+2 c F[x])^{2 p}} == 0$$

Rule: If  $b^2-4 a c == 0 \wedge p \notin \mathbb{Z}$ , then

$$\int \text{Tan}[d+e x]^m (a+b \text{Tan}[d+e x]^n+c \text{Tan}[d+e x]^{2n})^p dx \rightarrow \frac{(a+b \text{Tan}[d+e x]^n+c \text{Tan}[d+e x]^{2n})^p}{(b+2c \text{Tan}[d+e x]^n)^{2p}} \int \text{Tan}[d+e x]^m (b+2c \text{Tan}[d+e x]^n)^{2p} dx$$

### Program code:

```
Int[tan[d_+e_.*x_]^m_.*(a_+b_.*tan[d_+e_.*x_]^n_+c_.*tan[d_+e_.*x_]^2n_)^p_,x_Symbol] :=
(a+b*Tan[d+e*x]^n+c*Tan[d+e*x]^(2*n))^p/(b+2*c*Tan[d+e*x]^n)^(2*p)*Int[Tan[d+e*x]^m*(b+2*c*Tan[d+e*x]^n)^(2*p),x] /;
FreeQ[{a,b,c,d,e,m,n,p},x] && EqQ[n2,2*n] && EqQ[b^2-4*a*c,0] && Not[IntegerQ[p]]
```

```
Int[cot[d_+e_.*x_]^m_.*(a_+b_.*cot[d_+e_.*x_]^n_+c_.*cot[d_+e_.*x_]^2n_)^p_,x_Symbol] :=
(a+b*Cot[d+e*x]^n+c*Cot[d+e*x]^(2*n))^p/(b+2*c*Cot[d+e*x]^n)^(2*p)*Int[Cot[d+e*x]^m*(b+2*c*Cot[d+e*x]^n)^(2*p),x] /;
FreeQ[{a,b,c,d,e,m,n,p},x] && EqQ[n2,2*n] && EqQ[b^2-4*a*c,0] && Not[IntegerQ[p]]
```

2:  $\int \text{Tan}[d+e x]^m (a+b (f \text{Tan}[d+e x])^n+c (f \text{Tan}[d+e x])^{2n})^p dx$  when  $b^2 - 4ac \neq 0$

### Derivation: Integration by substitution

Basis:  $\text{Tan}[d+e x]^m F[f \text{Tan}[d+e x]] = \frac{f}{e} \text{Subst}\left[\left(\frac{x}{f}\right)^m \frac{F[x]}{f^2+x^2}, x, f \text{Tan}[d+e x]\right] \partial_x (f \text{Tan}[d+e x])$

Rule: If  $b^2 - 4ac \neq 0$ , then

$$\int \text{Tan}[d+e x]^m (a+b (f \text{Tan}[d+e x])^n+c (f \text{Tan}[d+e x])^{2n})^p dx \rightarrow \frac{f}{e} \text{Subst}\left[\int \left(\frac{x}{f}\right)^m \frac{(a+b x^n+c x^{2n})^p}{f^2+x^2} dx, x, f \text{Tan}[d+e x]\right]$$

### Program code:

```
Int[tan[d_+e_.*x_]^m_.*(a_+b_.*(f_.*tan[d_+e_.*x_]^n_+c_.*(f_.*tan[d_+e_.*x_]^2n_)^p_,x_Symbol] :=
f/e*Subst[Int[(x/f)^m*(a+b*x^n+c*x^(2*n))^p/(f^2+x^2),x],x,f*Tan[d+e*x]] /;
FreeQ[{a,b,c,d,e,f,m,n,p},x] && EqQ[n2,2*n] && NeQ[b^2-4*a*c,0]
```

```
Int[cot[d_+e_.*x_]^m_.*(a_+b_.*(f_.*cot[d_+e_.*x_]^n_+c_.*(f_.*cot[d_+e_.*x_]^2n_)^p_,x_Symbol] :=
-f/e*Subst[Int[(x/f)^m*(a+b*x^n+c*x^(2*n))^p/(f^2+x^2),x],x,f*Cot[d+e*x]] /;
FreeQ[{a,b,c,d,e,f,m,n,p},x] && EqQ[n2,2*n] && NeQ[b^2-4*a*c,0]
```



$$5. \int \text{Cot}[d+e x]^m (a+b \text{Tan}[d+e x]^n+c \text{Tan}[d+e x]^{2 n})^p dx$$

$$1. \int \text{Cot}[d+e x]^m (a+b \text{Tan}[d+e x]^n+c \text{Tan}[d+e x]^{2 n})^p dx \text{ when } b^2-4 a c == 0$$

$$1: \int \text{Cot}[d+e x]^m (a+b \text{Tan}[d+e x]^n+c \text{Tan}[d+e x]^{2 n})^p dx \text{ when } b^2-4 a c == 0 \wedge p \in \mathbb{Z}$$

Derivation: Algebraic simplification

$$\text{Basis: If } b^2-4 a c == 0, \text{ then } a+b z+c z^2 == \frac{(b+2 c z)^2}{4 c}$$

Rule: If  $b^2-4 a c == 0 \wedge p \in \mathbb{Z}$ , then

$$\int \text{Cot}[d+e x]^m (a+b \text{Tan}[d+e x]^n+c \text{Tan}[d+e x]^{2 n})^p dx \rightarrow \frac{1}{4^p c^p} \int \text{Cot}[d+e x]^m (b+2 c \text{Tan}[d+e x]^n)^{2 p} dx$$

Program code:

```
Int[cot[d_+e_.*x_]^m_.*(a_+b_.*tan[d_+e_.*x_]^n_+c_.*tan[d_+e_.*x_]^(2*n_)^p_.,x_Symbol] :=
  1/(4^p*c^p)*Int[Cot[d+e*x]^m*(b+2*c*Tan[d+e*x]^n)^(2*p),x] /;
FreeQ[{a,b,c,d,e,m,n},x] && EqQ[n2,2*n] && EqQ[b^2-4*a*c,0] && IntegerQ[p]
```

```
Int[tan[d_+e_.*x_]^m_.*(a_+b_.*cot[d_+e_.*x_]^n_+c_.*cot[d_+e_.*x_]^(2*n_)^p_.,x_Symbol] :=
  1/(4^p*c^p)*Int[Tan[d+e*x]^m*(b+2*c*Cot[d+e*x]^n)^(2*p),x] /;
FreeQ[{a,b,c,d,e,m,n},x] && EqQ[n2,2*n] && EqQ[b^2-4*a*c,0] && IntegerQ[p]
```

$$2: \int \text{Cot}[d+e x]^m (a+b \text{Tan}[d+e x]^n+c \text{Tan}[d+e x]^{2 n})^p dx \text{ when } b^2-4 a c == 0 \wedge p \notin \mathbb{Z}$$

Derivation: Piecewise constant extraction

$$\text{Basis: If } b^2-4 a c == 0, \text{ then } \partial_x \frac{(a+b F[x]+c F[x]^2)^p}{(b+2 c F[x])^{2 p}} == 0$$

Rule: If  $b^2-4 a c == 0 \wedge p \notin \mathbb{Z}$ , then

$$\int \text{Cot}[d+e x]^m (a+b \tan[d+e x]^n+c \tan[d+e x]^{2n})^p dx \rightarrow \frac{(a+b \tan[d+e x]^n+c \tan[d+e x]^{2n})^p}{(b+2c \tan[d+e x]^n)^{2p}} \int \text{Cot}[d+e x]^m (b+2c \tan[d+e x]^n)^{2p} dx$$

Program code:

```
Int[cot[d_+e_.*x_]^m_.*(a_+b_.*tan[d_+e_.*x_]^n_+c_.*tan[d_+e_.*x_]^(2*n_)^p_,x_Symbol] :=
(a+b*Tan[d+e*x]^n+c*Tan[d+e*x]^(2*n))^p/(b+2*c*Tan[d+e*x]^n)^(2*p)*Int[Cot[d+e*x]^m*(b+2*c*Tan[d+e*x]^n)^(2*p),x] /;
FreeQ[{a,b,c,d,e,m,n,p},x] && EqQ[n2,2*n] && EqQ[b^2-4*a*c,0] && Not[IntegerQ[p]]
```

```
Int[tan[d_+e_.*x_]^m_.*(a_+b_.*cot[d_+e_.*x_]^n_+c_.*cot[d_+e_.*x_]^(2*n_)^p_,x_Symbol] :=
(a+b*Cot[d+e*x]^n+c*Cot[d+e*x]^(2*n))^p/(b+2*c*Cot[d+e*x]^n)^(2*p)*Int[Tan[d+e*x]^m*(b+2*c*Cot[d+e*x]^n)^(2*p),x] /;
FreeQ[{a,b,c,d,e,m,n,p},x] && EqQ[n2,2*n] && EqQ[b^2-4*a*c,0] && Not[IntegerQ[p]]
```

2:  $\int \text{Cot}[d+e x]^m (a+b \tan[d+e x]^n+c \tan[d+e x]^{2n})^p dx$  when  $b^2-4ac \neq 0 \wedge \frac{n}{2} \in \mathbb{Z}$

Derivation: Integration by substitution

Basis:  $\text{Tan}[z]^2 \equiv \frac{1}{\text{Cot}[z]^2}$

Basis:  $\text{Cot}[d+e x]^m F[\text{Tan}[d+e x]^2] \equiv -\frac{1}{e} \text{Subst}\left[\frac{x^m F\left[\frac{1}{x^2}\right]}{1+x^2}, x, \text{Cot}[d+e x]\right] \partial_x \text{Cot}[d+e x]$

Rule: If  $b^2-4ac \neq 0 \wedge \frac{n}{2} \in \mathbb{Z}$ , then

$$\int \text{Cot}[d+e x]^m (a+b \tan[d+e x]^n+c \tan[d+e x]^{2n})^p dx \rightarrow -\frac{1}{e} \text{Subst}\left[\int \frac{x^{m-2np} (c+b x^n+a x^{2n})^p}{1+x^2} dx, x, \text{Cot}[d+e x]\right]$$

Program code:

```
Int[cot[d_+e_.*x_]^m_.*(a_+b_.*tan[d_+e_.*x_]^n_+c_.*tan[d_+e_.*x_]^(2*n_)^p_,x_Symbol] :=
Module[{g=FreeFactors[Cot[d+e*x],x]},
g/e*Subst[Int[(g*x)^(m-2*n*p)*(c+b*(g*x)^n+a*(g*x)^(2*n))^p/(1+g^2*x^2),x],x,Cot[d+e*x]/g] /;
FreeQ[{a,b,c,d,e,m,p},x] && EqQ[n2,2*n] && NeQ[b^2-4*a*c,0] && IntegerQ[n/2]
```

```
Int [tan [d_ .+e_ . *x_] ^m_ . * (a_ .+b_ . *cot [d_ .+e_ . *x_] ^n_ .+c_ . *cot [d_ .+e_ . *x_] ^n2_) ^p_ . ,x_Symbol] :=
Module [{g=FreeFactors [Tan [d+e*x] ,x] ,
-g/e*Subst [Int [(g*x) ^ (m-2*n*p) * (c+b*(g*x) ^n+a*(g*x) ^ (2*n)) ^p / (1+g^2*x^2) ,x] ,x,Tan [d+e*x] /g] ] /;
FreeQ[{a,b,c,d,e,m,p} ,x] && EqQ[n2,2*n] && NeQ[b^2-4*a*c,0] && IntegerQ[n/2]
```

$$6. \int (A + B \tan [d + e x]) (a + b \tan [d + e x] + c \tan [d + e x]^2)^n dx$$

$$1. \int (A + B \tan [d + e x]) (a + b \tan [d + e x] + c \tan [d + e x]^2)^n dx \text{ when } b^2 - 4 a c = 0$$

$$1: \int (A + B \tan [d + e x]) (a + b \tan [d + e x] + c \tan [d + e x]^2)^n dx \text{ when } b^2 - 4 a c = 0 \wedge n \in \mathbb{Z}$$

Derivation: Algebraic simplification

$$\text{Basis: If } b^2 - 4 a c = 0, \text{ then } a + b z + c z^2 = \frac{(b+2 c z)^2}{4 c}$$

Rule: If  $b^2 - 4 a c = 0 \wedge n \in \mathbb{Z}$ , then

$$\int (A + B \tan [d + e x]) (a + b \tan [d + e x] + c \tan [d + e x]^2)^n dx \rightarrow \frac{1}{4^n c^n} \int (A + B \tan [d + e x]) (b + 2 c \tan [d + e x])^{2 n} dx$$

Program code:

```
Int [(A_+B_.*tan [d_ .+e_ . *x_] ) * (a_+b_.*tan [d_ .+e_ . *x_] +c_.*tan [d_ .+e_ . *x_] ^2) ^n_ ,x_Symbol] :=
1 / (4^n*c^n) *Int [(A+B*Tan [d+e*x] ) * (b+2*c*Tan [d+e*x] ) ^ (2*n) ,x] /;
FreeQ[{a,b,c,d,e,A,B} ,x] && EqQ[b^2-4*a*c,0] && IntegerQ[n]
```

```
Int [(A_+B_.*cot [d_ .+e_ . *x_] ) * (a_+b_.*cot [d_ .+e_ . *x_] +c_.*cot [d_ .+e_ . *x_] ^2) ^n_ ,x_Symbol] :=
1 / (4^n*c^n) *Int [(A+B*Cot [d+e*x] ) * (b+2*c*Cot [d+e*x] ) ^ (2*n) ,x] /;
FreeQ[{a,b,c,d,e,A,B} ,x] && EqQ[b^2-4*a*c,0] && IntegerQ[n]
```

$$2: \int (A + B \tan [d + e x]) (a + b \tan [d + e x] + c \tan [d + e x]^2)^n dx \text{ when } b^2 - 4 a c = 0 \wedge n \notin \mathbb{Z}$$

Derivation: Piecewise constant extraction

Basis: If  $b^2 - 4 a c == 0$ , then  $\partial_x \frac{(a+b F[x]+c F[x]^2)^n}{(b+2 c F[x])^{2 n}} == 0$

Rule: If  $b^2 - 4 a c == 0 \wedge n \notin \mathbb{Z}$ , then

$$\int (A+B \tan [d+e x]) (a+b \tan [d+e x]+c \tan [d+e x]^2)^n dx \rightarrow \frac{(a+b \tan [d+e x]+c \tan [d+e x]^2)^n}{(b+2 c \tan [d+e x])^{2 n}} \int (A+B \tan [d+e x]) (b+2 c \tan [d+e x])^{2 n} dx$$

Program code:

```
Int[(A+B_.*tan[d_.+e_.*x_])*(a+b_.*tan[d_.+e_.*x_]+c_.*tan[d_.+e_.*x_]^2)^n_,x_Symbol] :=
(a+b*Tan[d+e*x]+C*Tan[d+e*x]^2)^n/(b+2*c*Tan[d+e*x])^(2*n)*Int[(A+B*Tan[d+e*x])*(b+2*c*Tan[d+e*x])^(2*n),x] /;
FreeQ[{a,b,c,d,e,A,B},x] && EqQ[b^2-4*a*c,0] && Not[IntegerQ[n]]
```

```
Int[(A+B_.*cot[d_.+e_.*x_])*(a+b_.*cot[d_.+e_.*x_]+c_.*cot[d_.+e_.*x_]^2)^n_,x_Symbol] :=
(a+b*Cot[d+e*x]+C*Cot[d+e*x]^2)^n/(b+2*c*Cot[d+e*x])^(2*n)*Int[(A+B*Cot[d+e*x])*(b+2*c*Cot[d+e*x])^(2*n),x] /;
FreeQ[{a,b,c,d,e,A,B},x] && EqQ[b^2-4*a*c,0] && Not[IntegerQ[n]]
```

$$2. \int (A + B \tan[d + e x]) (a + b \tan[d + e x] + c \tan[d + e x]^2)^n dx \text{ when } b^2 - 4ac \neq 0$$

$$1: \int \frac{A + B \tan[d + e x]}{a + b \tan[d + e x] + c \tan[d + e x]^2} dx \text{ when } b^2 - 4ac \neq 0$$

Derivation: Algebraic expansion

Basis: If  $q = \sqrt{b^2 - 4ac}$ , then  $\frac{A+Bz}{a+bz+cz^2} = \left(B + \frac{bB-2Ac}{q}\right) \frac{1}{b+q+2cz} + \left(B - \frac{bB-2Ac}{q}\right) \frac{1}{b-q+2cz}$

■ Rule: If  $b^2 - 4ac \neq 0$ , let  $q = \sqrt{b^2 - 4ac}$ , then

$$\int \frac{A + B \tan[d + e x]}{a + b \tan[d + e x] + c \tan[d + e x]^2} dx \rightarrow \left(B + \frac{bB - 2Ac}{q}\right) \int \frac{1}{b + q + 2c \tan[d + e x]} dx + \left(B - \frac{bB - 2Ac}{q}\right) \int \frac{1}{b - q + 2c \tan[d + e x]} dx$$

– Program code:

```
Int[(A+B_.*tan[d_.+e_.*x_])/(a_.+b_.*tan[d_.+e_.*x_]+c_.*tan[d_.+e_.*x_]^2),x_Symbol] :=
Module[{q=Rt[b^2-4*a*c,2]},
(B+(b*B-2*A*c)/q)*Int[1/Simp[b+q+2*c*Tan[d+e*x],x],x] +
(B-(b*B-2*A*c)/q)*Int[1/Simp[b-q+2*c*Tan[d+e*x],x],x] /;
FreeQ[{a,b,c,d,e,A,B},x] && NeQ[b^2-4*a*c,0]
```

```
Int[(A+B_.*cot[d_.+e_.*x_])/(a_.+b_.*cot[d_.+e_.*x_]+c_.*cot[d_.+e_.*x_]^2),x_Symbol] :=
Module[{q=Rt[b^2-4*a*c,2]},
(B+(b*B-2*A*c)/q)*Int[1/Simp[b+q+2*c*Cot[d+e*x],x],x] +
(B-(b*B-2*A*c)/q)*Int[1/Simp[b-q+2*c*Cot[d+e*x],x],x] /;
FreeQ[{a,b,c,d,e,A,B},x] && NeQ[b^2-4*a*c,0]
```

$$2: \int (A+B \tan [d+e x]) (a+b \tan [d+e x]+c \tan [d+e x]^2)^n dx \text{ when } b^2-4 a c \neq 0 \wedge n \in \mathbb{Z}$$

### Derivation: Algebraic expansion

Rule: If  $b^2-4 a c \neq 0 \wedge n \in \mathbb{Z}$

$$\int (A+B \tan [d+e x]) (a+b \tan [d+e x]+c \tan [d+e x]^2)^n dx \rightarrow \int \text{ExpandTrig}[(A+B \tan [d+e x]) (a+b \tan [d+e x]+c \tan [d+e x]^2)^n, x] dx$$

### Program code:

```
Int[(A+B_.*tan[d_.+e_.*x_])*(a_.+b_.*tan[d_.+e_.*x_]+c_.*tan[d_.+e_.*x_]^2)^n_,x_Symbol] :=
  Int[ExpandTrig[(A+B*tan[d+e*x])*(a+b*tan[d+e*x]+c*tan[d+e*x]^2)^n,x],x] /;
FreeQ[{a,b,c,d,e,A,B},x] && NeQ[b^2-4*a*c,0] && IntegerQ[n]
```

```
Int[(A+B_.*cot[d_.+e_.*x_])*(a_.+b_.*cot[d_.+e_.*x_]+c_.*cot[d_.+e_.*x_]^2)^n_,x_Symbol] :=
  Int[ExpandTrig[(A+B*cot[d+e*x])*(a+b*cot[d+e*x]+c*cot[d+e*x]^2)^n,x],x] /;
FreeQ[{a,b,c,d,e,A,B},x] && NeQ[b^2-4*a*c,0] && IntegerQ[n]
```