

Rules for integrands of the form $(d x)^m (a + b \operatorname{ArcTan}[c x^n])^p$

1. $\int x^m (a + b \operatorname{ArcTan}[c x^n])^p dx$ when $p \in \mathbb{Z}^+$

1. $\int \frac{(a + b \operatorname{ArcTan}[c x^n])^p}{x} dx$ when $p \in \mathbb{Z}^+$

1. $\int \frac{(a + b \operatorname{ArcTan}[c x])^p}{x} dx$ when $p \in \mathbb{Z}^+$

1: $\int \frac{a + b \operatorname{ArcTan}[c x]}{x} dx$

Derivation: Algebraic expansion

Basis: $\operatorname{ArcTan}[z] = \frac{i}{2} \operatorname{Log}[1 - iz] - \frac{i}{2} \operatorname{Log}[1 + iz]$

Basis: $\operatorname{ArcCot}[z] = \frac{i}{2} \operatorname{Log}[1 - \frac{i}{z}] - \frac{i}{2} \operatorname{Log}[1 + \frac{i}{z}]$

Rule:

$$\int \frac{a + b \operatorname{ArcTan}[c x]}{x} dx \rightarrow a \int \frac{1}{x} dx + \frac{ib}{2} \int \frac{\operatorname{Log}[1 - icx]}{x} dx - \frac{ib}{2} \int \frac{\operatorname{Log}[1 + icx]}{x} dx$$

$$\rightarrow a \operatorname{Log}[x] + \frac{ib}{2} \operatorname{PolyLog}[2, -icx] - \frac{ib}{2} \operatorname{PolyLog}[2, icx]$$

Program code:

```
Int[(a_.+b_.*ArcTan[c_.*x_])/x_,x_Symbol] :=
  a*Log[x] + I*b/2*Int[Log[1-I*c*x]/x,x] - I*b/2*Int[Log[1+I*c*x]/x,x] /;
FreeQ[{a,b,c},x]
```

```
Int[(a_.+b_.*ArcCot[c_.*x_])/x_,x_Symbol] :=
  a*Log[x] + I*b/2*Int[Log[1-I/(c*x)]/x,x] - I*b/2*Int[Log[1+I/(c*x)]/x,x] /;
FreeQ[{a,b,c},x]
```

$$2: \int \frac{(a + b \operatorname{ArcTan}[c x])^p}{x} dx \text{ when } p - 1 \in \mathbb{Z}^+$$

Derivation: Integration by parts

$$\text{Basis: } \frac{1}{x} = 2 \partial_x \operatorname{ArcTanh} \left[1 - \frac{2}{1 + i c x} \right]$$

Rule: If $p - 1 \in \mathbb{Z}^+$, then

$$\int \frac{(a + b \operatorname{ArcTan}[c x])^p}{x} dx \rightarrow 2 (a + b \operatorname{ArcTan}[c x])^p \operatorname{ArcTanh} \left[1 - \frac{2}{1 + i c x} \right] - 2 b c p \int \frac{(a + b \operatorname{ArcTan}[c x])^{p-1} \operatorname{ArcTanh} \left[1 - \frac{2}{1 + i c x} \right]}{1 + c^2 x^2} dx$$

Program code:

```
Int[(a_ + b_.*ArcTan[c_*x_])^p_/x_, x_Symbol] :=
  2*(a + b*ArcTan[c*x])^p*ArcTanh[1 - 2/(1 + I*c*x)] -
  2*b*c*p*Int[(a + b*ArcTan[c*x])^(p-1)*ArcTanh[1 - 2/(1 + I*c*x)]/(1 + c^2*x^2), x] /;
FreeQ[{a, b, c}, x] && IGtQ[p, 1]
```

```
Int[(a_ + b_.*ArcCot[c_*x_])^p_/x_, x_Symbol] :=
  2*(a + b*ArcCot[c*x])^p*ArcCoth[1 - 2/(1 + I*c*x)] +
  2*b*c*p*Int[(a + b*ArcCot[c*x])^(p-1)*ArcCoth[1 - 2/(1 + I*c*x)]/(1 + c^2*x^2), x] /;
FreeQ[{a, b, c}, x] && IGtQ[p, 1]
```

$$2: \int \frac{(a + b \operatorname{ArcTan}[c x^n])^p}{x} dx \text{ when } p \in \mathbb{Z}^+$$

Derivation: Integration by substitution

$$\text{Basis: } \frac{F[x^n]}{x} = \frac{1}{n} \operatorname{Subst}\left[\frac{F[x]}{x}, x, x^n\right] \partial_x x^n$$

Rule: If $p \in \mathbb{Z}^+$, then

$$\int \frac{(a + b \operatorname{ArcTan}[c x^n])^p}{x} dx \rightarrow \frac{1}{n} \operatorname{Subst}\left[\int \frac{(a + b \operatorname{ArcTan}[c x])^p}{x} dx, x, x^n\right]$$

Program code:

```
Int[(a_.+b_.*ArcTan[c_.*x^n_])^p_/x_,x_Symbol] :=
  1/n*Subst[Int[(a+b*ArcTan[c*x])^p/x,x],x,x^n] /;
FreeQ[{a,b,c,n},x] && IGtQ[p,0]
```

```
Int[(a_.+b_.*ArcCot[c_.*x^n_])^p_/x_,x_Symbol] :=
  1/n*Subst[Int[(a+b*ArcCot[c*x])^p/x,x],x,x^n] /;
FreeQ[{a,b,c,n},x] && IGtQ[p,0]
```

2: $\int x^m (a + b \operatorname{ArcTan}[c x^n])^p dx$ when $p \in \mathbb{Z}^+ \wedge (p = 1 \vee n = 1 \wedge m \in \mathbb{Z}) \wedge m \neq -1$

Derivation: Integration by parts

Basis: $\partial_x (a + b \operatorname{ArcTan}[c x^n])^p = b c n p \frac{x^{n-1} (a + b \operatorname{ArcTan}[c x^n])^{p-1}}{1 + c^2 x^{2n}}$

Rule: If $p \in \mathbb{Z}^+ \wedge (p = 1 \vee n = 1 \wedge m \in \mathbb{Z}) \wedge m \neq -1$, then

$$\int x^m (a + b \operatorname{ArcTan}[c x^n])^p dx \rightarrow \frac{x^{m+1} (a + b \operatorname{ArcTan}[c x^n])^p}{m+1} - \frac{b c n p}{m+1} \int \frac{x^{m+n} (a + b \operatorname{ArcTan}[c x^n])^{p-1}}{1 + c^2 x^{2n}} dx$$

Program code:

```
Int[x^m_.*(a_.+b_.*ArcTan[c_*x^n_])^p_.,x_Symbol] :=
  x^(m+1)*(a+b*ArcTan[c*x^n])^p/(m+1) -
  b*c*n*p/(m+1)*Int[x^(m+n)*(a+b*ArcTan[c*x^n])^(p-1)/(1+c^2*x^(2*n)),x] /;
FreeQ[{a,b,c,m,n},x] && IGtQ[p,0] && (EqQ[p,1] || EqQ[n,1] && IntegerQ[m]) && NeQ[m,-1]
```

```
Int[x^m_.*(a_.+b_.*ArcCot[c_*x^n_])^p_.,x_Symbol] :=
  x^(m+1)*(a+b*ArcCot[c*x^n])^p/(m+1) +
  b*c*n*p/(m+1)*Int[x^(m+n)*(a+b*ArcCot[c*x^n])^(p-1)/(1+c^2*x^(2*n)),x] /;
FreeQ[{a,b,c,m,n},x] && IGtQ[p,0] && (EqQ[p,1] || EqQ[n,1] && IntegerQ[m]) && NeQ[m,-1]
```

3: $\int x^m (a + b \operatorname{ArcTan}[c x^n])^p dx$ when $p - 1 \in \mathbb{Z}^+ \wedge \frac{m+1}{n} \in \mathbb{Z}$

Derivation: Integration by substitution

Basis: If $\frac{m+1}{n} \in \mathbb{Z}$, then $x^m F[x^n] = \frac{1}{n} \operatorname{Subst}[x^{\frac{m+1}{n}-1} F[x], x, x^n] dx x^n$

Rule: If $p - 1 \in \mathbb{Z}^+ \wedge \frac{m+1}{n} \in \mathbb{Z}$, then

$$\int x^m (a + b \operatorname{ArcTan}[c x^n])^p dx \rightarrow \frac{1}{n} \operatorname{Subst}\left[\int x^{\frac{m+1}{n}-1} (a + b \operatorname{ArcTan}[c x])^p dx, x, x^n\right]$$

Program code:

```
Int[x_^m_.*(a_.*b_.*ArcTan[c_.*x_^n_])^p_.,x_Symbol] :=
  1/n*Subst[Int[x^(Simplify[(m+1)/n]-1)*(a+b*ArcTan[c*x])^p,x],x,x^n] /;
FreeQ[{a,b,c,m,n},x] && IGtQ[p,1] && IntegerQ[Simplify[(m+1)/n]]
```

```
Int[x_^m_.*(a_.*b_.*ArcCot[c_.*x_^n_])^p_.,x_Symbol] :=
  1/n*Subst[Int[x^(Simplify[(m+1)/n]-1)*(a+b*ArcCot[c*x])^p,x],x,x^n] /;
FreeQ[{a,b,c,m,n},x] && IGtQ[p,1] && IntegerQ[Simplify[(m+1)/n]]
```

$$4. \int x^m (a + b \operatorname{ArcTan}[c x^n])^p dx \text{ when } p - 1 \in \mathbb{Z}^+ \wedge n \in \mathbb{Z}$$

$$1. \int x^m (a + b \operatorname{ArcTan}[c x^n])^p dx \text{ when } p - 1 \in \mathbb{Z}^+ \wedge n \in \mathbb{Z}^+$$

$$1: \int x^m (a + b \operatorname{ArcTan}[c x^n])^p dx \text{ when } p - 1 \in \mathbb{Z}^+ \wedge n \in \mathbb{Z}^+ \wedge m \in \mathbb{Z}$$

Derivation: Algebraic expansion

$$\text{Basis: } \operatorname{ArcTan}[z] == \frac{i \operatorname{Log}[1 - iz]}{2} - \frac{i \operatorname{Log}[1 + iz]}{2}$$

$$\text{Basis: } \operatorname{ArcCot}[z] == \frac{i \operatorname{Log}[1 - iz^{-1}]}{2} - \frac{i \operatorname{Log}[1 + iz^{-1}]}{2}$$

Rule: If $p - 1 \in \mathbb{Z}^+ \wedge n \in \mathbb{Z}^+ \wedge m \in \mathbb{Z}$, then

$$\int x^m (a + b \operatorname{ArcTan}[c x^n])^p dx \rightarrow \int \operatorname{ExpandIntegrand}\left[x^m \left(a + \frac{i b \operatorname{Log}[1 - i c x^n]}{2} - \frac{i b \operatorname{Log}[1 + i c x^n]}{2}\right)^p, x\right] dx$$

Program code:

```
Int[x^m.*(a_.+b_.*ArcTan[c_.*x^n_])^p_,x_Symbol] :=
  Int[ExpandIntegrand[x^m*(a+(I*b*Log[1-I*c*x^n])/2-(I*b*Log[1+I*c*x^n])/2]^p,x],x] /;
FreeQ[{a,b,c},x] && IGtQ[p,1] && IGtQ[n,0] && IntegerQ[m]
```

```
Int[x^m.*(a_.+b_.*ArcCot[c_.*x^n_])^p_,x_Symbol] :=
  Int[ExpandIntegrand[x^m*(a+(I*b*Log[1-I*x^(-n)/c])/2-(I*b*Log[1+I*x^(-n)/c])/2]^p,x],x] /;
FreeQ[{a,b,c},x] && IGtQ[p,1] && IGtQ[n,0] && IntegerQ[m]
```

$$2: \int x^m (a + b \operatorname{ArcTan}[c x^n])^p dx \text{ when } p - 1 \in \mathbb{Z}^+ \wedge n \in \mathbb{Z}^+ \wedge m \in \mathbb{F}$$

Derivation: Integration by substitution

Basis: If $k \in \mathbb{Z}^+$, then $F[x] = k \operatorname{Subst}[x^{k-1} F[x^k], x, x^{1/k}] \partial_x x^{1/k}$

Rule: If $p - 1 \in \mathbb{Z}^+ \wedge n \in \mathbb{Z}^+ \wedge m \in \mathbb{F}$, let $k \rightarrow \operatorname{Denominator}[m]$, then

$$\int x^m (a + b \operatorname{ArcTan}[c x^n])^p dx \rightarrow k \operatorname{Subst}\left[\int x^{k(m+1)-1} (a + b \operatorname{ArcTan}[c x^{kn}])^p dx, x, x^{1/k}\right]$$

Program code:

```
Int[x^m.*(a_.+b_.*ArcTan[c_.*x_^n_])^p_,x_Symbol] :=
  With[{k=Denominator[m]},
    k*Subst[Int[x^(k*(m+1)-1)*(a+b*ArcTan[c*x^(k*n)])^p,x],x,x^(1/k)] /;
  FreeQ[{a,b,c},x] && IGtQ[p,1] && IGtQ[n,0] && FractionQ[m]
```

```
Int[x^m.*(a_.+b_.*ArcCot[c_.*x_^n_])^p_,x_Symbol] :=
  With[{k=Denominator[m]},
    k*Subst[Int[x^(k*(m+1)-1)*(a+b*ArcCot[c*x^(k*n)])^p,x],x,x^(1/k)] /;
  FreeQ[{a,b,c},x] && IGtQ[p,1] && IGtQ[n,0] && FractionQ[m]
```

$$2: \int x^m (a + b \operatorname{ArcTan}[c x^n])^p dx \text{ when } p - 1 \in \mathbb{Z}^+ \wedge n \in \mathbb{Z}^-$$

Derivation: Algebraic simplification

$$\text{Basis: } \operatorname{ArcTan}[z] == \operatorname{ArcCot}\left[\frac{1}{z}\right]$$

Rule: If $p - 1 \in \mathbb{Z}^+ \wedge n \in \mathbb{Z}^-$, then

$$\int x^m (a + b \operatorname{ArcTan}[c x^n])^p dx \rightarrow \int x^m \left(a + b \operatorname{ArcCot}\left[\frac{x^{-n}}{c}\right]\right)^p dx$$

Program code:

```
Int[x^m_.*(a_.+b_.*ArcTan[c_.*x_^n_])^p_,x_Symbol] :=
  Int[x^m*(a+b*ArcCot[x^(-n)/c])^p,x] /;
FreeQ[{a,b,c},x] && IGtQ[p,1] && ILtQ[n,0]
```

```
Int[x^m_.*(a_.+b_.*ArcCot[c_.*x_^n_])^p_,x_Symbol] :=
  Int[x^m*(a+b*ArcTan[x^(-n)/c])^p,x] /;
FreeQ[{a,b,c},x] && IGtQ[p,1] && ILtQ[n,0]
```


5: $\int x^m (a + b \operatorname{ArcTan}[c x^n])^p dx$ when $p - 1 \in \mathbb{Z}^+ \wedge n \in \mathbb{F}$

Derivation: Integration by substitution

Basis: If $k \in \mathbb{Z}^+$, then $F[x] = k \operatorname{Subst}[x^{k-1} F[x^k], x, x^{1/k}] \partial_x x^{1/k}$

Rule: If $p - 1 \in \mathbb{Z}^+ \wedge n \in \mathbb{F}$, let $k \rightarrow \operatorname{Denominator}[n]$, then

$$\int x^m (a + b \operatorname{ArcTan}[c x^n])^p dx \rightarrow k \operatorname{Subst}\left[\int x^{k(m+1)-1} (a + b \operatorname{ArcTan}[c x^{kn}])^p dx, x, x^{1/k}\right]$$

Program code:

```
Int[x^m_.*(a_.+b_.*ArcTan[c_*x_^n_])^p_,x_Symbol] :=
  With[{k=Denominator[n]},
    k*Subst[Int[x^(k*(m+1)-1)*(a+b*ArcTan[c*x^(k*n)])^p,x],x,x^(1/k)] /;
  FreeQ[{a,b,c},x] && IGtQ[p,1] && FractionQ[n]
```

```
Int[x^m_.*(a_.+b_.*ArcCot[c_*x_^n_])^p_,x_Symbol] :=
  With[{k=Denominator[n]},
    k*Subst[Int[x^(k*(m+1)-1)*(a+b*ArcCot[c*x^(k*n)])^p,x],x,x^(1/k)] /;
  FreeQ[{a,b,c},x] && IGtQ[p,1] && FractionQ[n]
```

$$2: \int (dx)^m (a + b \operatorname{ArcTan}[cx^n]) dx \text{ when } n \in \mathbb{Z} \wedge m \neq -1$$

Derivation: Integration by parts

$$\text{Basis: If } n \in \mathbb{Z}, \text{ then } \partial_x (a + b \operatorname{ArcTan}[cx^n]) = \frac{bcn(dx)^{n-1}}{d^{n-1}(1+c^2x^{2n})}$$

Rule: If $n \in \mathbb{Z} \wedge m \neq -1$, then

$$\int (dx)^m (a + b \operatorname{ArcTan}[cx^n]) dx \rightarrow \frac{(dx)^{m+1} (a + b \operatorname{ArcTan}[cx^n])}{d(m+1)} - \frac{bcn}{d^n(m+1)} \int \frac{(dx)^{m+n}}{1+c^2x^{2n}} dx$$

Program code:

```
Int[(d*x_)^m*(a_.+b_.*ArcTan[c_.*x_^n_.]),x_Symbol] :=
  (d*x)^(m+1)*(a+b*ArcTan[c*x^n])/(d*(m+1)) -
  b*c*n/(d^n*(m+1))*Int[(d*x)^(m+n)/(1+c^2*x^(2*n)),x] /;
FreeQ[{a,b,c,d,m,n},x] && IntegerQ[n] && NeQ[m,-1]
```

```
Int[(d*x_)^m*(a_.+b_.*ArcCot[c_.*x_^n_.]),x_Symbol] :=
  (d*x)^(m+1)*(a+b*ArcCot[c*x^n])/(d*(m+1)) +
  b*c*n/(d^n*(m+1))*Int[(d*x)^(m+n)/(1+c^2*x^(2*n)),x] /;
FreeQ[{a,b,c,d,m,n},x] && IntegerQ[n] && NeQ[m,-1]
```

$$3: \int (dx)^m (a + b \operatorname{ArcTan}[cx^n])^p dx \text{ when } p \in \mathbb{Z}^+ \wedge (p = 1 \vee m \in \mathbb{R} \wedge n \in \mathbb{R})$$

Derivation: Piecewise constant extraction

$$\text{Basis: } a_x \frac{(dx)^m}{x^m} = 0$$

Rule: If $p \in \mathbb{Z}^+ \wedge (p = 1 \vee m \in \mathbb{R} \wedge n \in \mathbb{R})$, then

$$\int (dx)^m (a + b \operatorname{ArcTan}[cx^n])^p dx \rightarrow \frac{d^{\operatorname{IntPart}[m]} (dx)^{\operatorname{FracPart}[m]}}{x^{\operatorname{FracPart}[m]}} \int x^m (a + b \operatorname{ArcTan}[cx^n])^p dx$$

Program code:

```
Int[(d_.**x_)^m_*(a_.+b_.*ArcTan[c_.**x_^n_])^p_,x_Symbol] :=
  d^IntPart[m]*(d*x)^FracPart[m]/x^FracPart[m]*Int[x^m*(a+b*ArcTan[c*x])^p,x] /;
FreeQ[{a,b,c,d,m,n},x] && IGtQ[p,0] && (EqQ[p,1] || RationalQ[m,n])
```

```
Int[(d_.**x_)^m_*(a_.+b_.*ArcCot[c_.**x_^n_])^p_,x_Symbol] :=
  d^IntPart[m]*(d*x)^FracPart[m]/x^FracPart[m]*Int[x^m*(a+b*ArcCot[c*x])^p,x] /;
FreeQ[{a,b,c,d,m,n},x] && IGtQ[p,0] && (EqQ[p,1] || RationalQ[m,n])
```

$$U: \int (dx)^m (a + b \operatorname{ArcTan}[cx^n])^p dx$$

Rule:

$$\int (dx)^m (a + b \operatorname{ArcTan}[cx^n])^p dx \rightarrow \int (dx)^m (a + b \operatorname{ArcTan}[cx^n])^p dx$$

Program code:

```
Int[(d_.**x_)^m_*(a_.+b_.*ArcTan[c_.**x_^n_])^p_,x_Symbol] :=
  Unintegrable[(d*x)^m*(a+b*ArcTan[c*x^n])^p,x] /;
FreeQ[{a,b,c,d,m,n,p},x]
```

```
Int[(d.*x_)^m.*(a_.+b_.*ArcCot[c_.*x_^n_.])^p_,x_Symbol] :=  
  Unintegrable[(d*x)^m*(a+b*ArcCot[c*x^n])^p,x] /;  
FreeQ[{a,b,c,d,m,n,p},x]
```