

Rules for integrands of the form $(d + ex)^q (a + b \operatorname{ArcTan}[cx])^p$

1. $\int (d + ex)^q (a + b \operatorname{ArcTan}[cx])^p dx$ when $p \in \mathbb{Z}^+$

1. $\int \frac{(a + b \operatorname{ArcTan}[cx])^p}{d + ex} dx$ when $p \in \mathbb{Z}^+$

1: $\int \frac{(a + b \operatorname{ArcTan}[cx])^p}{d + ex} dx$ when $p \in \mathbb{Z}^+ \wedge c^2 d^2 + e^2 = 0$

Derivation: Integration by parts

Basis: $\frac{1}{d+ex} = -\frac{1}{e} \partial_x \operatorname{Log}\left[\frac{2}{1+\frac{ex}{d}}\right]$

Rule: If $p \in \mathbb{Z}^+ \wedge c^2 d^2 + e^2 = 0$, then

$$\int \frac{(a + b \operatorname{ArcTan}[cx])^p}{d + ex} dx \rightarrow -\frac{(a + b \operatorname{ArcTan}[cx])^p \operatorname{Log}\left[\frac{2}{1+\frac{ex}{d}}\right]}{e} + \frac{bc p}{e} \int \frac{(a + b \operatorname{ArcTan}[cx])^{p-1} \operatorname{Log}\left[\frac{2}{1+\frac{ex}{d}}\right]}{1 + c^2 x^2} dx$$

Program code:

```
Int[(a_.*b_.*ArcTan[c.*x_])^p_./(d_+e_.*x_),x_Symbol] :=
  -(a+b*ArcTan[c*x])^p*Log[2/(1+e*x/d)]/e +
  b*c*p/e*Int[(a+b*ArcTan[c*x])^(p-1)*Log[2/(1+e*x/d)]/(1+c^2*x^2),x] /;
FreeQ[{a,b,c,d,e},x] && IGtQ[p,0] && EqQ[c^2*d^2+e^2,0]
```

```
Int[(a_.*b_.*ArcCot[c.*x_])^p_./(d_+e_.*x_),x_Symbol] :=
  -(a+b*ArcCot[c*x])^p*Log[2/(1+e*x/d)]/e -
  b*c*p/e*Int[(a+b*ArcCot[c*x])^(p-1)*Log[2/(1+e*x/d)]/(1+c^2*x^2),x] /;
FreeQ[{a,b,c,d,e},x] && IGtQ[p,0] && EqQ[c^2*d^2+e^2,0]
```

$$2. \int \frac{(a + b \operatorname{ArcTan}[c x])^p}{d + e x} dx \text{ when } p \in \mathbb{Z}^+ \wedge c^2 d^2 + e^2 \neq 0$$

$$1: \int \frac{a + b \operatorname{ArcTan}[c x]}{d + e x} dx \text{ when } c^2 d^2 + e^2 \neq 0$$

Derivation: Algebraic expansion and integration by parts

$$\text{Basis: } \frac{1}{d+ex} = \frac{c}{e(i+cx)} - \frac{cd-ie}{e(i+cx)(d+ex)}$$

$$\text{Basis: } \frac{1}{i+cx} = -\frac{1}{c} \partial_x \operatorname{Log} \left[\frac{2}{1-icx} \right]$$

$$\text{Basis: } \frac{1}{(i+cx)(d+ex)} = -\frac{1}{cd-ie} \partial_x \operatorname{Log} \left[\frac{2c(d+ex)}{(cd+ie)(1-icx)} \right]$$

$$\text{Basis: } \partial_x (a + b \operatorname{ArcTan}[c x]) = \frac{bc}{1+c^2x^2}$$

Rule: If $c^2 d^2 + e^2 \neq 0$, then

$$\begin{aligned} \int \frac{a + b \operatorname{ArcTan}[c x]}{d + e x} dx &\rightarrow \frac{c}{e} \int \frac{a + b \operatorname{ArcTan}[c x]}{i + c x} dx - \frac{cd - ie}{e} \int \frac{a + b \operatorname{ArcTan}[c x]}{(i + c x)(d + e x)} dx \rightarrow \\ &-\frac{(a + b \operatorname{ArcTan}[c x]) \operatorname{Log} \left[\frac{2}{1-icx} \right]}{e} + \frac{bc}{e} \int \frac{\operatorname{Log} \left[\frac{2}{1-icx} \right]}{1 + c^2 x^2} dx + \frac{(a + b \operatorname{ArcTan}[c x]) \operatorname{Log} \left[\frac{2c(d+ex)}{(cd+ie)(1-icx)} \right]}{e} - \frac{bc}{e} \int \frac{\operatorname{Log} \left[\frac{2c(d+ex)}{(cd+ie)(1-icx)} \right]}{1 + c^2 x^2} dx \rightarrow \\ &-\frac{(a + b \operatorname{ArcTan}[c x]) \operatorname{Log} \left[\frac{2}{1-icx} \right]}{e} + \frac{ib \operatorname{PolyLog} \left[2, 1 - \frac{2}{1-icx} \right]}{2e} + \frac{(a + b \operatorname{ArcTan}[c x]) \operatorname{Log} \left[\frac{2c(d+ex)}{(cd+ie)(1-icx)} \right]}{e} - \frac{ib \operatorname{PolyLog} \left[2, 1 - \frac{2c(d+ex)}{(cd+ie)(1-icx)} \right]}{2e} \end{aligned}$$

Program code:

```
Int[(a_+b_.*ArcTan[c_.*x_])/(d_+e_.*x_),x_Symbol] :=
-(a+b*ArcTan[c*x])*Log[2/(1-I*c*x)]/e +
b*c/e*Int[Log[2/(1-I*c*x)]/(1+c^2*x^2),x] +
(a+b*ArcTan[c*x])*Log[2*c*(d+e*x)/((c*d+I*e)*(1-I*c*x))]/e -
b*c/e*Int[Log[2*c*(d+e*x)/((c*d+I*e)*(1-I*c*x))]/(1+c^2*x^2),x] /;
FreeQ[{a,b,c,d,e},x] && NeQ[c^2*d^2+e^2,0]
```

```

Int[(a_.+b_.*ArcCot[c_.*x_])/(d_.+e_.*x_),x_Symbol] :=
- (a+b*ArcCot[c*x]) * Log[2/(1-I*c*x)]/e -
b*c/e*Int[Log[2/(1-I*c*x)]/(1+c^2*x^2),x] +
(a+b*ArcCot[c*x]) * Log[2*c*(d+e*x)/((c*d+I*e)*(1-I*c*x))]/e +
b*c/e*Int[Log[2*c*(d+e*x)/((c*d+I*e)*(1-I*c*x))]/(1+c^2*x^2),x] /;
FreeQ[{a,b,c,d,e},x] && NeQ[c^2*d^2+e^2,0]

```

$$2: \int \frac{(a + b \operatorname{ArcTan}[c x])^2}{d + e x} dx \text{ when } c^2 d^2 + e^2 \neq 0$$

Derivation: Algebraic expansion and integration by parts

$$\text{Basis: } \frac{1}{d+ex} = \frac{c}{e(i+cx)} - \frac{cd-ie}{e(i+cx)(d+ex)}$$

$$\text{Basis: } \frac{1}{i+cx} = -\frac{1}{c} \partial_x \operatorname{Log}\left[\frac{2}{1-icx}\right]$$

$$\text{Basis: } \frac{1}{(i+cx)(d+ex)} = -\frac{1}{cd-ie} \partial_x \operatorname{Log}\left[\frac{2c(d+ex)}{(cd+ie)(1-icx)}\right]$$

$$\text{Basis: } \partial_x (a + b \operatorname{ArcTan}[c x])^2 = \frac{2bc(a+b \operatorname{ArcTan}[c x])}{1+c^2 x^2}$$

Rule: If $c^2 d^2 + e^2 \neq 0$, then

$$\begin{aligned} \int \frac{(a + b \operatorname{ArcTan}[c x])^2}{d + e x} dx &\rightarrow \frac{c}{e} \int \frac{(a + b \operatorname{ArcTan}[c x])^2}{i + c x} dx - \frac{cd - ie}{e} \int \frac{(a + b \operatorname{ArcTan}[c x])^2}{(i + c x)(d + e x)} dx \rightarrow \\ &-\frac{(a + b \operatorname{ArcTan}[c x])^2 \operatorname{Log}\left[\frac{2}{1-icx}\right]}{e} + \frac{2bc}{e} \int \frac{(a + b \operatorname{ArcTan}[c x]) \operatorname{Log}\left[\frac{2}{1-icx}\right]}{1 + c^2 x^2} dx + \\ &\frac{(a + b \operatorname{ArcTan}[c x])^2 \operatorname{Log}\left[\frac{2c(d+ex)}{(cd+ie)(1-icx)}\right]}{e} - \frac{2bc}{e} \int \frac{(a + b \operatorname{ArcTan}[c x]) \operatorname{Log}\left[\frac{2c(d+ex)}{(cd+ie)(1-icx)}\right]}{1 + c^2 x^2} dx \rightarrow \\ &-\frac{(a + b \operatorname{ArcTan}[c x])^2 \operatorname{Log}\left[\frac{2}{1-icx}\right]}{e} + \frac{ib(a + b \operatorname{ArcTan}[c x]) \operatorname{PolyLog}\left[2, 1 - \frac{2}{1-icx}\right]}{e} - \frac{b^2 \operatorname{PolyLog}\left[3, 1 - \frac{2}{1-icx}\right]}{2e} + \\ &\frac{(a + b \operatorname{ArcTan}[c x])^2 \operatorname{Log}\left[\frac{2c(d+ex)}{(cd+ie)(1-icx)}\right]}{e} - \frac{ib(a + b \operatorname{ArcTan}[c x]) \operatorname{PolyLog}\left[2, 1 - \frac{2c(d+ex)}{(cd+ie)(1-icx)}\right]}{e} + \frac{b^2 \operatorname{PolyLog}\left[3, 1 - \frac{2c(d+ex)}{(cd+ie)(1-icx)}\right]}{2e} \end{aligned}$$

Program code:

```

Int [(a_+b_.*ArcTan[c_.*x_])^2/(d_+e_.*x_),x_Symbol] :=
- (a+b*ArcTan[c*x])^2*Log[2/(1-I*c*x)]/e +
I*b*(a+b*ArcTan[c*x])*PolyLog[2,1-2/(1-I*c*x)]/e -
b^2*PolyLog[3,1-2/(1-I*c*x)]/(2*e) +
(a+b*ArcTan[c*x])^2*Log[2*c*(d+e*x)/((c*d+I*e)*(1-I*c*x))]/e -
I*b*(a+b*ArcTan[c*x])*PolyLog[2,1-2*c*(d+e*x)/((c*d+I*e)*(1-I*c*x))]/e +
b^2*PolyLog[3,1-2*c*(d+e*x)/((c*d+I*e)*(1-I*c*x))]/(2*e) /;
FreeQ[{a,b,c,d,e},x] && NeQ[c^2*d^2+e^2,0]

```

```

Int [(a_+b_.*ArcCot[c_.*x_])^2/(d_+e_.*x_),x_Symbol] :=
- (a+b*ArcCot[c*x])^2*Log[2/(1-I*c*x)]/e -
I*b*(a+b*ArcCot[c*x])*PolyLog[2,1-2/(1-I*c*x)]/e -
b^2*PolyLog[3,1-2/(1-I*c*x)]/(2*e) +
(a+b*ArcCot[c*x])^2*Log[2*c*(d+e*x)/((c*d+I*e)*(1-I*c*x))]/e +
I*b*(a+b*ArcCot[c*x])*PolyLog[2,1-2*c*(d+e*x)/((c*d+I*e)*(1-I*c*x))]/e +
b^2*PolyLog[3,1-2*c*(d+e*x)/((c*d+I*e)*(1-I*c*x))]/(2*e) /;
FreeQ[{a,b,c,d,e},x] && NeQ[c^2*d^2+e^2,0]

```

$$3: \int \frac{(a + b \operatorname{ArcTan}[c x])^3}{d + e x} dx \text{ when } c^2 d^2 + e^2 \neq 0$$

Derivation: Algebraic expansion and integration by parts

$$\text{Basis: } \frac{1}{d+ex} \equiv \frac{c}{e(i+cx)} - \frac{cd-ie}{e(i+cx)(d+ex)}$$

$$\text{Basis: } \frac{1}{i+cx} \equiv -\frac{1}{c} \partial_x \operatorname{Log} \left[\frac{2}{1-icx} \right]$$

$$\text{Basis: } \frac{1}{(i+cx)(d+ex)} \equiv -\frac{1}{cd-ie} \partial_x \operatorname{Log} \left[\frac{2c(d+ex)}{(cd+ie)(1-icx)} \right]$$

$$\text{Basis: } \partial_x (a + b \operatorname{ArcTan}[c x])^3 \equiv \frac{3bc(a+b \operatorname{ArcTan}[c x])^2}{1+c^2 x^2}$$

Rule: If $c^2 d^2 + e^2 \neq 0$, then

$$\int \frac{(a + b \operatorname{ArcTan}[c x])^3}{d + e x} dx \rightarrow \frac{c}{e} \int \frac{(a + b \operatorname{ArcTan}[c x])^3}{i + c x} dx - \frac{cd - ie}{e} \int \frac{(a + b \operatorname{ArcTan}[c x])^3}{(i + c x)(d + e x)} dx \rightarrow$$

$$\begin{aligned}
& -\frac{(a+b \operatorname{ArcTan}[c x])^3 \operatorname{Log}\left[\frac{2}{1-i c x}\right]}{e} + \frac{3 b c}{e} \int \frac{(a+b \operatorname{ArcTan}[c x])^2 \operatorname{Log}\left[\frac{2}{1-i c x}\right]}{1+c^2 x^2} dx + \\
& \frac{(a+b \operatorname{ArcTan}[c x])^3 \operatorname{Log}\left[\frac{2 c(d+e x)}{(c d+i e)(1-i c x)}\right]}{e} - \frac{3 b c}{e} \int \frac{(a+b \operatorname{ArcTan}[c x])^2 \operatorname{Log}\left[\frac{2 c(d+e x)}{(c d+i e)(1-i c x)}\right]}{1+c^2 x^2} dx \rightarrow \\
& -\frac{(a+b \operatorname{ArcTan}[c x])^3 \operatorname{Log}\left[\frac{2}{1-i c x}\right]}{e} + \frac{3 i b(a+b \operatorname{ArcTan}[c x])^2 \operatorname{PolyLog}\left[2, 1-\frac{2}{1-i c x}\right]}{2 e} - \\
& \frac{3 b^2(a+b \operatorname{ArcTan}[c x]) \operatorname{PolyLog}\left[3, 1-\frac{2}{1-i c x}\right]}{2 e} - \frac{3 i b^3 \operatorname{PolyLog}\left[4, 1-\frac{2}{1-i c x}\right]}{4 e} + \\
& \frac{(a+b \operatorname{ArcTan}[c x])^3 \operatorname{Log}\left[\frac{2 c(d+e x)}{(c d+i e)(1-i c x)}\right]}{e} - \frac{3 i b(a+b \operatorname{ArcTan}[c x])^2 \operatorname{PolyLog}\left[2, 1-\frac{2 c(d+e x)}{(c d+i e)(1-i c x)}\right]}{2 e} + \\
& \frac{3 b^2(a+b \operatorname{ArcTan}[c x]) \operatorname{PolyLog}\left[3, 1-\frac{2 c(d+e x)}{(c d+i e)(1-i c x)}\right]}{2 e} + \frac{3 i b^3 \operatorname{PolyLog}\left[4, 1-\frac{2 c(d+e x)}{(c d+i e)(1-i c x)}\right]}{4 e}
\end{aligned}$$

Program code:

```

Int[(a_.+b_.*ArcTan[c.*x_])^3/(d_+e_.x_),x_Symbol] :=
-(a+b*ArcTan[c*x])^3*Log[2/(1-I*c*x)]/e +
3*I*b*(a+b*ArcTan[c*x])^2*PolyLog[2,1-2/(1-I*c*x)]/(2*e) -
3*b^2*(a+b*ArcTan[c*x])*PolyLog[3,1-2/(1-I*c*x)]/(2*e) -
3*I*b^3*PolyLog[4,1-2/(1-I*c*x)]/(4*e) +
(a+b*ArcTan[c*x])^3*Log[2*c*(d+e*x)/((c*d+I*e)*(1-I*c*x))]/e -
3*I*b*(a+b*ArcTan[c*x])^2*PolyLog[2,1-2*c*(d+e*x)/((c*d+I*e)*(1-I*c*x))]/(2*e) +
3*b^2*(a+b*ArcTan[c*x])*PolyLog[3,1-2*c*(d+e*x)/((c*d+I*e)*(1-I*c*x))]/(2*e) +
3*I*b^3*PolyLog[4,1-2*c*(d+e*x)/((c*d+I*e)*(1-I*c*x))]/(4*e) /;
FreeQ[{a,b,c,d,e},x] && NeQ[c^2*d^2+e^2,0]

```

```

Int[(a_.+b_.*ArcCot[c.*x_])^3/(d_+e_.x_),x_Symbol] :=
-(a+b*ArcCot[c*x])^3*Log[2/(1-I*c*x)]/e -
3*I*b*(a+b*ArcCot[c*x])^2*PolyLog[2,1-2/(1-I*c*x)]/(2*e) -
3*b^2*(a+b*ArcCot[c*x])*PolyLog[3,1-2/(1-I*c*x)]/(2*e) +
3*I*b^3*PolyLog[4,1-2/(1-I*c*x)]/(4*e) +
(a+b*ArcCot[c*x])^3*Log[2*c*(d+e*x)/((c*d+I*e)*(1-I*c*x))]/e +
3*I*b*(a+b*ArcCot[c*x])^2*PolyLog[2,1-2*c*(d+e*x)/((c*d+I*e)*(1-I*c*x))]/(2*e) +
3*b^2*(a+b*ArcCot[c*x])*PolyLog[3,1-2*c*(d+e*x)/((c*d+I*e)*(1-I*c*x))]/(2*e) -
3*I*b^3*PolyLog[4,1-2*c*(d+e*x)/((c*d+I*e)*(1-I*c*x))]/(4*e) /;
FreeQ[{a,b,c,d,e},x] && NeQ[c^2*d^2+e^2,0]

```

2: $\int (d+ex)^q (a+b \operatorname{ArcTan}[cx]) dx$ when $q \neq -1$

Derivation: Integration by parts

Rule: If $q \neq -1$, then

$$\int (d+ex)^q (a+b \operatorname{ArcTan}[cx]) dx \rightarrow \frac{(d+ex)^{q+1} (a+b \operatorname{ArcTan}[cx])}{e(q+1)} - \frac{bc}{e(q+1)} \int \frac{(d+ex)^{q+1}}{1+c^2x^2} dx$$

Program code:

```
Int[(d+_e*_x_)^q_.*(a+_b_.*ArcTan[c*_x_]),x_Symbol] :=
  (d+e*x)^(q+1)*(a+b*ArcTan[c*x])/(e*(q+1)) -
  b*c/(e*(q+1))*Int[(d+e*x)^(q+1)/(1+c^2*x^2),x] /;
FreeQ[{a,b,c,d,e,q},x] && NeQ[q,-1]
```

```
Int[(d+_e*_x_)^q_.*(a+_b_.*ArcCot[c*_x_]),x_Symbol] :=
  (d+e*x)^(q+1)*(a+b*ArcCot[c*x])/(e*(q+1)) +
  b*c/(e*(q+1))*Int[(d+e*x)^(q+1)/(1+c^2*x^2),x] /;
FreeQ[{a,b,c,d,e,q},x] && NeQ[q,-1]
```

3: $\int (d+ex)^q (a+b \operatorname{ArcTan}[cx])^p dx$ when $p-1 \in \mathbb{Z}^+ \wedge q \in \mathbb{Z} \wedge q \neq -1$

Derivation: Integration by parts

Rule: If $p-1 \in \mathbb{Z}^+ \wedge q \in \mathbb{Z} \wedge q \neq -1$, then

$$\int (d+ex)^q (a+b \operatorname{ArcTan}[cx])^p dx \rightarrow \frac{(d+ex)^{q+1} (a+b \operatorname{ArcTan}[cx])^p}{e(q+1)} - \frac{bc p}{e(q+1)} \int (a+b \operatorname{ArcTan}[cx])^{p-1} \operatorname{ExpandIntegrand}\left[\frac{(d+ex)^{q+1}}{1+c^2 x^2}, x\right] dx$$

Program code:

```
Int[(d+_e*_x_)^q_.*(a+_b*_ArcTan[c*_x_])^p_,x_Symbol] :=
  (d+e*x)^(q+1)*(a+b*ArcTan[c*x])^p/(e*(q+1)) -
  b*c*p/(e*(q+1))*Int[ExpandIntegrand[(a+b*ArcTan[c*x])^(p-1),(d+e*x)^(q+1)/(1+c^2*x^2),x],x] /;
FreeQ[{a,b,c,d,e},x] && IGtQ[p,1] && IntegerQ[q] && NeQ[q,-1]
```

```
Int[(d+_e*_x_)^q_.*(a+_b*_ArcCot[c*_x_])^p_,x_Symbol] :=
  (d+e*x)^(q+1)*(a+b*ArcCot[c*x])^p/(e*(q+1)) +
  b*c*p/(e*(q+1))*Int[ExpandIntegrand[(a+b*ArcCot[c*x])^(p-1),(d+e*x)^(q+1)/(1+c^2*x^2),x],x] /;
FreeQ[{a,b,c,d,e},x] && IGtQ[p,1] && IntegerQ[q] && NeQ[q,-1]
```


$$2. \int (d+ex)^m (a+b \operatorname{ArcTan}[cx^n])^p dx \text{ when } p \in \mathbb{Z}^+$$

$$1. \int (d+ex)^m (a+b \operatorname{ArcTan}[cx^n]) dx$$

$$1. \int \frac{a+b \operatorname{ArcTan}[cx^n]}{d+ex} dx$$

$$1: \int \frac{a+b \operatorname{ArcTan}[cx^n]}{d+ex} dx \text{ when } n \in \mathbb{Z}$$

Derivation: Integration by parts

$$\text{Basis: } \partial_x (a+b \operatorname{ArcTan}[cx^n]) = bcn \frac{x^{n-1}}{1+c^2 x^{2n}}$$

Rule: If $n \in \mathbb{Z}$, then

$$\int \frac{a+b \operatorname{ArcTan}[cx^n]}{d+ex} dx \rightarrow \frac{\operatorname{Log}[d+ex] (a+b \operatorname{ArcTan}[cx^n])}{e} - \frac{bcn}{e} \int \frac{x^{n-1} \operatorname{Log}[d+ex]}{1+c^2 x^{2n}} dx$$

Program code:

```
Int[(a.+b_.*ArcTan[c.*x^n_)]/(d.+e.*x_),x_Symbol] :=
  Log[d+e*x]*(a+b*ArcTan[c*x^n])/e -
  b*c*n/e*Int[x^(n-1)*Log[d+e*x]/(1+c^2*x^(2*n)),x] /;
FreeQ[{a,b,c,d,e,n},x] && IntegerQ[n]
```

```
Int[(a.+b_.*ArcCot[c.*x^n_)]/(d.+e.*x_),x_Symbol] :=
  Log[d+e*x]*(a+b*ArcCot[c*x^n])/e +
  b*c*n/e*Int[x^(n-1)*Log[d+e*x]/(1+c^2*x^(2*n)),x] /;
FreeQ[{a,b,c,d,e,n},x] && IntegerQ[n]
```

$$2: \int \frac{a + b \operatorname{ArcTan}[c x^n]}{d + e x} dx \text{ when } n \in \mathbb{F}$$

Derivation: Integration by substitution

Basis: If $k \in \mathbb{Z}^+$, then $F[x] = k \operatorname{Subst}[x^{k-1} F[x^k], x, x^{1/k}] \partial_x x^{1/k}$

Rule: If $n \in \mathbb{F}$, let $k \rightarrow \operatorname{Denominator}[n]$, then

$$\int \frac{a + b \operatorname{ArcTan}[c x^n]}{d + e x} dx \rightarrow k \operatorname{Subst}\left[\int \frac{x^{k-1} (a + b \operatorname{ArcTan}[c x^{kn}])}{d + e x^k} dx, x, x^{1/k}\right]$$

Program code:

```
Int[(a_.+b_.*ArcTan[c_.*x^n_])/(d_.+e_.*x_),x_Symbol] :=
  With[{k=Denominator[n]},
    k*Subst[Int[x^(k-1)*(a+b*ArcTan[c*x^(k*n)])/(d+e*x^k),x],x,x^(1/k)]] /;
  FreeQ[{a,b,c,d,e},x] && FractionQ[n]
```

```
Int[(a_.+b_.*ArcCot[c_.*x^n_])/(d_.+e_.*x_),x_Symbol] :=
  With[{k=Denominator[n]},
    k*Subst[Int[x^(k-1)*(a+b*ArcCot[c*x^(k*n)])/(d+e*x^k),x],x,x^(1/k)]] /;
  FreeQ[{a,b,c,d,e},x] && FractionQ[n]
```

$$2: \int (d+ex)^m (a+b \operatorname{ArcTan}[cx^n]) dx \text{ when } m \neq -1$$

Derivation: Integration by parts

$$\text{Basis: } \partial_x (a+b \operatorname{ArcTan}[cx^n]) = bc n \frac{x^{n-1}}{1+c^2 x^{2n}}$$

Rule: If $m \neq -1$, then

$$\int (d+ex)^m (a+b \operatorname{ArcTan}[cx^n]) dx \rightarrow \frac{(d+ex)^{m+1} (a+b \operatorname{ArcTan}[cx^n])}{e(m+1)} - \frac{bc n}{e(m+1)} \int \frac{x^{n-1} (d+ex)^{m+1}}{1+c^2 x^{2n}} dx$$

Program code:

```
Int[(d+_e*_x_)^m_.*(a+_b_.*ArcTan[c*_x_^n_]),x_Symbol] :=
  (d+e*x)^(m+1)*(a+b*ArcTan[c*x^n])/(e*(m+1)) -
  b*c*n/(e*(m+1))*Int[x^(n-1)*(d+e*x)^(m+1)/(1+c^2*x^(2*n)),x] /;
FreeQ[{a,b,c,d,e,m,n},x] && NeQ[m,-1]
```

```
Int[(d+_e*_x_)^m_.*(a+_b_.*ArcCot[c*_x_^n_]),x_Symbol] :=
  (d+e*x)^(m+1)*(a+b*ArcCot[c*x^n])/(e*(m+1)) +
  b*c*n/(e*(m+1))*Int[x^(n-1)*(d+e*x)^(m+1)/(1+c^2*x^(2*n)),x] /;
FreeQ[{a,b,c,d,e,m,n},x] && NeQ[m,-1]
```

2: $\int (d+ex)^m (a+b \operatorname{ArcTan}[cx^n])^p dx$ when $p-1 \in \mathbb{Z}^+ \wedge m \in \mathbb{Z}^+$

Derivation: Algebraic expansion

Rule: If $p-1 \in \mathbb{Z}^+ \wedge m \in \mathbb{Z}^+$, then

$$\int (d+ex)^m (a+b \operatorname{ArcTan}[cx^n])^p dx \rightarrow \int (a+b \operatorname{ArcTan}[cx^n])^p \operatorname{ExpandIntegrand}[(d+ex)^m, x] dx$$

Program code:

```
Int[(d_+e_.*x_)^m_.*(a_+b_.*ArcTan[c_.*x_^n_])^p_,x_Symbol] :=
  Int[ExpandIntegrand[(a+b*ArcTan[c*x^n])^p,(d+e*x)^m,x],x] /;
  FreeQ[{a,b,c,d,e,n},x] && IGtQ[p,1] && IGtQ[m,0]
```

```
Int[(d_+e_.*x_)^m_.*(a_+b_.*ArcCot[c_.*x_^n_])^p_,x_Symbol] :=
  Int[ExpandIntegrand[(a+b*ArcCot[c*x^n])^p,(d+e*x)^m,x],x] /;
  FreeQ[{a,b,c,d,e,n},x] && IGtQ[p,1] && IGtQ[m,0]
```

U: $\int (d+ex)^m (a+b \operatorname{ArcTan}[cx^n])^p dx$

Rule:

$$\int (d+ex)^m (a+b \operatorname{ArcTan}[cx^n])^p dx \rightarrow \int (d+ex)^m (a+b \operatorname{ArcTan}[cx^n])^p dx$$

Program code:

```
Int[(d_+e_.*x_)^m_.*(a_+b_.*ArcTan[c_.*x_^n_])^p_,x_Symbol] :=
  Unintegrable[(d+e*x)^m*(a+b*ArcTan[c*x^n])^p,x] /;
  FreeQ[{a,b,c,d,e,m,n,p},x]
```

```
Int[(d_+e_*x_)^m_*(a_+b_*ArcCot[c_*x_^n_])^p_,x_Symbol] :=  
  Unintegrable[(d+e*x)^m*(a+b*ArcCot[c*x^n])^p,x] /;  
FreeQ[{a,b,c,d,e,m,n,p},x]
```