

Rules for integrands of the form $u (a + b \operatorname{ArcTan}[c + d x])^p$

1. $\int u (a + b \operatorname{ArcTan}[c + d x])^p dx$

1: $\int (a + b \operatorname{ArcTan}[c + d x])^p dx$ when $p \in \mathbb{Z}^+$

Derivation: Integration by substitution

Rule: If $p \in \mathbb{Z}^+$, then

$$\int (a + b \operatorname{ArcTan}[c + d x])^p dx \rightarrow \frac{1}{d} \operatorname{Subst} \left[\int (a + b \operatorname{ArcTan}[x])^p dx, x, c + d x \right]$$

Program code:

```
Int[(a_+b_.*ArcTan[c_+d_.*x_])^p_,x_Symbol] :=
  1/d*Subst[Int[(a+b*ArcTan[x])^p,x],x,c+d*x] /;
FreeQ[{a,b,c,d},x] && IGtQ[p,0]
```

```
Int[(a_+b_.*ArcCot[c_+d_.*x_])^p_,x_Symbol] :=
  1/d*Subst[Int[(a+b*ArcCot[x])^p,x],x,c+d*x] /;
FreeQ[{a,b,c,d},x] && IGtQ[p,0]
```

U: $\int (a + b \operatorname{ArcTan}[c + d x])^p dx$ when $p \notin \mathbb{Z}^+$

– Rule: If $p \notin \mathbb{Z}^+$, then

$$\int (a + b \operatorname{ArcTan}[c + d x])^p dx \rightarrow \int (a + b \operatorname{ArcTan}[c + d x])^p dx$$

– Program code:

```
Int[(a_+b_.*ArcTan[c_+d_.*x_])^p_,x_Symbol] :=
  Unintegrable[(a+b*ArcTan[c+d*x])^p,x] /;
FreeQ[{a,b,c,d,p},x] && Not[IGtQ[p,0]]
```

```

Int[(a_.+b_.*ArcCot[c_+d_.*x_])^p_,x_Symbol] :=
  Unintegrable[(a+b*ArcCot[c+d*x])^p,x] /;
  FreeQ[{a,b,c,d,p},x] && Not[IGtQ[p,0]]

```

2. $\int (e + f x)^m (a + b \operatorname{ArcTan}[c + d x])^p dx$

1: $\int (e + f x)^m (a + b \operatorname{ArcTan}[c + d x])^p dx$ when $d e - c f = 0 \wedge p \in \mathbb{Z}^+$

Derivation: Integration by substitution

Rule: If $d e - c f = 0 \wedge p \in \mathbb{Z}^+$, then

$$\int (e + f x)^m (a + b \operatorname{ArcTan}[c + d x])^p dx \rightarrow \frac{1}{d} \operatorname{Subst}\left[\int \left(\frac{f x}{d}\right)^m (a + b \operatorname{ArcTan}[x])^p dx, x, c + d x\right]$$

Program code:

```

Int[(e_.+f_.*x_)^m_.*(a_.+b_.*ArcTan[c_+d_.*x_])^p_,x_Symbol] :=
  1/d*Subst[Int[(f*x/d)^m*(a+b*ArcTan[x])^p,x],x,c+d*x] /;
  FreeQ[{a,b,c,d,e,f,m},x] && EqQ[d*e-c*f,0] && IGtQ[p,0]

```

```

Int[(e_.+f_.*x_)^m_.*(a_.+b_.*ArcCot[c_+d_.*x_])^p_,x_Symbol] :=
  1/d*Subst[Int[(f*x/d)^m*(a+b*ArcCot[x])^p,x],x,c+d*x] /;
  FreeQ[{a,b,c,d,e,f,m},x] && EqQ[d*e-c*f,0] && IGtQ[p,0]

```

2: $\int (e + f x)^m (a + b \operatorname{ArcTan}[c + d x])^p dx$ when $p \in \mathbb{Z}^+ \wedge m + 1 \in \mathbb{Z}^-$

Derivation: Integration by parts

Basis: $\partial_x (a + b \operatorname{ArcTan}[c + d x])^p = \frac{b d p (a+b \operatorname{ArcTan}[c+d x])^{p-1}}{1+(c+d x)^2}$

Rule: If $p \in \mathbb{Z}^+ \wedge m + 1 \in \mathbb{Z}^-$, then

$$\int (e + f x)^m (a + b \operatorname{ArcTan}[c + d x])^p dx \rightarrow \frac{(e + f x)^{m+1} (a + b \operatorname{ArcTan}[c + d x])^p}{f (m+1)} - \frac{b d p}{f (m+1)} \int \frac{(e + f x)^{m+1} (a + b \operatorname{ArcTan}[c + d x])^{p-1}}{1 + (c + d x)^2} dx$$

Program code:

```
Int[ (e_..+f_..*x_)^m*(a_..+b_..*ArcTan[c_..+d_..*x_])^p.,x_Symbol] :=  

  (e+f*x)^(m+1)*(a+b*ArcTan[c+d*x])^p/(f*(m+1)) -  

  b*d*p/(f*(m+1))*Int[ (e+f*x)^(m+1)*(a+b*ArcTan[c+d*x])^(p-1)/(1+(c+d*x)^2),x] /;  

FreeQ[{a,b,c,d,e,f},x] && IGtQ[p,0] && ILtQ[m,-1]
```

```
Int[ (e_..+f_..*x_)^m*(a_..+b_..*ArcCot[c_..+d_..*x_])^p.,x_Symbol] :=  

  (e+f*x)^(m+1)*(a+b*ArcCot[c+d*x])^p/(f*(m+1)) +  

  b*d*p/(f*(m+1))*Int[ (e+f*x)^(m+1)*(a+b*ArcCot[c+d*x])^(p-1)/(1+(c+d*x)^2),x] /;  

FreeQ[{a,b,c,d,e,f},x] && IGtQ[p,0] && ILtQ[m,-1]
```

3: $\int (e + f x)^m (a + b \operatorname{ArcTan}[c + d x])^p dx$ when $p \in \mathbb{Z}^+$

Derivation: Integration by substitution

Rule: If $p \in \mathbb{Z}^+$, then

$$\int (e + f x)^m (a + b \operatorname{ArcTan}[c + d x])^p dx \rightarrow \frac{1}{d} \operatorname{Subst} \left[\int \left(\frac{d e - c f}{d} + \frac{f x}{d} \right)^m (a + b \operatorname{ArcTan}[x])^p dx, x, c + d x \right]$$

Program code:

```
Int[ (e_..+f_..*x_)^m*(a_..+b_..*ArcTan[c_..+d_..*x_])^p.,x_Symbol] :=  

  1/d*Subst[Int[((d*e-c*f)/d+f*x/d)^m*(a+b*ArcTan[x])^p,x],x,c+d*x] /;  

FreeQ[{a,b,c,d,e,f,m,p},x] && IGtQ[p,0]
```

```
Int[ (e_..+f_..*x_)^m*(a_..+b_..*ArcCot[c_..+d_..*x_])^p.,x_Symbol] :=  

  1/d*Subst[Int[((d*e-c*f)/d+f*x/d)^m*(a+b*ArcCot[x])^p,x],x,c+d*x] /;  

FreeQ[{a,b,c,d,e,f,m,p},x] && IGtQ[p,0]
```

U: $\int (e + f x)^m (a + b \operatorname{ArcTan}[c + d x])^p dx$ when $p \notin \mathbb{Z}^+$

Rule: If $p \notin \mathbb{Z}^+$, then

$$\int (e + f x)^m (a + b \operatorname{ArcTan}[c + d x])^p dx \rightarrow \int (e + f x)^m (a + b \operatorname{ArcTan}[c + d x])^p dx$$

Program code:

```
Int[(e_..+f_..*x_)^m_..*(a_..+b_..*ArcTan[c_..+d_..*x_])^p_,x_Symbol]:=  
  Unintegrable[(e+f*x)^m*(a+b*ArcTan[c+d*x])^p,x] /;  
  FreeQ[{a,b,c,d,e,f,m,p},x] && Not[IGtQ[p,0]]
```

```
Int[(e_..+f_..*x_)^m_..*(a_..+b_..*ArcCot[c_..+d_..*x_])^p_,x_Symbol]:=  
  Unintegrable[(e+f*x)^m*(a+b*ArcCot[c+d*x])^p,x] /;  
  FreeQ[{a,b,c,d,e,f,m,p},x] && Not[IGtQ[p,0]]
```

$$3. \int (e + f x^n)^m (a + b \operatorname{ArcTan}[c + d x])^p dx$$

$$1. \int \frac{\operatorname{ArcTan}[a + b x]}{c + d x^n} dx$$

$$1: \int \frac{\operatorname{ArcTan}[a + b x]}{c + d x^n} dx \text{ when } n \in \mathbb{Q}$$

Derivation: Algebraic expansion

$$\text{Basis: } \operatorname{ArcTan}[z] = \frac{1}{2} i \operatorname{Log}[1 - iz] - \frac{1}{2} i \operatorname{Log}[1 + iz]$$

$$\text{Basis: } \operatorname{ArcCot}[z] = \frac{i}{2} \operatorname{Log}\left[1 - \frac{i}{z}\right] - \frac{i}{2} \operatorname{Log}\left[1 + \frac{i}{z}\right]$$

Rule: If $n \in \mathbb{Q}$, then

$$\int \frac{\operatorname{ArcTan}[a + b x]}{c + d x^n} dx \rightarrow \frac{i}{2} \int \frac{\operatorname{Log}[1 - ia - ibx]}{c + d x^n} dx - \frac{i}{2} \int \frac{\operatorname{Log}[1 + ia + ibx]}{c + d x^n} dx$$

Program code:

```
Int[ArcTan[a_+b_.*x_]/(c_+d_.*x_^n_),x_Symbol] :=
  I/2*Int[Log[1-I*a-I*b*x]/(c+d*x^n),x] -
  I/2*Int[Log[1+I*a+I*b*x]/(c+d*x^n),x] /;
FreeQ[{a,b,c,d},x] && RationalQ[n]
```

```
Int[ArcCot[a_+b_.*x_]/(c_+d_.*x_^n_),x_Symbol] :=
  I/2*Int[Log[(-I+a+b*x)/(a+b*x)]/(c+d*x^n),x] -
  I/2*Int[Log[(I+a+b*x)/(a+b*x)]/(c+d*x^n),x] /;
FreeQ[{a,b,c,d},x] && RationalQ[n]
```

2: $\int \frac{\text{ArcTan}[a + b x]}{c + d x^n} dx$ when $n \notin \mathbb{Q}$

Rule: If $n \notin \mathbb{Q}$, then

$$\int \frac{\text{ArcTan}[a + b x]}{c + d x^n} dx \rightarrow \int \frac{\text{ArcTan}[a + b x]}{c + d x^n} dx$$

Program code:

```
Int[ArcTan[a_+b_.*x_]/(c_+d_.*x_^n_),x_Symbol] :=
  Unintegrable[ArcTan[a+b*x]/(c+d*x^n),x] /;
  FreeQ[{a,b,c,d,n},x] && Not[RationalQ[n]]
```

```
Int[ArcCot[a_+b_.*x_]/(c_+d_.*x_^n_),x_Symbol] :=
  Unintegrable[ArcCot[a+b*x]/(c+d*x^n),x] /;
  FreeQ[{a,b,c,d,n},x] && Not[RationalQ[n]]
```

4: $\int (A + B x + C x^2)^q (a + b \arctan(c + d x))^p dx$ when $B (1 + c^2) - 2 A c d = 0 \wedge 2 c C - B d = 0$

Derivation: Integration by substitution

Basis: If $B (1 + c^2) - 2 A c d = 0 \wedge 2 c C - B d = 0$, then $A + B x + C x^2 = \frac{C}{d^2} + \frac{C}{d^2} (c + d x)^2$

Rule: If $B (1 + c^2) - 2 A c d = 0 \wedge 2 c C - B d = 0$, then

$$\int (A + B x + C x^2)^q (a + b \arctan(c + d x))^p dx \rightarrow \frac{1}{d} \text{Subst}\left[\int \left(\frac{C}{d^2} + \frac{C x^2}{d^2}\right)^q (a + b \arctan(x))^p dx, x, c + d x\right]$$

Program code:

```
Int[(A_+B_.*x_+C_.*x_^2)^q_.*(a_+b_.*ArcTan[c_+d_.*x_])^p_,x_Symbol] :=
  1/d*Subst[Int[(C/d^2+C/d^2*x^2)^q*(a+b*ArcTan[x])^p,x],x,c+d*x] /;
  FreeQ[{a,b,c,d,A,B,C,p,q},x] && EqQ[B*(1+c^2)-2*A*c*d,0] && EqQ[2*c*C-B*d,0]
```

```

Int[(A_+B_.*x_+C_.*x_^2)^q_.*(a_+b_.*ArcCot[c_+d_.*x_])^p_,x_Symbol]:=  

  1/d*Subst[Int[(C/d^2+C/d^2*x^2)^q*(a+b*ArcCot[x])^p,x],x,c+d*x] /;  

FreeQ[{a,b,c,d,A,B,C,p,q},x] && EqQ[B*(1+c^2)-2*A*c*d,0] && EqQ[2*c*C-B*d,0]

```

5: $\int (e + f x)^m (A + B x + C x^2)^q (a + b \operatorname{ArcTan}[c + d x])^p dx \text{ when } B (1 + c^2) - 2 A c d == 0 \wedge 2 c C - B d == 0$

Derivation: Integration by substitution

Basis: If $B (1 + c^2) - 2 A c d == 0 \wedge 2 c C - B d == 0$, then $A + B x + C x^2 == \frac{C}{d^2} + \frac{C}{d^2} (c + d x)^2$

Rule: If $B (1 + c^2) - 2 A c d == 0 \wedge 2 c C - B d == 0$, then

$$\int (e + f x)^m (A + B x + C x^2)^q (a + b \operatorname{ArcTan}[c + d x])^p dx \rightarrow \frac{1}{d} \operatorname{Subst}\left[\int \left(\frac{d e - c f}{d} + \frac{f x}{d}\right)^m \left(\frac{C}{d^2} + \frac{C x^2}{d^2}\right)^q (a + b \operatorname{ArcTan}[x])^p dx, x, c + d x\right]$$

Program code:

```

Int[(e_+f_.*x_)^m_.*(A_+B_.*x_+C_.*x_^2)^q_.*(a_+b_.*ArcTan[c_+d_.*x_])^p_,x_Symbol]:=  

  1/d*Subst[Int[((d*e-c*f)/d+f*x/d)^m*(C/d^2+C/d^2*x^2)^q*(a+b*ArcTan[x])^p,x],x,c+d*x] /;  

FreeQ[{a,b,c,d,e,f,A,B,C,m,p,q},x] && EqQ[B*(1+c^2)-2*A*c*d,0] && EqQ[2*c*C-B*d,0]

```

```

Int[(e_+f_.*x_)^m_.*(A_+B_.*x_+C_.*x_^2)^q_.*(a_+b_.*ArcCot[c_+d_.*x_])^p_,x_Symbol]:=  

  1/d*Subst[Int[((d*e-c*f)/d+f*x/d)^m*(C/d^2+C/d^2*x^2)^q*(a+b*ArcCot[x])^p,x],x,c+d*x] /;  

FreeQ[{a,b,c,d,e,f,A,B,C,m,p,q},x] && EqQ[B*(1+c^2)-2*A*c*d,0] && EqQ[2*c*C-B*d,0]

```