

Rules for integrands involving inverse hyperbolic tangents and cotangents

1. $\int u \operatorname{ArcTanh}[a + b x^n] dx$

1: $\int \operatorname{ArcTanh}[a + b x^n] dx$

- Derivation: Integration by parts

- Rule:

$$\int \operatorname{ArcTanh}[a + b x^n] dx \rightarrow x \operatorname{ArcTanh}[a + b x^n] - b n \int \frac{x^n}{1 - a^2 - 2 a b x^n - b^2 x^{2n}} dx$$

- Program code:

```
Int[ArcTanh[a+b_*x_^n_],x_Symbol] :=  
  x*ArcTanh[a+b*x^n] -  
  b*n*Int[x^n/(1-a^2-2*a*b*x^n-b^2*x^(2*n)),x] /;  
FreeQ[{a,b,n},x]
```

```
Int[ArcCoth[a+b_*x_^n_],x_Symbol] :=  
  x*ArcCoth[a+b*x^n] -  
  b*n*Int[x^n/(1-a^2-2*a*b*x^n-b^2*x^(2*n)),x] /;  
FreeQ[{a,b,n},x]
```

$$2. \int x^m \operatorname{ArcTanh}[a + b x^n] dx$$

$$1: \int \frac{\operatorname{ArcTanh}[a + b x^n]}{x} dx$$

Derivation: Algebraic expansion

Basis: $\operatorname{ArcTanh}[z] = \frac{1}{2} \operatorname{Log}[1+z] - \frac{1}{2} \operatorname{Log}[1-z]$

Basis: $\operatorname{ArcCoth}[z] = \frac{1}{2} \operatorname{Log}\left[1 + \frac{1}{z}\right] - \frac{1}{2} \operatorname{Log}\left[1 - \frac{1}{z}\right]$

Rule:

$$\int \frac{\operatorname{ArcTanh}[a + b x^n]}{x} dx \rightarrow \frac{1}{2} \int \frac{\operatorname{Log}[1 + a + b x^n]}{x} dx - \frac{1}{2} \int \frac{\operatorname{Log}[1 - a - b x^n]}{x} dx$$

Program code:

```
Int[ArcTanh[a_+b_*x_^n_]/x_,x_Symbol] :=
  1/2*Int[Log[1+a+b*x^n]/x,x] -
  1/2*Int[Log[1-a-b*x^n]/x,x] /;
FreeQ[{a,b,n},x]
```

```
Int[ArcCoth[a_+b_*x_^n_]/x_,x_Symbol] :=
  1/2*Int[Log[1+1/(a+b*x^n)]/x,x] -
  1/2*Int[Log[1-1/(a+b*x^n)]/x,x] /;
FreeQ[{a,b,n},x]
```

$$2: \int x^m \operatorname{ArcTanh}[a + b x^n] dx \text{ when } (m | n) \in \mathbb{Q} \wedge m + 1 \neq 0 \wedge m + 1 \neq n$$

Reference: CRC 588, A&S 4.6.54

Reference: CRC 590, A&S 4.6.60

Derivation: Integration by parts

Rule: If $(m | n) \in \mathbb{Q} \wedge m + 1 \neq 0 \wedge m + 1 \neq n$, then

$$\int x^m \operatorname{ArcTanh}[a + b x^n] dx \rightarrow \frac{x^{m+1} \operatorname{ArcTanh}[a + b x^n]}{m + 1} - \frac{b n}{m + 1} \int \frac{x^{m+n}}{1 - a^2 - 2 a b x^n - b^2 x^{2n}} dx$$

Program code:

```
Int[x^m_*ArcTanh[a+b_*x^n],x_Symbol] :=
  x^(m+1)*ArcTanh[a+b*x^n]/(m+1) -
  b*n/(m+1)*Int[x^(m+n)/(1-a^2-2*a*b*x^n-b^2*x^(2*n)),x] /;
FreeQ[{a,b},x] && RationalQ[m,n] && NeQ[m,-1] && NeQ[m+1,n]
```

```
Int[x^m_*ArcCoth[a+b_*x^n],x_Symbol] :=
  x^(m+1)*ArcCoth[a+b*x^n]/(m+1) -
  b*n/(m+1)*Int[x^(m+n)/(1-a^2-2*a*b*x^n-b^2*x^(2*n)),x] /;
FreeQ[{a,b},x] && RationalQ[m,n] && NeQ[m,-1] && NeQ[m+1,n]
```

$$2. \int u \operatorname{ArcTanh}[a + b f^{c+dx}] dx$$

$$1: \int \operatorname{ArcTanh}[a + b f^{c+dx}] dx$$

Derivation: Algebraic expansion

$$\text{Basis: } \operatorname{ArcTanh}[z] = \frac{1}{2} \operatorname{Log}[1+z] - \frac{1}{2} \operatorname{Log}[1-z]$$

$$\text{Basis: } \operatorname{ArcCoth}[z] = \frac{1}{2} \operatorname{Log}\left[1 + \frac{1}{z}\right] - \frac{1}{2} \operatorname{Log}\left[1 - \frac{1}{z}\right]$$

Rule:

$$\int \operatorname{ArcTanh}[a + b f^{c+dx}] dx \rightarrow \frac{1}{2} \int \operatorname{Log}[1 + a + b f^{c+dx}] dx - \frac{1}{2} \int \operatorname{Log}[1 - a - b f^{c+dx}] dx$$

Program code:

```
Int[ArcTanh[a_+b_*f^(c_+d_*x_)],x_Symbol] :=
  1/2*Int[Log[1+a+b*f^(c+d*x)],x] -
  1/2*Int[Log[1-a-b*f^(c+d*x)],x] /;
FreeQ[{a,b,c,d,f},x]
```

```
Int[ArcCoth[a_+b_*f^(c_+d_*x_)],x_Symbol] :=
  1/2*Int[Log[1+1/(a+b*f^(c+d*x))],x] -
  1/2*Int[Log[1-1/(a+b*f^(c+d*x))],x] /;
FreeQ[{a,b,c,d,f},x]
```

$$2: \int x^m \operatorname{ArcTanh}[a + b f^{c+dx}] dx \text{ when } m \in \mathbb{Z} \wedge m > 0$$

Derivation: Algebraic expansion

$$\text{Basis: } \operatorname{ArcTanh}[z] = \frac{1}{2} \operatorname{Log}[1+z] - \frac{1}{2} \operatorname{Log}[1-z]$$

$$\text{Basis: } \operatorname{ArcCoth}[z] = \frac{1}{2} \operatorname{Log}\left[1 + \frac{1}{z}\right] - \frac{1}{2} \operatorname{Log}\left[1 - \frac{1}{z}\right]$$

Rule: If $m \in \mathbb{Z} \wedge m > 0$, **then**

$$\int x^m \operatorname{ArcTanh}[a + b f^{c+dx}] dx \rightarrow \frac{1}{2} \int x^m \operatorname{Log}[1 + a + b f^{c+dx}] dx - \frac{1}{2} \int x^m \operatorname{Log}[1 - a - b f^{c+dx}] dx$$

Program code:

```
Int[x_^m_*ArcTanh[a_+b_*f^(c_+d_*x_)],x_Symbol] :=
  1/2*Int[x^m*Log[1+a+b*f^(c+d*x)],x] -
  1/2*Int[x^m*Log[1-a-b*f^(c+d*x)],x] /;
FreeQ[{a,b,c,d,f},x] && IGtQ[m,0]
```

```
Int[x_^m_*ArcCoth[a_+b_*f^(c_+d_*x_)],x_Symbol] :=
  1/2*Int[x^m*Log[1+1/(a+b*f^(c+d*x))],x] -
  1/2*Int[x^m*Log[1-1/(a+b*f^(c+d*x))],x] /;
FreeQ[{a,b,c,d,f},x] && IGtQ[m,0]
```

$$3: \int u \operatorname{ArcTanh}\left[\frac{c}{a + b x^n}\right]^m dx$$

Derivation: Algebraic simplification

$$\text{Basis: } \operatorname{ArcTanh}\left[\frac{1}{z}\right] = \operatorname{ArcCoth}\left[\frac{1}{z}\right]$$

Rule:

$$\int u \operatorname{ArcTanh}\left[\frac{c}{a + b x^n}\right]^m dx \rightarrow \int u \operatorname{ArcCoth}\left[\frac{a}{c} + \frac{b x^n}{c}\right]^m dx$$

Program code:

```
Int[u_*ArcTanh[c_/(a_+b_*x^n_)^m_],x_Symbol] :=
  Int[u*ArcCoth[a/c+b*x^n/c]^m,x] /;
FreeQ[{a,b,c,n,m},x]
```

```
Int[u.*ArcCoth[c./ (a.+b.*x^n.)]^m.,x_Symbol] :=
  Int[u*ArcTanh[a/c+b*x^n/c]^m,x] /;
FreeQ[{a,b,c,n,m},x]
```

4. $\int u \operatorname{ArcTanh}\left[\frac{c x}{\sqrt{a+b x^2}}\right] dx$ when $b = c^2$

1: $\int \operatorname{ArcTanh}\left[\frac{c x}{\sqrt{a+b x^2}}\right] dx$ when $b = c^2$

Derivation: Integration by parts

■ **Basis:** If $b = c^2$, then $\partial_x \operatorname{ArcTanh}\left[\frac{c x}{\sqrt{a+b x^2}}\right] = \frac{c}{\sqrt{a+b x^2}}$

– **Rule:** If $b = c^2$, then

$$\int \operatorname{ArcTanh}\left[\frac{c x}{\sqrt{a+b x^2}}\right] dx \rightarrow x \operatorname{ArcTanh}\left[\frac{c x}{\sqrt{a+b x^2}}\right] - c \int \frac{x}{\sqrt{a+b x^2}} dx$$

– **Program code:**

```
Int[ArcTanh[c.*x_/Sqrt[a.+b.*x^2]],x_Symbol] :=
  x*ArcTanh[(c*x)/Sqrt[a+b*x^2]] - c*Int[x/Sqrt[a+b*x^2],x] /;
FreeQ[{a,b,c},x] && EqQ[b,c^2]
```

```
Int[ArcCoth[c.*x_/Sqrt[a.+b.*x^2]],x_Symbol] :=
  x*ArcCoth[(c*x)/Sqrt[a+b*x^2]] - c*Int[x/Sqrt[a+b*x^2],x] /;
FreeQ[{a,b,c},x] && EqQ[b,c^2]
```

2. $\int (d x)^m \operatorname{ArcTanh}\left[\frac{c x}{\sqrt{a+b x^2}}\right] dx$ when $b = c^2$

1: $\int \frac{\operatorname{ArcTanh}\left[\frac{c x}{\sqrt{a+b x^2}}\right]}{x} dx$ when $b = c^2$

– **Derivation: Integration by parts**

■ **Basis:** If $b = c^2$, then $\partial_x \operatorname{ArcTanh}\left[\frac{c x}{\sqrt{a+b x^2}}\right] = \frac{c}{\sqrt{a+b x^2}}$

– **Rule:** If $b = c^2$, then

$$\int \frac{\text{ArcTanh}\left[\frac{c x}{\sqrt{a+b x^2}}\right]}{x} dx \rightarrow \text{ArcTanh}\left[\frac{c x}{\sqrt{a+b x^2}}\right] \text{Log}[x] - c \int \frac{\text{Log}[x]}{\sqrt{a+b x^2}} dx$$

Program code:

```
Int[ArcTanh[c.*x_/Sqrt[a_.+b_.*x_^2]]/x_,x_Symbol] :=
  ArcTanh[c*x/Sqrt[a+b*x^2]]*Log[x] - c*Int[Log[x]/Sqrt[a+b*x^2],x] /;
FreeQ[{a,b,c},x] && EqQ[b,c^2]
```

```
Int[ArcCoth[c.*x_/Sqrt[a_.+b_.*x_^2]]/x_,x_Symbol] :=
  ArcCoth[c*x/Sqrt[a+b*x^2]]*Log[x] - c*Int[Log[x]/Sqrt[a+b*x^2],x] /;
FreeQ[{a,b,c},x] && EqQ[b,c^2]
```

2: $\int (dx)^m \text{ArcTanh}\left[\frac{c x}{\sqrt{a+b x^2}}\right] dx$ when $b = c^2 \wedge m \neq -1$

Derivation: Integration by parts

■ Basis: If $b = c^2$, then $\partial_x \text{ArcTanh}\left[\frac{c x}{\sqrt{a+b x^2}}\right] = \frac{c}{\sqrt{a+b x^2}}$

Rule: If $b = c^2 \wedge m \neq -1$, then

$$\int (dx)^m \text{ArcTanh}\left[\frac{c x}{\sqrt{a+b x^2}}\right] dx \rightarrow \frac{(dx)^{m+1} \text{ArcTanh}\left[\frac{c x}{\sqrt{a+b x^2}}\right]}{d(m+1)} - \frac{c}{d(m+1)} \int \frac{(dx)^{m+1}}{\sqrt{a+b x^2}} dx$$

Program code:

```
Int[(d_.*x_)^m_.*ArcTanh[c.*x_/Sqrt[a_.+b_.*x_^2]],x_Symbol] :=
  (d*x)^(m+1)*ArcTanh[(c*x)/Sqrt[a+b*x^2]]/(d*(m+1)) - c/(d*(m+1))*Int[(d*x)^(m+1)/Sqrt[a+b*x^2],x] /;
FreeQ[{a,b,c,d,m},x] && EqQ[b,c^2] && NeQ[m,-1]
```

```
Int[(d_.*x_)^m_.*ArcCoth[c.*x_/Sqrt[a_.+b_.*x_^2]],x_Symbol] :=
  (d*x)^(m+1)*ArcCoth[(c*x)/Sqrt[a+b*x^2]]/(d*(m+1)) - c/(d*(m+1))*Int[(d*x)^(m+1)/Sqrt[a+b*x^2],x] /;
FreeQ[{a,b,c,d,m},x] && EqQ[b,c^2] && NeQ[m,-1]
```

3. $\int \frac{\text{ArcTanh}\left[\frac{c x}{\sqrt{a+b x^2}}\right]^m}{\sqrt{d+e x^2}} dx$ when $b = c^2 \wedge b d - a e = 0$

$$1. \int \frac{\text{ArcTanh}\left[\frac{cx}{\sqrt{a+bx^2}}\right]^m}{\sqrt{a+bx^2}} dx \text{ when } b = c^2$$

$$1: \int \frac{1}{\sqrt{a+bx^2} \text{ArcTanh}\left[\frac{cx}{\sqrt{a+bx^2}}\right]} dx \text{ when } b = c^2$$

Derivation: Reciprocal rule for integration

■ Basis: If $b = c^2$, then $\partial_x \text{ArcTanh}\left[\frac{cx}{\sqrt{a+bx^2}}\right] = \frac{c}{\sqrt{a+bx^2}}$

Rule: If $b = c^2$, then

$$\int \frac{1}{\sqrt{a+bx^2} \text{ArcTanh}\left[\frac{cx}{\sqrt{a+bx^2}}\right]} dx \rightarrow \frac{1}{c} \text{Log}\left[\text{ArcTanh}\left[\frac{cx}{\sqrt{a+bx^2}}\right]\right]$$

Program code:

```
Int[1/(Sqrt[a_+b_.*x_^2]*ArcTanh[c_.*x_/Sqrt[a_+b_.*x_^2]]),x_Symbol] :=
  1/c*Log[ArcTanh[c*x/Sqrt[a+b*x^2]]] /;
FreeQ[{a,b,c},x] && EqQ[b,c^2]
```

```
Int[1/(Sqrt[a_+b_.*x_^2]*ArcCoth[c_.*x_/Sqrt[a_+b_.*x_^2]]),x_Symbol] :=
  -1/c*Log[ArcCoth[c*x/Sqrt[a+b*x^2]]] /;
FreeQ[{a,b,c},x] && EqQ[b,c^2]
```

$$2: \int \frac{\text{ArcTanh}\left[\frac{cx}{\sqrt{a+bx^2}}\right]^m}{\sqrt{a+bx^2}} dx \text{ when } b = c^2 \wedge m \neq -1$$

Derivation: Power rule for integration

$$\text{Basis: If } b = c^2, \text{ then } \partial_x \text{ArcTanh}\left[\frac{cx}{\sqrt{a+bx^2}}\right] = \frac{c}{\sqrt{a+bx^2}}$$

Rule: If $b = c^2 \wedge m \neq -1$, then

$$\int \frac{\text{ArcTanh}\left[\frac{cx}{\sqrt{a+bx^2}}\right]^m}{\sqrt{a+bx^2}} dx \rightarrow \frac{\text{ArcTanh}\left[\frac{cx}{\sqrt{a+bx^2}}\right]^{m+1}}{c(m+1)}$$

Program code:

```
Int[ArcTanh[c.*x_/Sqrt[a_.+b_.*x_^2]]^m_/Sqrt[a_.+b_.*x_^2],x_Symbol] :=
  ArcTanh[c*x/Sqrt[a+b*x^2]]^(m+1)/(c*(m+1)) /;
FreeQ[{a,b,c,m},x] && EqQ[b,c^2] && NeQ[m,-1]
```

```
Int[ArcCoth[c.*x_/Sqrt[a_.+b_.*x_^2]]^m_/Sqrt[a_.+b_.*x_^2],x_Symbol] :=
  -ArcCoth[c*x/Sqrt[a+b*x^2]]^(m+1)/(c*(m+1)) /;
FreeQ[{a,b,c,m},x] && EqQ[b,c^2] && NeQ[m,-1]
```

$$2: \int \frac{\text{ArcTanh}\left[\frac{cx}{\sqrt{a+bx^2}}\right]^m}{\sqrt{d+ex^2}} dx \text{ when } b = c^2 \wedge bd - ae = 0$$

Derivation: Piecewise constant extraction

■ Basis: If $bd - ae = 0$, then $\partial_x \frac{\sqrt{a+bx^2}}{\sqrt{d+ex^2}} = 0$

Rule: If $b = c^2 \wedge bd - ae = 0$, then

$$\int \frac{\text{ArcTanh}\left[\frac{cx}{\sqrt{a+bx^2}}\right]^m}{\sqrt{d+ex^2}} dx \rightarrow \frac{\sqrt{a+bx^2}}{\sqrt{d+ex^2}} \int \frac{\text{ArcTanh}\left[\frac{cx}{\sqrt{a+bx^2}}\right]^m}{\sqrt{a+bx^2}} dx$$

Program code:

```
Int[ArcTanh[c.*x_/Sqrt[a_.+b_.*x_^2]]^m./Sqrt[d_.+e_.*x_^2],x_Symbol] :=
  Sqrt[a+b*x^2]/Sqrt[d+e*x^2]*Int[ArcTanh[c*x/Sqrt[a+b*x^2]]^m/Sqrt[a+b*x^2],x] /;
FreeQ[{a,b,c,d,e,m},x] && EqQ[b,c^2] && EqQ[b*d-a*e,0]
```

```
Int[ArcCoth[c.*x_/Sqrt[a_.+b_.*x_^2]]^m./Sqrt[d_.+e_.*x_^2],x_Symbol] :=
  Sqrt[a+b*x^2]/Sqrt[d+e*x^2]*Int[ArcCoth[c*x/Sqrt[a+b*x^2]]^m/Sqrt[a+b*x^2],x] /;
FreeQ[{a,b,c,d,e,m},x] && EqQ[b,c^2] && EqQ[b*d-a*e,0]
```

$$5: \int \frac{f[x, \text{ArcTanh}[a + b x]]}{1 - (a + b x)^2} dx$$

Derivation: Integration by substitution

- **Basis:** $\frac{f[z]}{1-z^2} == f[\text{Tanh}[\text{ArcTanh}[z]]] \text{ArcTanh}'[z]$
- **Basis:** $r + s x + t x^2 == -\frac{s^2-4 r t}{4 t} \left(1 - \frac{(s+2 t x)^2}{s^2-4 r t}\right)$
- **Basis:** $1 - \text{Tanh}[z]^2 == \text{Sech}[z]^2$

Rule:

$$\int \frac{f[x, \text{ArcTanh}[a + b x]]}{1 - (a + b x)^2} dx \rightarrow \frac{1}{b} \text{Subst}\left[\int f\left[-\frac{a}{b} + \frac{\text{Tanh}[x]}{b}, x\right] dx, x, \text{ArcTanh}[a + b x]\right]$$

Program code:

```
If[TrueQ[$LoadShowSteps],

Int[u_*v_^n_.,x_Symbol] :=
  With[{tmp=InverseFunctionOfLinear[u,x]},
  ShowStep["", "Int[f[x,ArcTanh[a+b*x]]/(1-(a+b*x)^2),x]",
    "Subst[Int[f[-a/b+Tanh[x]/b,x],x,ArcTanh[a+b*x]]/b",Hold[
  (-Discriminant[v,x]/(4*Coefficient[v,x,2]))^n/Coefficient[tmp[[1]],x,1]*
  Subst[Int[SimplifyIntegrand[SubstForInverseFunction[u,tmp,x]*Sech[x]^(2*(n+1)),x],x], x, tmp]] /;
  Not[FalseQ[tmp]] && EqQ[Head[tmp],ArcTanh] && EqQ[Discriminant[v,x]*tmp[[1]]^2-D[v,x]^2,0] /;
  SimplifyFlag && QuadraticQ[v,x] && ILtQ[n,0] && PosQ[Discriminant[v,x]] && MatchQ[u,r_.*f_^w_ /; FreeQ[f,x]],

Int[u_*v_^n_.,x_Symbol] :=
  With[{tmp=InverseFunctionOfLinear[u,x]},
  (-Discriminant[v,x]/(4*Coefficient[v,x,2]))^n/Coefficient[tmp[[1]],x,1]*
  Subst[Int[SimplifyIntegrand[SubstForInverseFunction[u,tmp,x]*Sech[x]^(2*(n+1)),x],x], x, tmp] /;
  Not[FalseQ[tmp]] && EqQ[Head[tmp],ArcTanh] && EqQ[Discriminant[v,x]*tmp[[1]]^2-D[v,x]^2,0] /;
  QuadraticQ[v,x] && ILtQ[n,0] && PosQ[Discriminant[v,x]] && MatchQ[u,r_.*f_^w_ /; FreeQ[f,x]]]
```

```

If[TrueQ[$LoadShowSteps],

Int[u_*v_^n_.,x_Symbol] :=
  With[{tmp=InverseFunctionOfLinear[u,x]},
  ShowStep["", "Int[f[x,ArcCoth[a+b*x]]/(1-(a+b*x)^2),x]",
    "Subst[Int[f[-a/b+Coth[x]/b,x],x,ArcCoth[a+b*x]]/b",Hold[
  (-Discriminant[v,x]/(4*Coefficient[v,x,2]))^n/Coefficient[tmp[[1]],x,1]*
  Subst[Int[SimplifyIntegrand[SubstForInverseFunction[u,tmp,x]*(-Csch[x]^2)^(n+1),x],x],x,tmp]] /;
  Not[FalseQ[tmp]] && EqQ[Head[tmp],ArcCoth] && EqQ[Discriminant[v,x]*tmp[[1]]^2-D[v,x]^2,0] /;
  SimplifyFlag && QuadraticQ[v,x] && ILtQ[n,0] && PosQ[Discriminant[v,x]] && MatchQ[u,r_.*f_^w_ /; FreeQ[f,x]],

Int[u_*v_^n_.,x_Symbol] :=
  With[{tmp=InverseFunctionOfLinear[u,x]},
  (-Discriminant[v,x]/(4*Coefficient[v,x,2]))^n/Coefficient[tmp[[1]],x,1]*
  Subst[Int[SimplifyIntegrand[SubstForInverseFunction[u,tmp,x]*(-Csch[x]^2)^(n+1),x],x],x,tmp] /;
  Not[FalseQ[tmp]] && EqQ[Head[tmp],ArcCoth] && EqQ[Discriminant[v,x]*tmp[[1]]^2-D[v,x]^2,0] /;
  QuadraticQ[v,x] && ILtQ[n,0] && PosQ[Discriminant[v,x]] && MatchQ[u,r_.*f_^w_ /; FreeQ[f,x]]]

```

$$6. \int u \operatorname{ArcTanh}[c + d \operatorname{Tanh}[a + b x]] dx$$

$$1. \int \operatorname{ArcTanh}[c + d \operatorname{Tanh}[a + b x]] dx$$

$$1: \int \operatorname{ArcTanh}[c + d \operatorname{Tanh}[a + b x]] dx \text{ when } (c - d)^2 = 1$$

- Derivation: Integration by parts

- Basis: If $(c - d)^2 = 1$, then $\partial_x \operatorname{ArcTanh}[c + d \operatorname{Tanh}[a + b x]] = -\frac{b}{c - d + c e^{2a + 2bx}}$

- Rule: If $(c - d)^2 = 1$, then

$$\int \operatorname{ArcTanh}[c + d \operatorname{Tanh}[a + b x]] dx \rightarrow x \operatorname{ArcTanh}[c + d \operatorname{Tanh}[a + b x]] + b \int \frac{x}{c - d + c e^{2a + 2bx}} dx$$

Program code:

```

Int[ArcTanh[c_.*d_.*Tanh[a_.*b_.*x_]],x_Symbol] :=
  x*ArcTanh[c+d*Tanh[a+b*x]] +
  b*Int[x/(c-d+c*E^(2*a+2*b*x)),x] /;
FreeQ[{a,b,c,d},x] && EqQ[(c-d)^2,1]

```

```

Int[ArcCoth[c_.+d_.*Tanh[a_.+b_.*x_]],x_Symbol] :=
  x*ArcCoth[c+d*Tanh[a+b*x]] +
  b*Int[x/(c-d+c*E^(2*a+2*b*x)),x] /;
FreeQ[{a,b,c,d},x] && EqQ[(c-d)^2,1]

```

```

Int[ArcTanh[c_.+d_.*Coth[a_.+b_.*x_]],x_Symbol] :=
  x*ArcTanh[c+d*Coth[a+b*x]] +
  b*Int[x/(c-d-c*E^(2*a+2*b*x)),x] /;
FreeQ[{a,b,c,d},x] && EqQ[(c-d)^2,1]

```

```

Int[ArcCoth[c_.+d_.*Coth[a_.+b_.*x_]],x_Symbol] :=
  x*ArcCoth[c+d*Coth[a+b*x]] +
  b*Int[x/(c-d-c*E^(2*a+2*b*x)),x] /;
FreeQ[{a,b,c,d},x] && EqQ[(c-d)^2,1]

```

$$2: \int \text{ArcTanh}[c + d \text{Tanh}[a + b x]] dx \text{ when } (c - d)^2 \neq 1$$

Derivation: Integration by parts

$$\text{Basis: } \partial_x \text{ArcTanh}[c + d \text{Tanh}[a + b x]] = -\frac{b(1-c-d)e^{2a+2bx}}{1-c+d+(1-c-d)e^{2(a+bx)}} + \frac{b(1+c+d)e^{2a+2bx}}{1+c-d+(1+c+d)e^{2a+2bx}}$$

Rule: If $(c - d)^2 \neq 1$, **then**

$$\int \text{ArcTanh}[c + d \text{Tanh}[a + b x]] dx \rightarrow x \text{ArcTanh}[c + d \text{Tanh}[a + b x]] + b(1-c-d) \int \frac{x e^{2a+2bx}}{1-c+d+(1-c-d)e^{2a+2bx}} dx - b(1+c+d) \int \frac{x e^{2a+2bx}}{1+c-d+(1+c+d)e^{2a+2bx}} dx$$

$$\text{Basis: } \partial_x \text{ArcTanh}[c + d \text{Tanh}[a + b x]] = -\frac{b(1+c-d)}{1+c-d+(1+c+d)e^{2a+2bx}} + \frac{b(1-c+d)}{1-c+d+(1-c-d)e^{2a+2bx}}$$

Note: Although this formula appears simpler, it either introduces superfluous terms that have to be cancelled out, or results in a slightly more complicated antiderivative.

Rule: If $(c - d)^2 \neq 1$, **then**

$$\int \text{ArcTanh}[c + d \text{Tanh}[a + b x]] dx \rightarrow x \text{ArcTanh}[c + d \text{Tanh}[a + b x]] + b(1+c-d) \int \frac{x}{1+c-d+(1+c+d)e^{2a+2bx}} dx - b(1-c+d) \int \frac{x}{1-c+d+(1-c-d)e^{2a+2bx}} dx$$

Program code:

```
Int[ArcTanh[c_+d_*Tanh[a_+b_*x_]],x_Symbol] :=
  x*ArcTanh[c+d*Tanh[a+b*x]] +
  b*(1-c-d)*Int[x*E^(2*a+2*b*x)/(1-c+d+(1-c-d)*E^(2*a+2*b*x)),x] -
  b*(1+c+d)*Int[x*E^(2*a+2*b*x)/(1+c-d+(1+c+d)*E^(2*a+2*b*x)),x] /;
FreeQ[{a,b,c,d},x] && NeQ[(c-d)^2,1]
```

```
Int[ArcCoth[c_+d_*Tanh[a_+b_*x_]],x_Symbol] :=
  x*ArcCoth[c+d*Tanh[a+b*x]] +
  b*(1-c-d)*Int[x*E^(2*a+2*b*x)/(1-c+d+(1-c-d)*E^(2*a+2*b*x)),x] -
  b*(1+c+d)*Int[x*E^(2*a+2*b*x)/(1+c-d+(1+c+d)*E^(2*a+2*b*x)),x] /;
FreeQ[{a,b,c,d},x] && NeQ[(c-d)^2,1]
```

```
Int[ArcTanh[c_.+d_.*Coth[a_.+b_.*x_]],x_Symbol] :=
  x*ArcTanh[c+d*Coth[a+b*x]] +
  b*(1+c+d)*Int[x*E^(2*a+2*b*x)/(1+c-d-(1+c+d)*E^(2*a+2*b*x)),x] -
  b*(1-c-d)*Int[x*E^(2*a+2*b*x)/(1-c+d-(1-c-d)*E^(2*a+2*b*x)),x] /;
FreeQ[{a,b,c,d},x] && NeQ[(c-d)^2,1]
```

```
Int[ArcCoth[c_.+d_.*Coth[a_.+b_.*x_]],x_Symbol] :=
  x*ArcCoth[c+d*Coth[a+b*x]] +
  b*(1+c+d)*Int[x*E^(2*a+2*b*x)/(1+c-d-(1+c+d)*E^(2*a+2*b*x)),x] -
  b*(1-c-d)*Int[x*E^(2*a+2*b*x)/(1-c+d-(1-c-d)*E^(2*a+2*b*x)),x] /;
FreeQ[{a,b,c,d},x] && NeQ[(c-d)^2,1]
```

2. $\int (e + f x)^m \text{ArcTanh}[c + d \text{Tanh}[a + b x]] dx$ when $m \in \mathbb{Z}^+$

1: $\int (e + f x)^m \text{ArcTanh}[c + d \text{Tanh}[a + b x]] dx$ when $m \in \mathbb{Z}^+ \wedge (c - d)^2 = 1$

Derivation: Integration by parts

Basis: If $(c - d)^2 = 1$, then $\partial_x \text{ArcTanh}[c + d \text{Tanh}[a + b x]] = -\frac{b}{c-d+c e^{2a+2bx}}$

Rule: If $m \in \mathbb{Z}^+ \wedge (c - d)^2 = 1$, then

$$\int (e + f x)^m \text{ArcTanh}[c + d \text{Tanh}[a + b x]] dx \rightarrow \frac{(e + f x)^{m+1} \text{ArcTanh}[c + d \text{Tanh}[a + b x]]}{f(m+1)} + \frac{b}{f(m+1)} \int \frac{(e + f x)^{m+1}}{c - d + c e^{2a+2bx}} dx$$

Program code:

```
Int[(e_.+f_.*x_)^m_.*ArcTanh[c_.+d_.*Tanh[a_.+b_.*x_]],x_Symbol] :=
  (e+f*x)^(m+1)*ArcTanh[c+d*Tanh[a+b*x]]/(f*(m+1)) +
  b/(f*(m+1))*Int[(e+f*x)^(m+1)/(c-d+c*E^(2*a+2*b*x)),x] /;
FreeQ[{a,b,c,d,e,f},x] && IGtQ[m,0] && EqQ[(c-d)^2,1]
```

```
Int[(e_.+f_.*x_)^m_.*ArcCoth[c_.+d_.*Tanh[a_.+b_.*x_]],x_Symbol] :=
  (e+f*x)^(m+1)*ArcCoth[c+d*Tanh[a+b*x]]/(f*(m+1)) +
  b/(f*(m+1))*Int[(e+f*x)^(m+1)/(c-d+c*E^(2*a+2*b*x)),x] /;
FreeQ[{a,b,c,d,e,f},x] && IGtQ[m,0] && EqQ[(c-d)^2,1]
```

```
Int[(e_.+f_.**x_)^m_.*ArcTanh[c_.+d_.*Coth[a_.+b_.**x_]],x_Symbol] :=
  (e+f*x)^(m+1)*ArcTanh[c+d*Coth[a+b*x]]/(f*(m+1)) +
  b/(f*(m+1))*Int[(e+f*x)^(m+1)/(c-d-c*E^(2*a+2*b*x)),x] /;
FreeQ[{a,b,c,d,e,f},x] && IGtQ[m,0] && EqQ[(c-d)^2,1]
```

```
Int[(e_.+f_.**x_)^m_.*ArcCoth[c_.+d_.*Coth[a_.+b_.**x_]],x_Symbol] :=
  (e+f*x)^(m+1)*ArcCoth[c+d*Coth[a+b*x]]/(f*(m+1)) +
  b/(f*(m+1))*Int[(e+f*x)^(m+1)/(c-d-c*E^(2*a+2*b*x)),x] /;
FreeQ[{a,b,c,d,e,f},x] && IGtQ[m,0] && EqQ[(c-d)^2,1]
```

$$2: \int (e + f x)^m \operatorname{ArcTanh}[c + d \operatorname{Tanh}[a + b x]] dx \text{ when } m \in \mathbb{Z}^+ \wedge (c - d)^2 \neq 1$$

Derivation: Integration by parts

■ **Basis:** $\partial_x \operatorname{ArcTanh}[c + d \operatorname{Tanh}[a + b x]] = -\frac{b(1-c-d)e^{2a+2bx}}{1-c+d+(1-c-d)e^{2(a+bx)}} + \frac{b(1+c+d)e^{2a+2bx}}{1+c-d+(1+c+d)e^{2a+2bx}}$

– **Rule:** If $m \in \mathbb{Z}^+ \wedge (c - d)^2 \neq 1$, then

$$\int (e + f x)^m \operatorname{ArcTanh}[c + d \operatorname{Tanh}[a + b x]] dx \rightarrow \frac{(e + f x)^{m+1} \operatorname{ArcTanh}[c + d \operatorname{Tanh}[a + b x]]}{f(m+1)} + \frac{b(1-c-d)}{f(m+1)} \int \frac{(e + f x)^{m+1} e^{2a+2bx}}{1-c+d+(1-c-d)e^{2a+2bx}} dx - \frac{b(1+c+d)}{f(m+1)} \int \frac{(e + f x)^{m+1} e^{2a+2bx}}{1+c-d+(1+c+d)e^{2a+2bx}} dx$$

Program code:

```
Int[(e_.+f_.**x_)^m_.*ArcTanh[c_.+d_.*Tanh[a_.+b_.**x_]],x_Symbol] :=
  (e+f*x)^(m+1)*ArcTanh[c+d*Tanh[a+b*x]]/(f*(m+1)) +
  b*(1-c-d)/(f*(m+1))*Int[(e+f*x)^(m+1)*E^(2*a+2*b*x)/(1-c+d+(1-c-d)*E^(2*a+2*b*x)),x] -
  b*(1+c+d)/(f*(m+1))*Int[(e+f*x)^(m+1)*E^(2*a+2*b*x)/(1+c-d+(1+c+d)*E^(2*a+2*b*x)),x] /;
FreeQ[{a,b,c,d,e,f},x] && IGtQ[m,0] && NeQ[(c-d)^2,1]
```

```
Int[(e_.+f_.**x_)^m_.*ArcCoth[c_.+d_.*Tanh[a_.+b_.**x_]],x_Symbol] :=
  (e+f*x)^(m+1)*ArcCoth[c+d*Tanh[a+b*x]]/(f*(m+1)) +
  b*(1-c-d)/(f*(m+1))*Int[(e+f*x)^(m+1)*E^(2*a+2*b*x)/(1-c+d+(1-c-d)*E^(2*a+2*b*x)),x] -
  b*(1+c+d)/(f*(m+1))*Int[(e+f*x)^(m+1)*E^(2*a+2*b*x)/(1+c-d+(1+c+d)*E^(2*a+2*b*x)),x] /;
FreeQ[{a,b,c,d,e,f},x] && IGtQ[m,0] && NeQ[(c-d)^2,1]
```

```

Int[(e_.+f_.**x_)^m_.*ArcTanh[c_.+d_.*Coth[a_.+b_.**x_] ,x_Symbol] :=
  (e+f*x)^(m+1)*ArcTanh[c+d*Coth[a+b*x]]/(f*(m+1)) +
  b*(1+c+d)/(f*(m+1))*Int[(e+f*x)^(m+1)*E^(2*a+2*b*x)/(1+c-d-(1+c+d)*E^(2*a+2*b*x)),x] -
  b*(1-c-d)/(f*(m+1))*Int[(e+f*x)^(m+1)*E^(2*a+2*b*x)/(1-c+d-(1-c-d)*E^(2*a+2*b*x)),x] /;
FreeQ[{a,b,c,d,e,f},x] && IGtQ[m,0] && NeQ[(c-d)^2,1]

```

```

Int[(e_.+f_.**x_)^m_.*ArcCoth[c_.+d_.*Coth[a_.+b_.**x_] ,x_Symbol] :=
  (e+f*x)^(m+1)*ArcCoth[c+d*Coth[a+b*x]]/(f*(m+1)) +
  b*(1+c+d)/(f*(m+1))*Int[(e+f*x)^(m+1)*E^(2*a+2*b*x)/(1+c-d-(1+c+d)*E^(2*a+2*b*x)),x] -
  b*(1-c-d)/(f*(m+1))*Int[(e+f*x)^(m+1)*E^(2*a+2*b*x)/(1-c+d-(1-c-d)*E^(2*a+2*b*x)),x] /;
FreeQ[{a,b,c,d,e,f},x] && IGtQ[m,0] && NeQ[(c-d)^2,1]

```

$$7. \int u \operatorname{ArcTanh}[c + d \operatorname{Tan}[a + b x]] dx$$

$$1. \int u \operatorname{ArcTanh}[\operatorname{Tan}[a + b x]] dx$$

$$1: \int \operatorname{ArcTanh}[\operatorname{Tan}[a + b x]] dx$$

Derivation: Integration by parts

Basis: $\partial_x \operatorname{ArcTanh}[\operatorname{Tan}[a + b x]] = b \operatorname{Sec}[2 a + 2 b x]$

Rule:

$$\int \operatorname{ArcTanh}[\operatorname{Tan}[a + b x]] dx \rightarrow x \operatorname{ArcTanh}[\operatorname{Tan}[a + b x]] - b \int x \operatorname{Sec}[2 a + 2 b x] dx$$

Program code:

```

Int[ArcTanh[Tan[a_.+b_.**x_] ,x_Symbol] :=
  x*ArcTanh[Tan[a+b*x]] - b*Int[x*Sec[2*a+2*b*x],x] /;
FreeQ[{a,b},x]

```

```

Int[ArcCoth[Tan[a_.+b_.**x_] ,x_Symbol] :=
  x*ArcCoth[Tan[a+b*x]] - b*Int[x*Sec[2*a+2*b*x],x] /;
FreeQ[{a,b},x]

```

```

Int[ArcTanh[Cot[a_.+b_.**x_] ,x_Symbol] :=
  x*ArcTanh[Cot[a+b*x]] - b*Int[x*Sec[2*a+2*b*x],x] /;
FreeQ[{a,b},x]

```

```
Int[ArcCoth[Cot[a_+b_.*x_]],x_Symbol] :=
  x*ArcCoth[Cot[a+b*x]] - b*Int[x*Sec[2*a+2*b*x],x] /;
FreeQ[{a,b},x]
```

$$2: \int (e + f x)^m \operatorname{ArcTanh}[\operatorname{Tan}[a + b x]] dx \text{ when } m \in \mathbb{Z}^+$$

Derivation: Integration by parts

Basis: $\partial_x \operatorname{ArcTanh}[\operatorname{Tan}[a + b x]] = b \operatorname{Sec}[2 a + 2 b x]$

Rule: If $m \in \mathbb{Z}^+$, then

$$\int (e + f x)^m \operatorname{ArcTanh}[\operatorname{Tan}[a + b x]] dx \rightarrow \frac{(e + f x)^{m+1} \operatorname{ArcTanh}[\operatorname{Tan}[a + b x]]}{f (m + 1)} - \frac{b}{f (m + 1)} \int (e + f x)^{m+1} \operatorname{Sec}[2 a + 2 b x] dx$$

Program code:

```
Int[(e_+f_.*x_)^m_.*ArcTanh[Tan[a_+b_.*x_]],x_Symbol] :=
  (e+f*x)^(m+1)*ArcTanh[Tan[a+b*x]]/(f*(m+1)) - b/(f*(m+1))*Int[(e+f*x)^(m+1)*Sec[2*a+2*b*x],x] /;
FreeQ[{a,b,e,f},x] && IGtQ[m,0]
```

```
Int[(e_+f_.*x_)^m_.*ArcCoth[Tan[a_+b_.*x_]],x_Symbol] :=
  (e+f*x)^(m+1)*ArcCoth[Tan[a+b*x]]/(f*(m+1)) - b/(f*(m+1))*Int[(e+f*x)^(m+1)*Sec[2*a+2*b*x],x] /;
FreeQ[{a,b,e,f},x] && IGtQ[m,0]
```

```
Int[(e_+f_.*x_)^m_.*ArcTanh[Cot[a_+b_.*x_]],x_Symbol] :=
  (e+f*x)^(m+1)*ArcTanh[Cot[a+b*x]]/(f*(m+1)) - b/(f*(m+1))*Int[(e+f*x)^(m+1)*Sec[2*a+2*b*x],x] /;
FreeQ[{a,b,e,f},x] && IGtQ[m,0]
```

```
Int[(e_+f_.*x_)^m_.*ArcCoth[Cot[a_+b_.*x_]],x_Symbol] :=
  (e+f*x)^(m+1)*ArcCoth[Cot[a+b*x]]/(f*(m+1)) - b/(f*(m+1))*Int[(e+f*x)^(m+1)*Sec[2*a+2*b*x],x] /;
FreeQ[{a,b,e,f},x] && IGtQ[m,0]
```

$$2. \int u \operatorname{ArcTanh}[c + d \operatorname{Tan}[a + b x]] dx$$

$$1. \int \operatorname{ArcTanh}[c + d \operatorname{Tan}[a + b x]] dx$$

$$1: \int \operatorname{ArcTanh}[c + d \operatorname{Tan}[a + b x]] dx \text{ when } (c + i d)^2 = 1$$

Derivation: Integration by parts

$$\text{Basis: If } (c + i d)^2 = 1, \text{ then } \partial_x \operatorname{ArcTanh}[c + d \operatorname{Tan}[a + b x]] = - \frac{i b}{c + i d + c e^{2 i a + 2 i b x}}$$

Rule: If $(c + i d)^2 = 1$, then

$$\int \operatorname{ArcTanh}[c + d \operatorname{Tan}[a + b x]] dx \rightarrow x \operatorname{ArcTanh}[c + d \operatorname{Tan}[a + b x]] + i b \int \frac{x}{c + i d + c e^{2 i a + 2 i b x}} dx$$

Program code:

```
Int[ArcTanh[c_.+d_.*Tan[a_.+b_.*x_]],x_Symbol] :=
  x*ArcTanh[c+d*Tan[a+b*x]] +
  I*b*Int[x/(c+I*d+c*E^(2*I*a+2*I*b*x)),x] /;
FreeQ[{a,b,c,d},x] && EqQ[(c+I*d)^2,1]
```

```
Int[ArcCoth[c_.+d_.*Tan[a_.+b_.*x_]],x_Symbol] :=
  x*ArcCoth[c+d*Tan[a+b*x]] +
  I*b*Int[x/(c+I*d+c*E^(2*I*a+2*I*b*x)),x] /;
FreeQ[{a,b,c,d},x] && EqQ[(c+I*d)^2,1]
```

```
Int[ArcTanh[c_.+d_.*Cot[a_.+b_.*x_]],x_Symbol] :=
  x*ArcTanh[c+d*Cot[a+b*x]] +
  I*b*Int[x/(c-I*d-c*E^(2*I*a+2*I*b*x)),x] /;
FreeQ[{a,b,c,d},x] && EqQ[(c-I*d)^2,1]
```

```
Int[ArcCoth[c_.+d_.*Cot[a_.+b_.*x_]],x_Symbol] :=
  x*ArcCoth[c+d*Cot[a+b*x]] +
  I*b*Int[x/(c-I*d-c*E^(2*I*a+2*I*b*x)),x] /;
FreeQ[{a,b,c,d},x] && EqQ[(c-I*d)^2,1]
```

$$2: \int \text{ArcTanh}[c + d \tan[a + b x]] dx \text{ when } (c + i d)^2 \neq 1$$

Derivation: Integration by parts

$$\text{Basis: } \partial_x \text{ArcTanh}[c + d \tan[a + b x]] = -\frac{i b (1-c+i d) e^{2 i a+2 i b x}}{1-c-i d+(1-c+i d) e^{2 i a+2 i b x}} + \frac{i b (1+c-i d) e^{2 i a+2 i b x}}{1+c+i d+(1+c-i d) e^{2 i a+2 i b x}}$$

Rule: If $(c + i d)^2 \neq 1$, then

$$\int \text{ArcTanh}[c + d \tan[a + b x]] dx \rightarrow x \text{ArcTanh}[c + d \tan[a + b x]] + i b (1 - c + i d) \int \frac{x e^{2 i a+2 i b x}}{1 - c - i d + (1 - c + i d) e^{2 i a+2 i b x}} dx - i b (1 + c - i d) \int \frac{x e^{2 i a+2 i b x}}{1 + c + i d + (1 + c - i d) e^{2 i a+2 i b x}} dx$$

Program code:

```
Int[ArcTanh[c_.+d_.*Tan[a_.+b_.*x_]],x_Symbol] :=
  x*ArcTanh[c+d*Tan[a+b*x]] +
  I*b*(1-c+I*d)*Int[x*E^(2*I*a+2*I*b*x)/(1-c-I*d+(1-c+I*d)*E^(2*I*a+2*I*b*x)),x] -
  I*b*(1+c-I*d)*Int[x*E^(2*I*a+2*I*b*x)/(1+c+I*d+(1+c-I*d)*E^(2*I*a+2*I*b*x)),x] /;
FreeQ[{a,b,c,d},x] && NeQ[(c+I*d)^2,1]
```

```
Int[ArcCoth[c_.+d_.*Tan[a_.+b_.*x_]],x_Symbol] :=
  x*ArcCoth[c+d*Tan[a+b*x]] +
  I*b*(1-c+I*d)*Int[x*E^(2*I*a+2*I*b*x)/(1-c-I*d+(1-c+I*d)*E^(2*I*a+2*I*b*x)),x] -
  I*b*(1+c-I*d)*Int[x*E^(2*I*a+2*I*b*x)/(1+c+I*d+(1+c-I*d)*E^(2*I*a+2*I*b*x)),x] /;
FreeQ[{a,b,c,d},x] && NeQ[(c+I*d)^2,1]
```

```
Int[ArcTanh[c_.+d_.*Cot[a_.+b_.*x_]],x_Symbol] :=
  x*ArcTanh[c+d*Cot[a+b*x]] -
  I*b*(1-c-I*d)*Int[x*E^(2*I*a+2*I*b*x)/(1-c+I*d-(1-c-I*d)*E^(2*I*a+2*I*b*x)),x] +
  I*b*(1+c+I*d)*Int[x*E^(2*I*a+2*I*b*x)/(1+c-I*d-(1+c+I*d)*E^(2*I*a+2*I*b*x)),x] /;
FreeQ[{a,b,c,d},x] && NeQ[(c-I*d)^2,1]
```

```
Int[ArcCoth[c_.+d_.*Cot[a_.+b_.*x_]],x_Symbol] :=
  x*ArcCoth[c+d*Cot[a+b*x]] -
  I*b*(1-c-I*d)*Int[x*E^(2*I*a+2*I*b*x)/(1-c+I*d-(1-c-I*d)*E^(2*I*a+2*I*b*x)),x] +
  I*b*(1+c+I*d)*Int[x*E^(2*I*a+2*I*b*x)/(1+c-I*d-(1+c+I*d)*E^(2*I*a+2*I*b*x)),x] /;
FreeQ[{a,b,c,d},x] && NeQ[(c-I*d)^2,1]
```

$$2. \int (e + f x)^m \operatorname{ArcTanh}[c + d \operatorname{Tan}[a + b x]] dx \text{ when } m \in \mathbb{Z}^+$$

$$1: \int (e + f x)^m \operatorname{ArcTanh}[c + d \operatorname{Tan}[a + b x]] dx \text{ when } m \in \mathbb{Z}^+ \wedge (c + i d)^2 = 1$$

Derivation: Integration by parts

Basis: If $(c + i d)^2 = 1$, then $\partial_x \operatorname{ArcTanh}[c + d \operatorname{Tan}[a + b x]] = -\frac{i b}{c + i d + c e^{2 i a + 2 i b x}}$

Rule: If $m \in \mathbb{Z}^+ \wedge (c + i d)^2 = 1$, then

$$\int (e + f x)^m \operatorname{ArcTanh}[c + d \operatorname{Tan}[a + b x]] dx \rightarrow \frac{(e + f x)^{m+1} \operatorname{ArcTanh}[c + d \operatorname{Tan}[a + b x]]}{f (m + 1)} + \frac{i b}{f (m + 1)} \int \frac{(e + f x)^{m+1}}{c + i d + c e^{2 i a + 2 i b x}} dx$$

Program code:

```
Int[(e_+f_*x_)^m_*ArcTanh[c_+d_*Tan[a_+b_*x_]],x_Symbol] :=
  (e+f*x)^(m+1)*ArcTanh[c+d*Tan[a+b*x]]/(f*(m+1)) +
  I*b/(f*(m+1))*Int[(e+f*x)^(m+1)/(c+I*d+c*E^(2*I*a+2*I*b*x)),x] /;
FreeQ[{a,b,c,d,e,f},x] && IGtQ[m,0] && EqQ[(c+I*d)^2,1]
```

```
Int[(e_+f_*x_)^m_*ArcCoth[c_+d_*Tan[a_+b_*x_]],x_Symbol] :=
  (e+f*x)^(m+1)*ArcCoth[c+d*Tan[a+b*x]]/(f*(m+1)) +
  I*b/(f*(m+1))*Int[(e+f*x)^(m+1)/(c+I*d+c*E^(2*I*a+2*I*b*x)),x] /;
FreeQ[{a,b,c,d,e,f},x] && IGtQ[m,0] && EqQ[(c+I*d)^2,1]
```

```
Int[(e_+f_*x_)^m_*ArcTanh[c_+d_*Cot[a_+b_*x_]],x_Symbol] :=
  (e+f*x)^(m+1)*ArcTanh[c+d*Cot[a+b*x]]/(f*(m+1)) +
  I*b/(f*(m+1))*Int[(e+f*x)^(m+1)/(c-I*d-c*E^(2*I*a+2*I*b*x)),x] /;
FreeQ[{a,b,c,d,e,f},x] && IGtQ[m,0] && EqQ[(c-I*d)^2,1]
```

```
Int[(e_+f_*x_)^m_*ArcCoth[c_+d_*Cot[a_+b_*x_]],x_Symbol] :=
  (e+f*x)^(m+1)*ArcCoth[c+d*Cot[a+b*x]]/(f*(m+1)) +
  I*b/(f*(m+1))*Int[(e+f*x)^(m+1)/(c-I*d-c*E^(2*I*a+2*I*b*x)),x] /;
FreeQ[{a,b,c,d,e,f},x] && IGtQ[m,0] && EqQ[(c-I*d)^2,1]
```

$$2: \int (e + f x)^m \operatorname{ArcTanh}[c + d \operatorname{Tan}[a + b x]] dx \text{ when } m \in \mathbb{Z}^+ \wedge (c + i d)^2 \neq 1$$

Derivation: Integration by parts

$$\blacksquare \text{Basis: } \partial_x \operatorname{ArcTanh}[c + d \operatorname{Tan}[a + b x]] = -\frac{i b (1 - c + i d) e^{2 i a + 2 i b x}}{1 - c - i d + (1 - c + i d) e^{2 i a + 2 i b x}} + \frac{i b (1 + c - i d) e^{2 i a + 2 i b x}}{1 + c + i d + (1 + c - i d) e^{2 i a + 2 i b x}}$$

\blacksquare **Rule:** If $m \in \mathbb{Z}^+ \wedge (c + i d)^2 \neq 1$, then

$$\int (e + f x)^m \operatorname{ArcTanh}[c + d \operatorname{Tan}[a + b x]] dx \rightarrow \frac{(e + f x)^{m+1} \operatorname{ArcTanh}[c + d \operatorname{Tan}[a + b x]]}{f (m + 1)} + \frac{i b (1 - c + i d)}{f (m + 1)} \int \frac{(e + f x)^{m+1} e^{2 i a + 2 i b x}}{1 - c - i d + (1 - c + i d) e^{2 i a + 2 i b x}} dx - \frac{i b (1 + c - i d)}{f (m + 1)} \int \frac{(e + f x)^{m+1} e^{2 i a + 2 i b x}}{1 + c + i d + (1 + c - i d) e^{2 i a + 2 i b x}} dx$$

Program code:

```
Int[(e_+f_.*x_)^m_.*ArcTanh[c_+d_.*Tan[a_+b_.*x_] ,x_Symbol] :=
(e+f*x)^(m+1)*ArcTanh[c+d*Tan[a+b*x]]/(f*(m+1)) +
I*b*(1-c+I*d)/(f*(m+1))*Int[(e+f*x)^(m+1)*E^(2*I*a+2*I*b*x)/(1-c-I*d+(1-c+I*d)*E^(2*I*a+2*I*b*x)),x] -
I*b*(1+c-I*d)/(f*(m+1))*Int[(e+f*x)^(m+1)*E^(2*I*a+2*I*b*x)/(1+c+I*d+(1+c-I*d)*E^(2*I*a+2*I*b*x)),x] /;
FreeQ[{a,b,c,d,e,f},x] && IGtQ[m,0] && NeQ[(c+I*d)^2,1]
```

```
Int[(e_+f_.*x_)^m_.*ArcCoth[c_+d_.*Tan[a_+b_.*x_] ,x_Symbol] :=
(e+f*x)^(m+1)*ArcCoth[c+d*Tan[a+b*x]]/(f*(m+1)) +
I*b*(1-c+I*d)/(f*(m+1))*Int[(e+f*x)^(m+1)*E^(2*I*a+2*I*b*x)/(1-c-I*d+(1-c+I*d)*E^(2*I*a+2*I*b*x)),x] -
I*b*(1+c-I*d)/(f*(m+1))*Int[(e+f*x)^(m+1)*E^(2*I*a+2*I*b*x)/(1+c+I*d+(1+c-I*d)*E^(2*I*a+2*I*b*x)),x] /;
FreeQ[{a,b,c,d,e,f},x] && IGtQ[m,0] && NeQ[(c+I*d)^2,1]
```

```
Int[(e_+f_.*x_)^m_.*ArcTanh[c_+d_.*Cot[a_+b_.*x_] ,x_Symbol] :=
(e+f*x)^(m+1)*ArcTanh[c+d*Cot[a+b*x]]/(f*(m+1)) -
I*b*(1-c-I*d)/(f*(m+1))*Int[(e+f*x)^(m+1)*E^(2*I*a+2*I*b*x)/(1-c+I*d-(1-c-I*d)*E^(2*I*a+2*I*b*x)),x] +
I*b*(1+c+I*d)/(f*(m+1))*Int[(e+f*x)^(m+1)*E^(2*I*a+2*I*b*x)/(1+c-I*d-(1+c+I*d)*E^(2*I*a+2*I*b*x)),x] /;
FreeQ[{a,b,c,d,e,f},x] && IGtQ[m,0] && NeQ[(c-I*d)^2,1]
```

```
Int[(e_+f_.*x_)^m_.*ArcCoth[c_+d_.*Cot[a_+b_.*x_] ,x_Symbol] :=
(e+f*x)^(m+1)*ArcCoth[c+d*Cot[a+b*x]]/(f*(m+1)) -
I*b*(1-c-I*d)/(f*(m+1))*Int[(e+f*x)^(m+1)*E^(2*I*a+2*I*b*x)/(1-c+I*d-(1-c-I*d)*E^(2*I*a+2*I*b*x)),x] +
I*b*(1+c+I*d)/(f*(m+1))*Int[(e+f*x)^(m+1)*E^(2*I*a+2*I*b*x)/(1+c-I*d-(1+c+I*d)*E^(2*I*a+2*I*b*x)),x] /;
FreeQ[{a,b,c,d,e,f},x] && IGtQ[m,0] && NeQ[(c-I*d)^2,1]
```

8. $\int v (a + b \operatorname{ArcTanh}[u]) dx$ when u is free of inverse functions

1: $\int \operatorname{ArcTanh}[u] dx$ when u is free of inverse functions

Derivation: Integration by parts

Rule: If u is free of inverse functions, then

$$\int \operatorname{ArcTanh}[u] dx \rightarrow x \operatorname{ArcTanh}[u] - \int \frac{x \partial_x u}{1 - u^2} dx$$

Program code:

```
Int[ArcTanh[u_], x_Symbol] :=
  x*ArcTanh[u] -
  Int[SimplifyIntegrand[x*D[u,x]/(1-u^2),x],x] /;
InverseFunctionFreeQ[u,x]
```

```
Int[ArcCoth[u_], x_Symbol] :=
  x*ArcCoth[u] -
  Int[SimplifyIntegrand[x*D[u,x]/(1-u^2),x],x] /;
InverseFunctionFreeQ[u,x]
```

2: $\int (c + dx)^m (a + b \operatorname{ArcTanh}[u]) dx$ when $m \neq -1 \wedge u$ is free of inverse functions

Derivation: Integration by parts

Rule: If $m \neq -1 \wedge u$ is free of inverse functions, then

$$\int (c + dx)^m (a + b \operatorname{ArcTanh}[u]) dx \rightarrow \frac{(c + dx)^{m+1} (a + b \operatorname{ArcTanh}[u])}{d(m+1)} - \frac{b}{d(m+1)} \int \frac{(c + dx)^{m+1} \partial_x u}{1 - u^2} dx$$

Program code:

```
Int[(c_+d_*x_)^m_.*(a_+b_*ArcTanh[u_]),x_Symbol] :=
  (c+d*x)^(m+1)*(a+b*ArcTanh[u])/(d*(m+1)) -
  b/(d*(m+1))*Int[SimplifyIntegrand[(c+d*x)^(m+1)*D[u,x]/(1-u^2),x],x] /;
FreeQ[{a,b,c,d,m},x] && NeQ[m,-1] && InverseFunctionFreeQ[u,x] && Not[FunctionOfQ[(c+d*x)^(m+1),u,x]] && FalseQ[PowerVariableExpn[u
```

```

Int[(c_.+d_.*x_)^m_.*(a_.+b_.*ArcCoth[u_]),x_Symbol] :=
  (c+d*x)^(m+1)*(a+b*ArcCoth[u])/(d*(m+1)) -
  b/(d*(m+1))*Int[SimplifyIntegrand[(c+d*x)^(m+1)*D[u,x]/(1-u^2),x],x] /;
FreeQ[{a,b,c,d,m},x] && NeQ[m,-1] && InverseFunctionFreeQ[u,x] && Not[FunctionOfQ[(c+d*x)^(m+1),u,x]] && FalseQ[PowerVariableExpn[u

```

3: $\int v (a + b \operatorname{ArcTanh}[u]) dx$ when u and $\int v dx$ are free of inverse functions

Derivation: Integration by parts

Rule: If u is free of inverse functions, let $w = \int v dx$, if w is free of inverse functions, then

$$\int v (a + b \operatorname{ArcTanh}[u]) dx \rightarrow w (a + b \operatorname{ArcTanh}[u]) - b \int \frac{w \partial_x u}{1 - u^2} dx$$

Program code:

```

Int[v_*(a_.+b_.*ArcTanh[u_]),x_Symbol] :=
  With[{w=IntHide[v,x]},
    Dist[(a+b*ArcTanh[u]),w,x] - b*Int[SimplifyIntegrand[w*D[u,x]/(1-u^2),x],x] /;
    InverseFunctionFreeQ[w,x] /;
    FreeQ[{a,b},x] && InverseFunctionFreeQ[u,x] && Not[MatchQ[v, (c_.+d_.*x)^m_. /; FreeQ[{c,d,m},x]]] && FalseQ[FunctionOfLinear[v*(a+

```

```

Int[v_*(a_.+b_.*ArcCoth[u_]),x_Symbol] :=
  With[{w=IntHide[v,x]},
    Dist[(a+b*ArcCoth[u]),w,x] - b*Int[SimplifyIntegrand[w*D[u,x]/(1-u^2),x],x] /;
    InverseFunctionFreeQ[w,x] /;
    FreeQ[{a,b},x] && InverseFunctionFreeQ[u,x] && Not[MatchQ[v, (c_.+d_.*x)^m_. /; FreeQ[{c,d,m},x]]] && FalseQ[FunctionOfLinear[v*(a+

```