

# Mathematica 11.3 Integration Test Results

on the problems in the test-suite directory "1 Algebraic functions\1.2  
Trinomial products\1.2.4 Improper"

Test results for the 140 problems in "1.2.4.2 (d x)^m (a x^q+b x^n+c x^(2 n-q))^p.m"

Problem 104: Result more than twice size of optimal antiderivative.

$$\int \frac{x}{\sqrt{a x + b x^3 + c x^5}} dx$$

Optimal (type 6, 142 leaves, 3 steps):

$$\frac{2 x^2 \sqrt{1 + \frac{2 c x^2}{b - \sqrt{b^2 - 4 a c}}} \sqrt{1 + \frac{2 c x^2}{b + \sqrt{b^2 - 4 a c}}} \operatorname{AppellF1}\left[\frac{3}{4}, \frac{1}{2}, \frac{1}{2}, \frac{7}{4}, -\frac{2 c x^2}{b - \sqrt{b^2 - 4 a c}}, -\frac{2 c x^2}{b + \sqrt{b^2 - 4 a c}}\right]}{3 \sqrt{a x + b x^3 + c x^5}}$$

Result (type 6, 383 leaves):

$$\begin{aligned} & - \left( \left( 14 a^2 x^3 \left( b - \sqrt{b^2 - 4 a c} + 2 c x^2 \right) \left( b + \sqrt{b^2 - 4 a c} + 2 c x^2 \right) \operatorname{AppellF1}\left[\frac{3}{4}, \frac{1}{2}, \frac{1}{2}, \frac{7}{4}, -\frac{2 c x^2}{b + \sqrt{b^2 - 4 a c}}, \frac{2 c x^2}{-b + \sqrt{b^2 - 4 a c}}\right] \right) / \right. \\ & \left( 3 \left( b - \sqrt{b^2 - 4 a c} \right) \left( b + \sqrt{b^2 - 4 a c} \right) \left( x \left( a + b x^2 + c x^4 \right) \right)^{3/2} \right. \\ & \left. \left( -7 a \operatorname{AppellF1}\left[\frac{3}{4}, \frac{1}{2}, \frac{1}{2}, \frac{7}{4}, -\frac{2 c x^2}{b + \sqrt{b^2 - 4 a c}}, \frac{2 c x^2}{-b + \sqrt{b^2 - 4 a c}}\right] + x^2 \left( \left( b + \sqrt{b^2 - 4 a c} \right) \operatorname{AppellF1}\left[\frac{7}{4}, \frac{1}{2}, \frac{3}{2}, \frac{11}{4}, \right. \right. \right. \right. \\ & \left. \left. \left. -\frac{2 c x^2}{b + \sqrt{b^2 - 4 a c}}, \frac{2 c x^2}{-b + \sqrt{b^2 - 4 a c}} \right] + \left( b - \sqrt{b^2 - 4 a c} \right) \operatorname{AppellF1}\left[\frac{7}{4}, \frac{3}{2}, \frac{1}{2}, \frac{11}{4}, -\frac{2 c x^2}{b + \sqrt{b^2 - 4 a c}}, \frac{2 c x^2}{-b + \sqrt{b^2 - 4 a c}} \right] \right) \right) \right) \end{aligned}$$

Problem 105: Result unnecessarily involves imaginary or complex numbers.

$$\int x^{3/2} \sqrt{a x + b x^3 + c x^5} dx$$

Optimal (type 4, 380 leaves, 5 steps):

$$\begin{aligned}
 & - \frac{2 (b^2 - 3 a c) x^{3/2} (a + b x^2 + c x^4)}{15 c^{3/2} (\sqrt{a} + \sqrt{c} x^2) \sqrt{a x + b x^3 + c x^5}} + \frac{\sqrt{x} (b + 3 c x^2) \sqrt{a x + b x^3 + c x^5}}{15 c} + \\
 & \frac{2 a^{1/4} (b^2 - 3 a c) \sqrt{x} (\sqrt{a} + \sqrt{c} x^2) \sqrt{\frac{a + b x^2 + c x^4}{(\sqrt{a} + \sqrt{c} x^2)^2}} \operatorname{EllipticE}\left[2 \operatorname{ArcTan}\left[\frac{c^{1/4} x}{a^{1/4}}\right], \frac{1}{4} \left(2 - \frac{b}{\sqrt{a} \sqrt{c}}\right)\right]}{15 c^{7/4} \sqrt{a x + b x^3 + c x^5}} - \frac{1}{30 c^{7/4} \sqrt{a x + b x^3 + c x^5}} \\
 & a^{1/4} \left(2 b^2 + \sqrt{a} b \sqrt{c} - 6 a c\right) \sqrt{x} (\sqrt{a} + \sqrt{c} x^2) \sqrt{\frac{a + b x^2 + c x^4}{(\sqrt{a} + \sqrt{c} x^2)^2}} \operatorname{EllipticF}\left[2 \operatorname{ArcTan}\left[\frac{c^{1/4} x}{a^{1/4}}\right], \frac{1}{4} \left(2 - \frac{b}{\sqrt{a} \sqrt{c}}\right)\right]
 \end{aligned}$$

Result (type 4, 486 leaves):

$$\begin{aligned}
 & \frac{1}{30 c^2 \sqrt{\frac{c}{b + \sqrt{b^2 - 4 a c}}} \sqrt{x (a + b x^2 + c x^4)}} \\
 & \sqrt{x} \left( 2 c \sqrt{\frac{c}{b + \sqrt{b^2 - 4 a c}}} x (b + 3 c x^2) (a + b x^2 + c x^4) - i (b^2 - 3 a c) \left(-b + \sqrt{b^2 - 4 a c}\right) \sqrt{\frac{b + \sqrt{b^2 - 4 a c} + 2 c x^2}{b + \sqrt{b^2 - 4 a c}}} \sqrt{\frac{2 b - 2 \sqrt{b^2 - 4 a c} + 4 c x^2}{b - \sqrt{b^2 - 4 a c}}} \right. \\
 & \left. \operatorname{EllipticE}\left[i \operatorname{ArcSinh}\left[\sqrt{2} \sqrt{\frac{c}{b + \sqrt{b^2 - 4 a c}}} x\right], \frac{b + \sqrt{b^2 - 4 a c}}{b - \sqrt{b^2 - 4 a c}}\right] + i \left(-b^3 + 4 a b c + b^2 \sqrt{b^2 - 4 a c} - 3 a c \sqrt{b^2 - 4 a c}\right) \right. \\
 & \left. \sqrt{\frac{b + \sqrt{b^2 - 4 a c} + 2 c x^2}{b + \sqrt{b^2 - 4 a c}}} \sqrt{\frac{2 b - 2 \sqrt{b^2 - 4 a c} + 4 c x^2}{b - \sqrt{b^2 - 4 a c}}} \operatorname{EllipticF}\left[i \operatorname{ArcSinh}\left[\sqrt{2} \sqrt{\frac{c}{b + \sqrt{b^2 - 4 a c}}} x\right], \frac{b + \sqrt{b^2 - 4 a c}}{b - \sqrt{b^2 - 4 a c}}\right] \right)
 \end{aligned}$$

**Problem 107: Result unnecessarily involves imaginary or complex numbers.**

$$\int \frac{\sqrt{a x + b x^3 + c x^5}}{\sqrt{x}} dx$$

Optimal (type 4, 347 leaves, 5 steps):

$$\frac{b x^{3/2} (a + b x^2 + c x^4)}{3 \sqrt{c} (\sqrt{a} + \sqrt{c} x^2) \sqrt{a x + b x^3 + c x^5}} + \frac{1}{3} \sqrt{x} \sqrt{a x + b x^3 + c x^5} -$$

$$\frac{a^{1/4} b \sqrt{x} (\sqrt{a} + \sqrt{c} x^2) \sqrt{\frac{a + b x^2 + c x^4}{(\sqrt{a} + \sqrt{c} x^2)^2}} \text{EllipticE}\left[2 \text{ArcTan}\left[\frac{c^{1/4} x}{a^{1/4}}\right], \frac{1}{4} \left(2 - \frac{b}{\sqrt{a} \sqrt{c}}\right)\right]}{3 c^{3/4} \sqrt{a x + b x^3 + c x^5}} +$$

$$\frac{a^{1/4} (b + 2 \sqrt{a} \sqrt{c}) \sqrt{x} (\sqrt{a} + \sqrt{c} x^2) \sqrt{\frac{a + b x^2 + c x^4}{(\sqrt{a} + \sqrt{c} x^2)^2}} \text{EllipticF}\left[2 \text{ArcTan}\left[\frac{c^{1/4} x}{a^{1/4}}\right], \frac{1}{4} \left(2 - \frac{b}{\sqrt{a} \sqrt{c}}\right)\right]}{6 c^{3/4} \sqrt{a x + b x^3 + c x^5}}$$

Result (type 4, 452 leaves):

$$\frac{1}{12 c \sqrt{\frac{c}{b + \sqrt{b^2 - 4 a c}}} \sqrt{x (a + b x^2 + c x^4)}}$$

$$\sqrt{x} \left( 4 c \sqrt{\frac{c}{b + \sqrt{b^2 - 4 a c}}} x (a + b x^2 + c x^4) + i b (-b + \sqrt{b^2 - 4 a c}) \sqrt{\frac{b + \sqrt{b^2 - 4 a c} + 2 c x^2}{b + \sqrt{b^2 - 4 a c}}} \sqrt{\frac{2 b - 2 \sqrt{b^2 - 4 a c} + 4 c x^2}{b - \sqrt{b^2 - 4 a c}}} \right.$$

$$\left. \text{EllipticE}\left[i \text{ArcSinh}\left[\sqrt{2} \sqrt{\frac{c}{b + \sqrt{b^2 - 4 a c}}} x\right], \frac{b + \sqrt{b^2 - 4 a c}}{b - \sqrt{b^2 - 4 a c}}\right] - i (-b^2 + 4 a c + b \sqrt{b^2 - 4 a c}) \right.$$

$$\left. \sqrt{\frac{b + \sqrt{b^2 - 4 a c} + 2 c x^2}{b + \sqrt{b^2 - 4 a c}}} \sqrt{\frac{2 b - 2 \sqrt{b^2 - 4 a c} + 4 c x^2}{b - \sqrt{b^2 - 4 a c}}} \text{EllipticF}\left[i \text{ArcSinh}\left[\sqrt{2} \sqrt{\frac{c}{b + \sqrt{b^2 - 4 a c}}} x\right], \frac{b + \sqrt{b^2 - 4 a c}}{b - \sqrt{b^2 - 4 a c}}\right] \right)$$

**Problem 110: Result unnecessarily involves imaginary or complex numbers.**

$$\int \sqrt{x} (a x + b x^3 + c x^5)^{3/2} dx$$

Optimal (type 4, 487 leaves, 6 steps):

$$\frac{(8b^4 - 57ab^2c + 84a^2c^2)x^{3/2}(a + bx^2 + cx^4)}{315c^{5/2}(\sqrt{a} + \sqrt{c}x^2)\sqrt{ax + bx^3 + cx^5}} - \frac{\sqrt{x}(b(4b^2 - 9ac) + 6c(2b^2 - 7ac)x^2)\sqrt{ax + bx^3 + cx^5}}{315c^2} + \frac{(3b + 7cx^2)(ax + bx^3 + cx^5)^{3/2}}{63c\sqrt{x}} -$$

$$\left( a^{1/4}(8b^4 - 57ab^2c + 84a^2c^2)\sqrt{x}(\sqrt{a} + \sqrt{c}x^2)\sqrt{\frac{a + bx^2 + cx^4}{(\sqrt{a} + \sqrt{c}x^2)^2}} \text{EllipticE}\left[2 \text{ArcTan}\left[\frac{c^{1/4}x}{a^{1/4}}\right], \frac{1}{4}\left(2 - \frac{b}{\sqrt{a}\sqrt{c}}\right)\right] \right) /$$

$$\left( 315c^{11/4}\sqrt{ax + bx^3 + cx^5} \right) +$$

$$\left( a^{1/4}(8b^4 - 57ab^2c + 84a^2c^2 + 4\sqrt{a}b\sqrt{c}(b^2 - 6ac))\sqrt{x}(\sqrt{a} + \sqrt{c}x^2)\sqrt{\frac{a + bx^2 + cx^4}{(\sqrt{a} + \sqrt{c}x^2)^2}} \text{EllipticF}\left[2 \text{ArcTan}\left[\frac{c^{1/4}x}{a^{1/4}}\right], \frac{1}{4}\left(2 - \frac{b}{\sqrt{a}\sqrt{c}}\right)\right] \right) /$$

$$\left( 630c^{11/4}\sqrt{ax + bx^3 + cx^5} \right)$$

Result (type 4, 609 leaves):

$$\frac{1}{1260c^3\sqrt{\frac{c}{b + \sqrt{b^2 - 4ac}}}\sqrt{x(a + bx^2 + cx^4)}}\sqrt{x}$$

$$\left( 4c\sqrt{\frac{c}{b + \sqrt{b^2 - 4ac}}}\sqrt{x(-4b^4x^2 - b^3cx^4 + 53b^2c^2x^6 + 85b^3c^3x^8 + 35c^4x^{10} + a^2c(24b + 77cx^2) + a(-4b^3 + 27b^2cx^2 + 151bc^2x^4 + 112c^3x^6))} \right) +$$

$$i(8b^4 - 57ab^2c + 84a^2c^2)(-b + \sqrt{b^2 - 4ac})\sqrt{\frac{b + \sqrt{b^2 - 4ac} + 2cx^2}{b + \sqrt{b^2 - 4ac}}}$$

$$\sqrt{\frac{2b - 2\sqrt{b^2 - 4ac} + 4cx^2}{b - \sqrt{b^2 - 4ac}}}\text{EllipticE}\left[i \text{ArcSinh}\left[\sqrt{2}\sqrt{\frac{c}{b + \sqrt{b^2 - 4ac}}}\sqrt{x}\right], \frac{b + \sqrt{b^2 - 4ac}}{b - \sqrt{b^2 - 4ac}}\right] -$$

$$i(-8b^5 + 65ab^3c - 132a^2bc^2 + 8b^4\sqrt{b^2 - 4ac} - 57ab^2c\sqrt{b^2 - 4ac} + 84a^2c^2\sqrt{b^2 - 4ac})\sqrt{\frac{b + \sqrt{b^2 - 4ac} + 2cx^2}{b + \sqrt{b^2 - 4ac}}}$$

$$\sqrt{\frac{2b - 2\sqrt{b^2 - 4ac} + 4cx^2}{b - \sqrt{b^2 - 4ac}}}\text{EllipticF}\left[i \text{ArcSinh}\left[\sqrt{2}\sqrt{\frac{c}{b + \sqrt{b^2 - 4ac}}}\sqrt{x}\right], \frac{b + \sqrt{b^2 - 4ac}}{b - \sqrt{b^2 - 4ac}}\right]$$

### Problem 112: Result unnecessarily involves imaginary or complex numbers.

$$\int \frac{(a x + b x^3 + c x^5)^{3/2}}{x^{3/2}} dx$$

Optimal (type 4, 425 leaves, 6 steps):

$$\begin{aligned} & -\frac{2 b (b^2 - 8 a c) x^{3/2} (a + b x^2 + c x^4)}{35 c^{3/2} (\sqrt{a} + \sqrt{c} x^2) \sqrt{a x + b x^3 + c x^5}} + \frac{\sqrt{x} (b^2 + 10 a c + 3 b c x^2) \sqrt{a x + b x^3 + c x^5}}{35 c} + \frac{(a x + b x^3 + c x^5)^{3/2}}{7 \sqrt{x}} + \\ & \frac{2 a^{1/4} b (b^2 - 8 a c) \sqrt{x} (\sqrt{a} + \sqrt{c} x^2) \sqrt{\frac{a + b x^2 + c x^4}{(\sqrt{a} + \sqrt{c} x^2)^2}} \text{EllipticE}\left[2 \text{ArcTan}\left[\frac{c^{1/4} x}{a^{1/4}}\right], \frac{1}{4} \left(2 - \frac{b}{\sqrt{a} \sqrt{c}}\right)\right]}{35 c^{7/4} \sqrt{a x + b x^3 + c x^5}} - \frac{1}{70 c^{7/4} \sqrt{a x + b x^3 + c x^5}} \\ & a^{1/4} \left(\sqrt{a} \sqrt{c} (b^2 - 20 a c) + 2 b (b^2 - 8 a c)\right) \sqrt{x} (\sqrt{a} + \sqrt{c} x^2) \sqrt{\frac{a + b x^2 + c x^4}{(\sqrt{a} + \sqrt{c} x^2)^2}} \text{EllipticF}\left[2 \text{ArcTan}\left[\frac{c^{1/4} x}{a^{1/4}}\right], \frac{1}{4} \left(2 - \frac{b}{\sqrt{a} \sqrt{c}}\right)\right] \end{aligned}$$

Result (type 4, 540 leaves):

$$\begin{aligned} & \frac{1}{70 c^2 \sqrt{\frac{c}{b + \sqrt{b^2 - 4 a c}}} \sqrt{x (a + b x^2 + c x^4)}} \\ & \sqrt{x} \left( 2 c \sqrt{\frac{c}{b + \sqrt{b^2 - 4 a c}}} x (15 a^2 c + a (b^2 + 23 b c x^2 + 20 c^2 x^4) + x^2 (b^3 + 9 b^2 c x^2 + 13 b c^2 x^4 + 5 c^3 x^6)) - i b (b^2 - 8 a c) \left(-b + \sqrt{b^2 - 4 a c}\right) \right. \\ & \left. \sqrt{\frac{b + \sqrt{b^2 - 4 a c} + 2 c x^2}{b + \sqrt{b^2 - 4 a c}}} \sqrt{\frac{2 b - 2 \sqrt{b^2 - 4 a c} + 4 c x^2}{b - \sqrt{b^2 - 4 a c}}} \text{EllipticE}\left[i \text{ArcSinh}\left[\sqrt{2} \sqrt{\frac{c}{b + \sqrt{b^2 - 4 a c}}} x\right], \frac{b + \sqrt{b^2 - 4 a c}}{b - \sqrt{b^2 - 4 a c}}\right] + \right. \\ & \left. i \left(-b^4 + 9 a b^2 c - 20 a^2 c^2 + b^3 \sqrt{b^2 - 4 a c} - 8 a b c \sqrt{b^2 - 4 a c}\right) \sqrt{\frac{b + \sqrt{b^2 - 4 a c} + 2 c x^2}{b + \sqrt{b^2 - 4 a c}}} \right. \\ & \left. \sqrt{\frac{2 b - 2 \sqrt{b^2 - 4 a c} + 4 c x^2}{b - \sqrt{b^2 - 4 a c}}} \text{EllipticF}\left[i \text{ArcSinh}\left[\sqrt{2} \sqrt{\frac{c}{b + \sqrt{b^2 - 4 a c}}} x\right], \frac{b + \sqrt{b^2 - 4 a c}}{b - \sqrt{b^2 - 4 a c}}\right] \right) \end{aligned}$$

**Problem 114:** Result unnecessarily involves imaginary or complex numbers.

$$\int \frac{\sqrt{x}}{\sqrt{ax + bx^3 + cx^5}} dx$$

Optimal (type 4, 121 leaves, 2 steps):

$$\frac{\sqrt{x} \left( \sqrt{a} + \sqrt{c} x^2 \right) \sqrt{\frac{a+bx^2+cx^4}{(\sqrt{a}+\sqrt{c}x^2)^2}} \operatorname{EllipticF}\left[2 \operatorname{ArcTan}\left[\frac{c^{1/4}x}{a^{1/4}}\right], \frac{1}{4} \left(2 - \frac{b}{\sqrt{a}\sqrt{c}}\right)\right]}{2 a^{1/4} c^{1/4} \sqrt{ax + bx^3 + cx^5}}$$

Result (type 4, 193 leaves):

$$\frac{i \sqrt{x} \sqrt{\frac{b+\sqrt{b^2-4ac}+2cx^2}{b+\sqrt{b^2-4ac}}} \sqrt{1 + \frac{2cx^2}{b-\sqrt{b^2-4ac}}} \operatorname{EllipticF}\left[i \operatorname{ArcSinh}\left[\sqrt{2} \sqrt{\frac{c}{b+\sqrt{b^2-4ac}}} x\right], \frac{b+\sqrt{b^2-4ac}}{b-\sqrt{b^2-4ac}}\right]}{\sqrt{2} \sqrt{\frac{c}{b+\sqrt{b^2-4ac}}} \sqrt{x(a+bx^2+cx^4)}}$$

**Problem 116:** Result unnecessarily involves imaginary or complex numbers.

$$\int \frac{1}{x^{3/2} \sqrt{ax + bx^3 + cx^5}} dx$$

Optimal (type 4, 330 leaves, 6 steps):

$$\frac{\sqrt{c} x^{3/2} (a + bx^2 + cx^4)}{a (\sqrt{a} + \sqrt{c} x^2) \sqrt{ax + bx^3 + cx^5}} - \frac{\sqrt{ax + bx^3 + cx^5}}{a x^{3/2}} - \frac{c^{1/4} \sqrt{x} (\sqrt{a} + \sqrt{c} x^2) \sqrt{\frac{a+bx^2+cx^4}{(\sqrt{a}+\sqrt{c}x^2)^2}} \operatorname{EllipticE}\left[2 \operatorname{ArcTan}\left[\frac{c^{1/4}x}{a^{1/4}}\right], \frac{1}{4} \left(2 - \frac{b}{\sqrt{a}\sqrt{c}}\right)\right]}{a^{3/4} \sqrt{ax + bx^3 + cx^5}} +$$

$$\frac{c^{1/4} \sqrt{x} (\sqrt{a} + \sqrt{c} x^2) \sqrt{\frac{a+bx^2+cx^4}{(\sqrt{a}+\sqrt{c}x^2)^2}} \operatorname{EllipticF}\left[2 \operatorname{ArcTan}\left[\frac{c^{1/4}x}{a^{1/4}}\right], \frac{1}{4} \left(2 - \frac{b}{\sqrt{a}\sqrt{c}}\right)\right]}{2 a^{3/4} \sqrt{ax + bx^3 + cx^5}}$$

Result (type 4, 303 leaves):

$$\left( -4 (a + b x^2 + c x^4) + \frac{1}{\sqrt{\frac{c}{b + \sqrt{b^2 - 4ac}}}} \right. \\ \left. i \sqrt{2} \left( -b + \sqrt{b^2 - 4ac} \right) x \sqrt{\frac{b + \sqrt{b^2 - 4ac} + 2cx^2}{b + \sqrt{b^2 - 4ac}}} \sqrt{1 + \frac{2cx^2}{b - \sqrt{b^2 - 4ac}}} \left( \text{EllipticE} \left[ i \text{ArcSinh} \left[ \sqrt{2} \sqrt{\frac{c}{b + \sqrt{b^2 - 4ac}}} x \right], \frac{b + \sqrt{b^2 - 4ac}}{b - \sqrt{b^2 - 4ac}} \right] - \right. \right. \\ \left. \left. \text{EllipticF} \left[ i \text{ArcSinh} \left[ \sqrt{2} \sqrt{\frac{c}{b + \sqrt{b^2 - 4ac}}} x \right], \frac{b + \sqrt{b^2 - 4ac}}{b - \sqrt{b^2 - 4ac}} \right] \right) \right) / \left( 4a \sqrt{x} \sqrt{x(a + bx^2 + cx^4)} \right)$$

**Problem 117: Result unnecessarily involves imaginary or complex numbers.**

$$\int \frac{x^{3/2}}{(ax + bx^3 + cx^5)^{3/2}} dx$$

Optimal (type 4, 391 leaves, 5 steps):

$$\frac{x^{3/2} (b^2 - 2ac + bcx^2)}{a(b^2 - 4ac) \sqrt{ax + bx^3 + cx^5}} - \frac{b \sqrt{c} x^{3/2} (a + bx^2 + cx^4)}{a(b^2 - 4ac) (\sqrt{a} + \sqrt{c} x^2) \sqrt{ax + bx^3 + cx^5}} + \\ \frac{bc^{1/4} \sqrt{x} (\sqrt{a} + \sqrt{c} x^2) \sqrt{\frac{a+bx^2+cx^4}{(\sqrt{a} + \sqrt{c} x^2)^2}} \text{EllipticE} \left[ 2 \text{ArcTan} \left[ \frac{c^{1/4} x}{a^{1/4}} \right], \frac{1}{4} \left( 2 - \frac{b}{\sqrt{a} \sqrt{c}} \right) \right]}{a^{3/4} (b^2 - 4ac) \sqrt{ax + bx^3 + cx^5}} - \\ \frac{c^{1/4} \sqrt{x} (\sqrt{a} + \sqrt{c} x^2) \sqrt{\frac{a+bx^2+cx^4}{(\sqrt{a} + \sqrt{c} x^2)^2}} \text{EllipticF} \left[ 2 \text{ArcTan} \left[ \frac{c^{1/4} x}{a^{1/4}} \right], \frac{1}{4} \left( 2 - \frac{b}{\sqrt{a} \sqrt{c}} \right) \right]}{2a^{3/4} (b - 2\sqrt{a} \sqrt{c}) \sqrt{ax + bx^3 + cx^5}}$$

Result (type 4, 463 leaves):

$$\begin{aligned}
& - \frac{1}{4 a (b^2 - 4 a c) \sqrt{\frac{c}{b + \sqrt{b^2 - 4 a c}}} \sqrt{x (a + b x^2 + c x^4)}} \\
& \sqrt{x} \left( -4 \sqrt{\frac{c}{b + \sqrt{b^2 - 4 a c}}} x (b^2 - 2 a c + b c x^2) + i b (-b + \sqrt{b^2 - 4 a c}) \sqrt{\frac{b + \sqrt{b^2 - 4 a c} + 2 c x^2}{b + \sqrt{b^2 - 4 a c}}} \sqrt{\frac{2 b - 2 \sqrt{b^2 - 4 a c} + 4 c x^2}{b - \sqrt{b^2 - 4 a c}}} \right. \\
& \text{EllipticE} \left[ i \text{ArcSinh} \left[ \sqrt{2} \sqrt{\frac{c}{b + \sqrt{b^2 - 4 a c}}} x \right], \frac{b + \sqrt{b^2 - 4 a c}}{b - \sqrt{b^2 - 4 a c}} \right] - i (-b^2 + 4 a c + b \sqrt{b^2 - 4 a c}) \\
& \left. \sqrt{\frac{b + \sqrt{b^2 - 4 a c} + 2 c x^2}{b + \sqrt{b^2 - 4 a c}}} \sqrt{\frac{2 b - 2 \sqrt{b^2 - 4 a c} + 4 c x^2}{b - \sqrt{b^2 - 4 a c}}} \text{EllipticF} \left[ i \text{ArcSinh} \left[ \sqrt{2} \sqrt{\frac{c}{b + \sqrt{b^2 - 4 a c}}} x \right], \frac{b + \sqrt{b^2 - 4 a c}}{b - \sqrt{b^2 - 4 a c}} \right] \right)
\end{aligned}$$

**Problem 119: Result unnecessarily involves imaginary or complex numbers.**

$$\int \frac{1}{\sqrt{x} (a x + b x^3 + c x^5)^{3/2}} dx$$

Optimal (type 4, 468 leaves, 6 steps):

$$\begin{aligned}
& \frac{b^2 - 2 a c + b c x^2}{a (b^2 - 4 a c) \sqrt{x} \sqrt{a x + b x^3 + c x^5}} + \frac{2 \sqrt{c} (b^2 - 3 a c) x^{3/2} (a + b x^2 + c x^4)}{a^2 (b^2 - 4 a c) (\sqrt{a} + \sqrt{c} x^2) \sqrt{a x + b x^3 + c x^5}} - \frac{2 (b^2 - 3 a c) \sqrt{a x + b x^3 + c x^5}}{a^2 (b^2 - 4 a c) x^{3/2}} - \\
& \frac{2 c^{1/4} (b^2 - 3 a c) \sqrt{x} (\sqrt{a} + \sqrt{c} x^2) \sqrt{\frac{a + b x^2 + c x^4}{(\sqrt{a} + \sqrt{c} x^2)^2}} \text{EllipticE} \left[ 2 \text{ArcTan} \left[ \frac{c^{1/4} x}{a^{1/4}} \right], \frac{1}{4} \left( 2 - \frac{b}{\sqrt{a} \sqrt{c}} \right) \right]}{a^{7/4} (b^2 - 4 a c) \sqrt{a x + b x^3 + c x^5}} + \\
& \left( c^{1/4} (2 b^2 + \sqrt{a} b \sqrt{c} - 6 a c) \sqrt{x} (\sqrt{a} + \sqrt{c} x^2) \sqrt{\frac{a + b x^2 + c x^4}{(\sqrt{a} + \sqrt{c} x^2)^2}} \text{EllipticF} \left[ 2 \text{ArcTan} \left[ \frac{c^{1/4} x}{a^{1/4}} \right], \frac{1}{4} \left( 2 - \frac{b}{\sqrt{a} \sqrt{c}} \right) \right] \right) / \\
& (2 a^{7/4} (b^2 - 4 a c) \sqrt{a x + b x^3 + c x^5})
\end{aligned}$$

Result (type 4, 519 leaves):



$$\begin{aligned}
& - \frac{1}{2 a^2 (b^2 - 4 a c) \sqrt{\frac{c}{b + \sqrt{b^2 - 4 a c}}} \sqrt{x} \sqrt{x (a + b x^2 + c x^4)}} \\
& \left( 2 \sqrt{\frac{c}{b + \sqrt{b^2 - 4 a c}}} (-4 a^2 c + 2 b^2 x^2 (b + c x^2) + a (b^2 - 7 b c x^2 - 6 c^2 x^4)) - i (b^2 - 3 a c) (-b + \sqrt{b^2 - 4 a c}) x \right. \\
& \left. \sqrt{\frac{b + \sqrt{b^2 - 4 a c} + 2 c x^2}{b + \sqrt{b^2 - 4 a c}}} \sqrt{\frac{2 b - 2 \sqrt{b^2 - 4 a c} + 4 c x^2}{b - \sqrt{b^2 - 4 a c}}} \text{EllipticE}\left[i \text{ArcSinh}\left[\sqrt{2} \sqrt{\frac{c}{b + \sqrt{b^2 - 4 a c}}} x\right], \frac{b + \sqrt{b^2 - 4 a c}}{b - \sqrt{b^2 - 4 a c}}\right] + \right. \\
& \left. i (-b^3 + 4 a b c + b^2 \sqrt{b^2 - 4 a c} - 3 a c \sqrt{b^2 - 4 a c}) x \sqrt{\frac{b + \sqrt{b^2 - 4 a c} + 2 c x^2}{b + \sqrt{b^2 - 4 a c}}} \sqrt{\frac{2 b - 2 \sqrt{b^2 - 4 a c} + 4 c x^2}{b - \sqrt{b^2 - 4 a c}}} \right. \\
& \left. \text{EllipticF}\left[i \text{ArcSinh}\left[\sqrt{2} \sqrt{\frac{c}{b + \sqrt{b^2 - 4 a c}}} x\right], \frac{b + \sqrt{b^2 - 4 a c}}{b - \sqrt{b^2 - 4 a c}}\right] \right)
\end{aligned}$$

**Problem 122: Result more than twice size of optimal antiderivative.**

$$\int \frac{x (d + e x^2)}{\sqrt{a x + b x^3 + c x^5}} dx$$

Optimal (type 6, 287 leaves, 7 steps):

$$\begin{aligned}
& \frac{2 d x^2 \sqrt{1 + \frac{2 c x^2}{b - \sqrt{b^2 - 4 a c}}} \sqrt{1 + \frac{2 c x^2}{b + \sqrt{b^2 - 4 a c}}} \text{AppellF1}\left[\frac{3}{4}, \frac{1}{2}, \frac{1}{2}, \frac{7}{4}, -\frac{2 c x^2}{b - \sqrt{b^2 - 4 a c}}, -\frac{2 c x^2}{b + \sqrt{b^2 - 4 a c}}\right]}{3 \sqrt{a x + b x^3 + c x^5}} + \\
& \frac{2 e x^4 \sqrt{1 + \frac{2 c x^2}{b - \sqrt{b^2 - 4 a c}}} \sqrt{1 + \frac{2 c x^2}{b + \sqrt{b^2 - 4 a c}}} \text{AppellF1}\left[\frac{7}{4}, \frac{1}{2}, \frac{1}{2}, \frac{11}{4}, -\frac{2 c x^2}{b - \sqrt{b^2 - 4 a c}}, -\frac{2 c x^2}{b + \sqrt{b^2 - 4 a c}}\right]}{7 \sqrt{a x + b x^3 + c x^5}}
\end{aligned}$$

Result (type 6, 639 leaves):

$$\frac{1}{42 c (x (a + b x^2 + c x^4))^{3/2}}$$

$$a x^3 \left( b - \sqrt{b^2 - 4 a c} + 2 c x^2 \right) \left( b + \sqrt{b^2 - 4 a c} + 2 c x^2 \right) \left( - \left( \left( 49 d \operatorname{AppellF1} \left[ \frac{3}{4}, \frac{1}{2}, \frac{1}{2}, \frac{7}{4}, -\frac{2 c x^2}{b + \sqrt{b^2 - 4 a c}}, \frac{2 c x^2}{-b + \sqrt{b^2 - 4 a c}} \right] \right) / \right. \right.$$

$$\left. \left( -7 a \operatorname{AppellF1} \left[ \frac{3}{4}, \frac{1}{2}, \frac{1}{2}, \frac{7}{4}, -\frac{2 c x^2}{b + \sqrt{b^2 - 4 a c}}, \frac{2 c x^2}{-b + \sqrt{b^2 - 4 a c}} \right] + x^2 \left( \left( b + \sqrt{b^2 - 4 a c} \right) \operatorname{AppellF1} \left[ \frac{7}{4}, \frac{1}{2}, \frac{3}{2}, \frac{11}{4}, -\frac{2 c x^2}{b + \sqrt{b^2 - 4 a c}}, \frac{2 c x^2}{-b + \sqrt{b^2 - 4 a c}} \right], \right. \right.$$

$$\left. \left. \frac{2 c x^2}{-b + \sqrt{b^2 - 4 a c}} \right) + \left( b - \sqrt{b^2 - 4 a c} \right) \operatorname{AppellF1} \left[ \frac{7}{4}, \frac{3}{2}, \frac{1}{2}, \frac{11}{4}, -\frac{2 c x^2}{b + \sqrt{b^2 - 4 a c}}, \frac{2 c x^2}{-b + \sqrt{b^2 - 4 a c}} \right] \right) \right) \left. \right) -$$

$$\left( 33 e x^2 \operatorname{AppellF1} \left[ \frac{7}{4}, \frac{1}{2}, \frac{1}{2}, \frac{11}{4}, -\frac{2 c x^2}{b + \sqrt{b^2 - 4 a c}}, \frac{2 c x^2}{-b + \sqrt{b^2 - 4 a c}} \right] \right) / \left( -11 a \operatorname{AppellF1} \left[ \frac{7}{4}, \frac{1}{2}, \frac{1}{2}, \frac{11}{4}, -\frac{2 c x^2}{b + \sqrt{b^2 - 4 a c}}, \frac{2 c x^2}{-b + \sqrt{b^2 - 4 a c}} \right], \right.$$

$$\left. \frac{2 c x^2}{-b + \sqrt{b^2 - 4 a c}} \right) + x^2 \left( \left( b + \sqrt{b^2 - 4 a c} \right) \operatorname{AppellF1} \left[ \frac{11}{4}, \frac{1}{2}, \frac{3}{2}, \frac{15}{4}, -\frac{2 c x^2}{b + \sqrt{b^2 - 4 a c}}, \frac{2 c x^2}{-b + \sqrt{b^2 - 4 a c}} \right] + \right.$$

$$\left. \left( b - \sqrt{b^2 - 4 a c} \right) \operatorname{AppellF1} \left[ \frac{11}{4}, \frac{3}{2}, \frac{1}{2}, \frac{15}{4}, -\frac{2 c x^2}{b + \sqrt{b^2 - 4 a c}}, \frac{2 c x^2}{-b + \sqrt{b^2 - 4 a c}} \right] \right) \left. \right)$$

Problem 140: Unable to integrate problem.

$$\int \frac{x^{-1+\frac{q}{2}}}{\sqrt{b x^n + c x^{2n-q} + a x^q}} dx$$

Optimal (type 3, 70 leaves, 2 steps):


$$\frac{\operatorname{ArcTanh} \left[ \frac{x^{q/2} (2 a + b x^{n-q})}{2 \sqrt{a} \sqrt{b x^n + c x^{2n-q} + a x^q}} \right]}{\sqrt{a} (n - q)}$$

Result (type 8, 38 leaves):

$$\int \frac{x^{-1+\frac{q}{2}}}{\sqrt{b x^n + c x^{2n-q} + a x^q}} dx$$

## Summary of Integration Test Results

140 integration problems



A - 129 optimal antiderivatives

B - 2 more than twice size of optimal antiderivatives

C - 8 unnecessarily complex antiderivatives

D - 1 unable to integrate problems

E - 0 integration timeouts